

Safer's®

SLUG AND SNAIL BAIT II

PELLET

COMMERCIAL

READY TO USE

CONTROLS: Slugs and Snails

For use around vegetables, fruit trees, berries, ornamentals, lawns and greenhouses

EASY TO USE

ACTIVE INGREDIENT: Ferric Sodium EDTA.....6%

READ THE LABEL BEFORE USING

Warning, contains the allergen wheat

REGISTRATION NO. 28774 PEST CONTROL PRODUCTS ACT

NET CONTENTS 10 Kg

Woodstream Canada Corporation

Unit 1, 25 Bramtree Court, Brampton, ON L6S 6G2

1-800-800-1819

DIRECTIONS FOR USE:

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

WHEN TO APPLY: Early morning or evening is the best time to apply the bait, as slugs and snails travel and feed mostly by night or early morning. Re-apply as the bait is consumed or at 14-day intervals if slugs and snails continue to be a problem.

WHERE TO APPLY: All likely areas of infestation (i.e., damp places) should be treated. Also treat around the perimeter of crop areas to intercept any slugs or snails travelling into these areas from daytime refuge sites. Slugs and snails favor damp places around and within vegetable crops, in and around weeds or ditches around field margins, and leafy herbaceous perennials. Although phytotoxicity is not expected, it is recommended that a small group of test plants or small area be treated with Safer's Slug & Snail Bait II before treating an entire crop.

HOW TO APPLY: Safer's Slug & Snail Bait II is to be applied evenly on the soil surface around or near plants and areas to be protected. Do not place bait in piles. The soil should be moist, but with little or no standing water. Product can be applied using ground equipment such as a handheld spreader or rotary type spreader. Use the higher rates to control slugs and snails at higher levels of pest pressure.

APPLICATION RATES

Crops/Use	Rate	Directions
Vegetables	11-22 kg/ha	Evenly scatter the bait around the perimeter of vegetables and between rows where slugs or snails are observed.
Orchard fruits	11-22 kg/ha	For mature trees in orchards, evenly scatter bait evenly around the base of the trees where slugs or snails are observed. For new seedlings, spread bait evenly in a 15 cm band around the base of the stem where slugs or snails are observed.
Berries	11-22 kg/ha	Evenly scatter by hand or with granular spreaders around the perimeter of plot and between rows where slugs and snails are observed. For small plots, evenly apply the bait around the base of the plants to be protected from slugs and snails. Do not broadcast entire area, but treat selectively.
Field crops	11-22 kg/ha	At all stages of growth, evenly apply the bait between the rows and around the field perimeter where slugs or snails are observed.
Vineyards	11-22 kg/ha	Evenly scatter bait at the base of vines and between rows where slugs or snails are observed.
Wheat	11-22 kg/ha	Evenly scatter bait with a granular spreader in areas where slugs or snails are observed.
Grass grown for seed production	11-22 kg/ha	Evenly scatter bait with a granular spreader in areas where slugs or snails are observed.

(turf and feed crops)		
Greenhouse vegetables	2 g/m ² or 1 g/ 10 pots measuring 23 cm diameters	Evenly scatter bait on the soil surface of potted plants being damaged by slugs or snails. Bait can also be evenly scattered around pots on greenhouse benches.
Greenhouse ornamentals	2 g/m ² or 1 g/ 10 pots measuring 23 cm diameters	For large areas, evenly scatter bait on the soil surface of potted plants being damaged by slugs or snails. Bait may also be evenly scattered around pots on greenhouse benches.
Outdoor ornamentals	2 g/m ²	Evenly scatter bait in a 15 cm wide circular band around the base of the plants to be protected where slugs or snails are observed.
Outdoor container-grown nursery plants	2 g/m ² or 3 g/ 10 pots measuring 46 cm diameters	Evenly scatter bait on the soil surface of containers where slugs or snails are observed. Bait may also be evenly scattered on the soil surface around pots.
Turf, golf courses, sod farms	2 g/m ²	Evenly scatter bait with a granular spreader in areas where slugs or snails are observed.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.

May cause eye irritation. Avoid contact with skin, eyes, and clothing. Wear waterproof gloves and goggles during application, or during mixing, loading, and repair/clean-up activities. Avoid hand-to-mouth contact. Avoid use in areas frequented by pets. Wash hands with soap and water after handling.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at:

www.croplife.ca

Do not apply by air.

FIRST AID: If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. Take the affected person to a safe and uncontaminated area. Seek medical attention IMMEDIATELY if redness, swelling, itching, burning or shortness of breath occurs. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

ENVIRONMENTAL HAZARDS: Do not apply directly to water.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Dispose of the rinsings in accordance with provincial requirements.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product contact the manufacturer or provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offense under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.