

(Sleeve)

GROUP	4	9	HERBICIDES
--------------	----------	----------	-------------------

NUFARM

GlyKamba® Herbicide

For use in pre-seeding applications in cereals and summerfallow
Solution

AGRICULTURAL

READ THE LABEL AND BOOKLET BEFORE USING.

ACTIVE INGREDIENT:

Glyphosate, present as isopropylamine salt ... 194 g a.e./L
Dicamba, present as isopropylamine salt ... 46 g a.e./L

REGISTRATION NO. 30870

PEST CONTROL PRODUCTS ACT.

CAUTION POISON
DANGER CORROSIVE TO EYES

AVOID CONTACT WITH FOLIAGE OF CROP OF OTHER DESIRABLE VEGETATION SINCE SEVERE INJURY OR DESTRUCTION MAY
RESULT

Net Contents: 1 L to Bulk

Nufarm Agriculture Inc.
5101, 333 - 96th Ave NE
Calgary, Alberta T3K 0S3
1-800-868-5444

24 Hour Emergency Response Number
1-800-424-9300

GlyKamba® is a registered trademark of Nufarm Agriculture Inc.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. May cause irreversible damage to eyes. May cause allergic skin reaction. Harmful if inhaled or swallowed. Avoid inhaling dust, sprays, etc. Do not get in eyes, on skin or on clothing.

DO NOT enter treated fields until 12 hours after application.

DO NOT permit lactating dairy animals to graze fields within 7 days after application.

DO NOT harvest forage or cut hay within 30 days after application. Withdraw meat animals from treated fields at least 3 days before slaughter.

DO NOT use in residential areas, which are defined as sites where bystanders may be present during or after spraying, including homes, schools, parks, playgrounds, playing fields and public building.

DO NOT apply by air.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings

PROTECTIVE CLOTHING AND EQUIPMENT:

Wear a long-sleeved shirt, long pants, goggles or a face shield and chemical-resistant gloves during mixing, loading, clean-up or repair activities. Applicators must wear a long-sleeved shirt, long pants and chemical-resistant gloves.

Wash thoroughly with soap and water after handling.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Dicamba may cause severe irritation to the eyes, and irritation to the skin and mucous membranes. Symptoms of overexposure to dicamba may include dizziness, muscle weakness, loss of appetite, weight loss, vomiting, decreased heart rate, shortness of breath, excitement, tenseness, depression, incontinence, cyanosis, muscle spasms, exhaustion and loss of voice. Treat symptomatically.

EMERGENCY NUMBERS

For spills or transportation accidents, Chemtrec, 1-800-424-9300.

For health or environmental emergencies, ProPharma Group, 1-877-325-1840.

For product and use information, Nufarm Agriculture Inc., 1-800-868-5444.

PHYSICAL OR CHEMICAL PRECAUTIONS:

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fibreglass, plastic and plastic lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or

explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ENVIRONMENTAL HAZARDS:

- TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.
- Avoid direct applications to any body of water. Do not use in areas where adverse impact on domestic water or aquatic species is likely. Do not contaminate water by disposal of waste or cleaning of equipment.
- Avoid all drift to or contact with other vegetation for which treatment is not intended as damage or destruction may occur.
- DO NOT apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands), estuarine or marine habitats.
- DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Surface Runoff

- To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include but are not limited to heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted, fine textured, or low in organic matter such as clay).
- Potential for contamination of aquatic areas as a result of runoff may be reduced by including an untreated vegetative strip between the treated area and the edge of the water body.
- Avoid applying this product when heavy rain is forecast.

Leaching

- The use of this chemical may result in contamination of groundwater, particularly in areas where soils are permeable (e.g. sand, loamy sand and sandy loam soils) and/or the depth to the water table is shallow.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Store above 5 °C to keep product in solution. If crystals form, place in a warm room (20 °C), allow the product to reach room temperature and roll or shake periodically until crystals have re-dissolved. Keep container closed to prevent spills and contamination.

DISPOSAL:

For Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site.

Contact your local distributor/dealer or municipality for the location of the nearest collection site.

Before taking the container to the collection site:

- Triple- or pressure-rinse the empty container. Add the rinsing to the spray mixture in the tank.
- Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Returnable-Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the

distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.

(booklet)

GROUP	4	9	HERBICIDES
--------------	----------	----------	-------------------

NUFARM

GlyKamba® Herbicide

For use in pre-seeding applications in cereals and summerfallow

Solution

AGRICULTURAL

READ THE LABEL AND BOOKLET BEFORE USING.

ACTIVE INGREDIENT:

Glyphosate, present as isopropylamine salt ... 194 g a.e./L
Dicamba, present as isopropylamine salt ... 46 g a.e./L

REGISTRATION NO. 30870 PEST CONTROL PRODUCTS ACT.

**CAUTION POISON
DANGER CORROSIVE TO EYES**

AVOID CONTACT WITH FOLIAGE OF CROP OF OTHER DESIRABLE VEGETATION SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT

Net Contents: 1 L to Bulk

Nufarm Agriculture Inc.
5101, 333 - 96th Ave NE
Calgary, Alberta T3K 0S3
1-800-868-5444

24 Hour Emergency Response Number
1-800-424-9300

GlyKamba® is a registered trademark of Nufarm Agriculture Inc.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. May cause irreversible damage to eyes. May cause allergic skin reaction. Harmful if inhaled or swallowed. Avoid inhaling dust, sprays, etc. Do not get in eyes, on skin or on clothing.

DO NOT enter treated fields until 12 hours after application.

DO NOT permit lactating dairy animals to graze fields within 7 days after application.

DO NOT harvest forage or cut hay within 30 days after application. Withdraw meat animals from treated fields at least 3 days before slaughter.

DO NOT use in residential areas, which are defined as sites where bystanders may be present during or after spraying, including homes, schools, parks, playgrounds, playing fields and public building.

DO NOT apply by air.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings

PROTECTIVE CLOTHING AND EQUIPMENT:

Wear a long-sleeved shirt, long pants, goggles or a face shield and chemical-resistant gloves during mixing, loading, clean-up or repair activities. Applicators must wear a long-sleeved shirt, long pants and chemical-resistant gloves.

Wash thoroughly with soap and water after handling.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Dicamba may cause severe irritation to the eyes, and irritation to the skin and mucous membranes. Symptoms of overexposure to dicamba may include dizziness, muscle weakness, loss of appetite, weight loss, vomiting, decreased heart rate, shortness of breath, excitement, tenseness, depression, incontinence, cyanosis, muscle spasms, exhaustion and loss of voice. Treat symptomatically.

EMERGENCY NUMBERS

For spills or transportation accidents, Chemtrec, 1-800-424-9300.

For health or environmental emergencies, ProPharma Group, 1-877-325-1840.

For product and use information, Nufarm Agriculture Inc., 1-800-868-5444.

PHYSICAL OR CHEMICAL PRECAUTIONS:

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fibreglass, plastic and plastic lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or

explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Store above 5 °C to keep product in solution. If crystals form, place in a warm room (20 °C), allow the product to reach room temperature and roll or shake periodically until crystals have re-dissolved. Keep container closed to prevent spills and contamination.

DISPOSAL:

For Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site.

Contact your local distributor/dealer or municipality for the location of the nearest collection site.

Before taking the container to the collection site:

- Triple- or pressure-rinse the empty container. Add the rinsing to the spray mixture in the tank.
- Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Returnable-Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.

1. GENERAL INFORMATION

PRODUCT INFORMATION:

- **Glyphosate** is not to be applied using hand-wicking or hand-daubing methods.
- **GlyKamba Herbicide** is a post-emergent herbicide for control or suppression of emerged weeds prior to seeding cereals in fallow and reduced tillage systems.
- This product enters the plant through the foliage and moves throughout the plant. Visual effects of control are a gradual wilting or yellowing of the plant, which advances to complete browning of above-ground growth and deterioration of affected underground plant parts. Visible symptoms will usually develop on labelled weeds within 5 to 7 days after applications but may not occur for more than 7 days. Extremely cool or cloudy weather following treatment or prolonged drought conditions may slow activity of this product and delay the visual effects of control.
- Always use the higher rate of this product under poor growing conditions, such as drought.
- Reduced control may result if treatments are made during poor growing conditions such as drought stress, disease or insect damage, or if weeds have been mowed, grazed or cut. Heavy dust on foliage or an overstory canopy covering targeted weeds may also reduce control.
- This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the precautionary statements and all other information appearing on the labels of all herbicides used.

- Rainfall occurring with 6 hours after application, particularly on weeds growing under stress conditions, may reduce the effectiveness of this product. Heavy rainfall within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required. For best results spray coverage should be uniform and complete.

-

Restricted Entry Intervals

The restricted entry interval is 12 hours after application for all agricultural uses.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, please note that **GlyKamba Herbicide** is both a Group 4 and Group 9 herbicide. Any weed population may contain plants naturally resistant **GlyKamba Herbicide** and other Group 4 and Group 9 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **GlyKamba Herbicide** or other Group 4 and 9 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information and to report suspected resistance, contact Nufarm Agriculture Inc. at 1-800-868-5444 or at www.nufarm.ca.

ENVIRONMENTAL HAZARDS:

- TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.
- Avoid direct applications to any body of water. Do not use in areas where adverse impact on domestic water or aquatic species is likely. Do not contaminate water by disposal of waste or cleaning of equipment.
- Avoid all drift to or contact with other vegetation for which treatment is not intended as damage or destruction may occur.
- DO NOT apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands), estuarine or marine habitats.

- DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Surface Runoff

- To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include but are not limited to heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted, fine textured, or low in organic matter such as clay).
- Potential for contamination of aquatic areas as a result of runoff may be reduced by including an untreated vegetative strip between the treated area and the edge of the water body.
- Avoid applying this product when heavy rain is forecast.

Leaching

- The use of this chemical may result in contamination of groundwater, particularly in areas where soils are permeable (e.g. sand, loamy sand and sandy loam soils) and/or the depth to the water table is shallow.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. Boom height must be 60 cm or less above the crop or ground.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURING DESIRABLE PLANTS AND CROPS. Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings. Do not allow spray mist to drift since even minute quantities of spray can cause severe damage or destruction to nearby crops, plants or other areas on which treatment is not intended, or may cause other unintended consequences. Do not apply when winds are gusty or in excess of 8 km/h or when other conditions, including lesser wind velocities, will allow drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in fine particles (mist) which are more likely to drift.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

BUFFER ZONES:

Use of the following spray methods or equipment **DO NOT** require a buffer zone: hand-held or backpack sprayer and spot treatment, inter-row hooded sprayer, low-clearance hooded or shielded sprayers that ensure spray drift does not come in contact with orchard crop fruit or foliage, soil drench and soil incorporation.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

Table 1 Buffer Zones for the Protection of Aquatic and Terrestrial Habitats from Spray Drift of Glyphosate Products Formulated with POEA

Agricultural and non-cropland systems	Maximum number of applications	Buffer Zones (metres) Required for the Protection of:	
		Aquatic habitats	Terrestrial habitats
Agricultural crop system and ground boom application method			
Pre-seeding applications for rye, summer fallow	1	1	1
Canola – Roundup Ready hybrid for seed production	2	1	1
Corn (glyphosate non-tolerant varieties including grain, silage and ornamental types)	2	1	2
Wheat, barley, oats, soybean (glyphosate non-tolerant varieties), corn-sweet (glyphosate tolerant varieties), canola (glyphosate non-tolerant varieties), flax (including low linoleic acid varieties), pearl millet, corn (glyphosate tolerant varieties)	3	1	2
Canola (glyphosate tolerant varieties), soybean (glyphosate tolerant varieties)	4	1	2

* Buffer zones for the protection of terrestrial habitats are not required for use on rights-of-way including railroad ballast, rail and hydro rights-of-way, utility easements and roads.

NR = Not Required.

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

2. DIRECTIONS FOR USE

- **As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests**
- **DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.**

Apply this product at 2.5 to 3.125 L/ha in 50 to 100 litres of clean water per hectare.

The higher rate of this product should be applied when weeds are under poor growing conditions such as drought or for those weeds specified in the table below.

- Recommended applications of this product can be made as required prior to seeding in cereal crops and to fallow land.
- This product or labelled tank mixtures should be applied to emerged, actively growing weeds.

- Application at earlier growth stages generally will provide best results.
- Reduced results may occur if application of this product or labelled tank mixtures is made to weeds which are not actively growing or weeds that are drought stressed.

MIXING INSTRUCTIONS:

1. Fill the spray tank to about ¾ of the desired volume with clean water.
2. Add the recommended amount of **GlyKamba Herbicide**.
3. Complete the filling process while maintaining agitation.

Remove the hose from the mix tank immediately after filling to avoid siphoning back into the carrier source. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, terminate by-pass and return lines at the tank bottom during mixing.

Reduced results may occur if water containing soil is used, such as water from ponds and unlined ditches.

SPRAYER CLEANUP

CLEAN THE ENTIRE SPRAYER AFTER APPLICATION OF THIS PRODUCT. Failure to clean the sprayer thoroughly may result in injury to desirable crops which are subsequently sprayed. First, add clean water to the tank and thoroughly rinse the entire sprayer system.

Secondly, fill the tank with water and ammonia. Add 1 litre of household ammonia per 100 litres of water. Pump enough solution through the hoses, boom and nozzles to fill these parts completely. Then fill the tank, close and leave for 24 hours before draining and rinsing thoroughly with water.

Application or use of other agricultural chemicals with the equipment used for this product may result in injury to desirable vegetation.

3. WEEDS CONTROLLED WITH GlyKamba Herbicide

Annual Grasses

Weeds controlled	L/ha	Application timing
Brome (Downy)* Cereals (volunteer) Darnel (Persian) Foxtail (Green) Oats (Wild)	2.5	Any time between emergence and heading

Annual Broadleaves

Weeds controlled	L/ha	Application timing
Buckwheat, wild	2.5	1 to 4 leaf stage
Canola (volunteer) - does not control volunteer Roundup Ready Canola Cowcockle Flixweed* Kochia Ladysthumb Lambsquarter Mustard, wild Pigweed (Redroot) Smartweed Stinkweed Thistle, Russian	2.5	Up to 15 cm height

Perennial

Weeds controlled	L/ha	Application timing
Barley (Foxtail)**	3.125	Before initiation of seed head or browning of lower leaves

* See "Special Weed Control Concerns" Section of label

** Suppression only

SPECIAL WEED CONTROL CONCERNS

Annual Weed Control Prior to Seeding

Crops: prior to seeding wheat, barley, rye, oats and field corn (do not apply prior to sweet corn planting)

Rate: 2.5 L/ha

Notes:

- Use on emerged annual weeds in reduced tillage systems
- Do not apply prior to seeding lentils, peas, canola, flax or other broadleaved crops.
- Do not apply after crop emergence.
- Seeding should follow soon after application since this product does not provide residual weed control.
- For field corn, apply to medium to fine textured soils containing more than 2.5% organic matter.
- Do not use on sandy or sandy loam soils.
- Use only the 2.5 L/ha rate for pre-seeding applications.

Winter Annual Weeds

For best control of winter annual broadleaf weeds, such as flixweed and stinkweed, 2,4-D should be applied to emerged, actively growing weeds in the fall previous to the fallow season or in early spring in the fallow season when winter annual weeds are less than 10 cm tall. Refer to the 2,4-D product label for appropriate rates and other use instructions and cautionary statements.

Downy Brome

For best control of these winter annual weeds, **GlyKamba Herbicide** can be applied after emergence in the fall previous to the fallow season or in spring of the fallow season up to seed head emergence.

Control of Weeds under Stress Conditions

Under certain stress conditions, such as drought, cool temperatures or where extremely hard water (> 700 ppm Ca + Mg) has been used, weed control may be reduced with this product.

However, the following list of application instructions will help improve results under these conditions.

- Use 50 L/ha water volume
- Use the higher rate of **GlyKamba Herbicide** within the recommended rate range (3.125 L/ha)

4. TANK MIXTURES WITH GlyKamba Herbicide

When applied as a tank-mix combination, read and observe all label directions, including rates, restrictions, and grazing limitations for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Nufarm Agriculture Inc. at 1-800- 868-5444 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do

not appear on this label or that are not specifically recommended by Nufarm Agriculture Inc.

GlyKamba Herbicide + 2,4-D

For control of various broadleaf weeds (i.e. volunteer glyphosate tolerant canola)

Crops: Prior to seeding and/or emergence of wheat (winter, spring), barley, rye in minimum tillage systems, summerfallow

Application Timing: summerfallow; prior to seeding or after seeding prior to crop emergence

Rate: Use only the 2.5 L/ha rate of GlyKamba Herbicide prior to cereal crop seeding and/or emergence.

Use 2.5 – 3.125 L/ha of GlyKamba Herbicide for a summerfallow application.

Water Volume: 50 to 100 L/ha

Notes: GlyKamba Herbicide may be tank mixed with 2,4-D low volatile ester or amine formulations.

Refer to the 2,4-D herbicide label for further safety precautions and handling instructions.

Product	Rate (L/ha)	Weeds Controlled		Application Stage
GlyKamba Herbicide + 2,4-D ^a	2.5 - 3.125 + 0.6 – 0.9 L/ha (280 – 420 g ai/ha)	Brome (Downy), Cereals (volunteer) Darnel (Persian)	Foxtail (Green) Oats (Wild) Barley (Foxtail)*	Any time between emergence and heading. *Use 3.125 L/ha before initiation of seed head or lower leaves browning. Suppression only.
		Buckwheat, wild** Canola, volunteer (not Roundup Ready canola), Cowcockle, Flixweed, Kochia,	Lady’s thumb, Lamb’s quarters, Mustard, wild Pigweed (Redroot) Smartweed, Stinkweed, Thistle, Russian	Up to 15 cm tall and actively growing for best results. ** 1-4 leaf stage use 2.5 L/ha
		Bluebur, Burdock (before the 4 leaf), Canola, Volunteer Roundup Ready (1-4 leaf stage), Cocklebur, Common plantain, Daisy fleabane, False flax, False ragweed, Goat’s beard,	Mustards (except dog and tansy), Prickly lettuce, Ragweed (common & giant), Russian pigweed, Shepherd’s purse, Stinging nettle, Sweet clover, Thyme-leaved spurge, Wild radish, Wild sunflower	2-4 leaf stage unless noted.
GlyKamba Herbicide + 2,4-D ^a	2.5 - 3.125	Weed listed above plus: Volunteer Roundup Ready canola (4-6 leaf stage), Annual sow thistle, Blue lettuce, Biennial wormwood,	Gumweed, Hairy galinsoga, Hawkweed, Heal-all, Hedge bindweed, Knotweed (before 4 leaf), Leafy spurge, Peppergrass,	2-4 leaf stage unless noted

	+ 1.2 – 1.5 L/ha (560 – 700 g ai/ha)	Common chickweed, Common purslane, Curled dock (before 4 leaf), Dog and tansy mustard, Oak-leaved goosefoot, Groundsel,	Pineapple weed, Prostrate pigweed, Purslane, Russian knapweed, Sheep sorrel, Smartweed, Tumble pigweed, Velvetleaf, Volunteer mustard, Yellow rocket (before 4 leaf)	Top growth control when applied after the 4 leaf stage
		Blue lettuce, Biennial wormwood, Gumweed, Hedge bindweed, Leafy spurge, Russian knapweed, Yellow rocket (before 4 leaf), Bull thistle, Burdock buttercup, Canada thistle,	Field bindweed, Field dandelion, Hoary cress, Mouse-eared chickweed, Perennial sow thistle, Tartary buckwheat, Teasel, Volunteer flax, Volunteer sunflower	

^a Based on 470 g a.i./litre 2,4-D. Adjust rates accordingly for other 2,4-D formulations. Use only low volatile ester or amine formulations of 2,4-D.

5. GlyKamba in Glyphosate Tolerant Corn (i.e., varieties with genetic resistance to glyphosate) – Post-emergent application

NOTE: ALWAYS USE PEDIGREED (CERTIFIED) CORN SEED DESIGNATED AS GLYPHOSATE TOLERANT. CORN WHICH IS NOT DESIGNATED AS GLYPHOSATE TOLERANT WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.

Do not apply by air

Do not apply more than one application per year

Product	Rate (L/ha)	Weeds Controlled	Application Stage
GlyKamba Herbicide	2.5	All Weeds listed in Section 3 at the 2.5 L/Ha rate	Glyphosate tolerant corn, 1 to 5-leaf stage

GlyKamba® is a Registered Trademark of Nufarm Agriculture Inc.