

2013-06-25

2012-4592

ZODIAC®

**POWERBAND PLUS
DUAL ACTION FLEA AND TICK COLLAR
FOR CATS & KITTENS**

DOMESTIC

GUARANTEE

Propoxur....10.0%

(S)-Methoprene.....2.10%

REG. NO. 28199 P.C.P. ACT

**KEEP OUT OF REACH OF CHILDREN
READ THE LABEL BEFORE USING
FOR USE ONLY ONLY ON CATS**

NET CONTENTS: 10.5g

WELLMARK INTERNATIONAL

100 Stone Road West, Suite 111

Guelph, Ontario N1G 5L3

(PRIMARY or SECONDARY PANEL)

For use in the control of cat fleas, dog fleas, American dog ticks and brown dog ticks on cats.

- *With Precor® [vIGRen®] Insect Growth Regulator [IGR]*
- *New Technology*
- *Plus IGR Kills Flea Eggs*
- *Kills adult fleas fast*
- *245 days [35 weeks] [8 months] protection against adult fleas*
- *245 days [35 weeks] [8 months] protection against flea eggs*
- *60 days [8 weeks] [2 months] protection against ticks*
- *Kills Adult Fleas and Ticks*
- *Kills Flea Eggs for 8 Months*
- *Contains Precor® [vIGRen®], an Insect Growth Regulator, which Kills Flea Eggs Laid on Your Pet for 8 Months*
- *Long Lasting Protection*
- *Works from head to tail*
- *Quick acting [action] [killing] for [of] adult fleas*
- *Kills adult fleas fast*
- *Ovicidal – kills flea eggs [before they hatch] [245 days] [35 weeks] [8 months]*
- *Unique one-piece buckle [integrated one-piece buckle]*
- *Integrated plastic buckle [Metal Buckle] [Expandable Collar][Break Away Feature][Stretch Band]*
- *Sold Through Veterinarians*

NOTE: Italics and [Bracketed] Tex = Alternate/Optional Text

This product will begin to kill fleas and ticks within hours. Control of ticks will take a few days to occur after placing the collar on the pet. The collar will continue to control fleas for 8 months by killing adult fleas and flea eggs on your pet. This product may be worn with a regular collar.

DIRECTIONS: Do NOT remove the collar from its package until ready to use. Place the collar around the cat's neck, adjust for proper fit and buckle in place *[by threading flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle.]* *[Since the plastic buckle is molded into the collar, the possibility of neck irritation and buckle failure is reduced.]* The collar must be worn loosely to allow for growth of the animal and to permit the collar to move around the neck. *[A properly fitted collar is one that, when fastened, will snugly slide over the cat's head.]* *[Leave 5 to 9 cm on the collar for extra adjustment. Cut off and dispose of extra length in empty pouch and discard in garbage. To release the collar, squeeze buckle between thumb and finger and pull sharply.]*

Do NOT use on sick or nursing animals, or on cats under 12 weeks of age or animals receiving drug or other pesticide treatment. Consult a veterinarian before using on aged or pregnant animals. Replace this collar no more than once every 8 months. Replace collar every 8 months for continual pest control in situations where there is an ongoing risk of pest infestation.

RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE. Treatment must only be repeated if pest problem reoccurs.

For effective flea and tick control, treatment of the pet should be combined with sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, an insecticidal treatment of the premises may be required (Zodiac Premise Plus, Zodiac Premise 1000/2000). A veterinarian should be consulted when pest infestation continues to be a problem on the pet.

PRECAUTIONS: HUMANS: KEEP OUT OF REACH OF CHILDREN.

DO NOT ALLOW CHILDREN TO HANDLE THIS COLLAR. Harmful if collar is sucked or chewed. Dust will form on this collar during storage. Dust may irritate eyes, nose and throat. Harmful if dust is inhaled or absorbed through skin. Avoid contact with mouth, eyes, skin and clothing. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Avoid product contact with food, food utensils or places where food is prepared or stored.

PETS: Harmful if collar is chewed or swallowed. Do NOT use this product at the same time or within 30 days before or after treatment with, or exposure to, other cholinesterase-inhibiting drugs, pesticides or chemicals. Dust may irritate eyes, nose and throat. Avoid letting pet inhale dust from collar. Avoid contact with cat's mouth, face, eyes and genitals. Some animals may be sensitive to this collar. Any collar, when fastened too tightly, may cause skin irritation. If irritation persists after proper adjustment, remove the collar and consult a veterinarian. Monitor your pet closely for the first few days in order to ensure there is no adverse reaction.

FIRST AID:

HUMANS: IF SWALLOWED, SUCKED OR CHEWED, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. IF DUST CONTACTS SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. IF DUST CONTACTS EYES, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove

contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. IF DUST FROM COLLAR IS INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. Take container, label or product name and PCP Registration Number with you when seeking medical or veterinary attention.

PETS:

Sensitivities may occur after using ANY pesticide for pets. If signs of sensitivity occur, remove collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. If swallowed or chewed, consult a veterinarian. Take container, label or product name and PCP Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION:

This product contains a pesticide that is a cholinesterase inhibitor (an anti-cholinesterase compound). Atropine is antidotal if symptoms of cholinesterase inhibition are present.

STORAGE: Store in original, unopened container at room temperature.

DISPOSAL: Do not reuse empty pouch. Dispose of any excess collar, packaging and unwanted (used) collars in household garbage.

100% satisfaction guaranteed.

Visit our website: www.zodiacpet.com

HOT LINE: 1-800-688-7378

Zodiac is a registered trademark of Wellmark International

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.