

2015-5424

2016-01-14

(Box Label)

Pro-Shield 10 Flea Adulticide

For dogs 12 weeks and older weighing 4.5 kg and under
[Once-A-Month] [Once Per Month] Topical Flea Treatment

LIQUID

[Rapidly kills fleas] [Kills fleas rapidly] [and] [may reduce flea allergy dermatitis (FAD)]
[Controls fleas within one (1) day of treatment]
[Flea infestation protection for [at least][a minimum of] four (4) weeks]
[Three] [Four] [Six] [3] [4] [6] months' supply

READ THE LABEL AND INSERT BEFORE USING

FOR USE ONLY ON DOGS

Do not use on other animals

KEEP OUT OF REACH OF CHILDREN

WARNING

POISON

EYE IRRITANT

DOMESTIC

GUARANTEE: Imidacloprid 9.1%

REGISTRATION NO. 32016 P.C.P. ACT

[Three] [Four] [Six] [x] 0.4 mL Tubes
TWIST N' USE TUBE™

NorVet Supplies Inc.
2325 Hurontario St., Suite 196
Mississauga, ON L5A 4KA
(905) 274-0009

Pro-Shield 10 Flea Adulticide will control fleas on dogs within one day of treatment and will continue to protect the dog from further flea infestation for at least four weeks. Reinfestation from the emergence of new fleas in the environment may continue to occur for six weeks or longer after treatment is initiated. Reapply every 4 weeks for continual pest control in situations where there is an ongoing risk of pest infestation. Treatment must only be repeated if pest problem reoccurs. If retreatment becomes necessary earlier than 4 weeks, do not retreat more than once weekly.

Shampooing of the dog may shorten the duration of flea protection, therefore, re-application following shampooing may be necessary.

For use on pregnant/lactating animals consult your veterinarian.

PRECAUTIONS:

HUMANS: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed, or absorbed through the skin. Causes eye irritation. **DO NOT** get in eyes. Avoid contact with skin and clothing. Wear rubber gloves when applying product. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Avoid contact with pets until dry. Do not contaminate water, feed or food by storage or disposal.

PETS: Do not use on animals under 12 weeks of age. Care should be taken to prevent exposure of other pets who may be in close physical contact with the treated animal. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. Smaller animals are more likely to experience an adverse reaction. Observe carefully. If signs persist, or become more severe, consult a veterinarian immediately.

FIRST AID:

HUMANS: IF IN EYES, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

STORAGE: Store container at room temperature.

DISPOSAL: Do not reuse empty container. Wrap empty tube and cap in newspaper and dispose with household garbage.

(Insert label)

Pro-Shield 10 Flea Adulticide

For dogs 12 weeks and older weighing 4.5 kg and under
Once-A-Month [Once Per Month] Topical Flea Treatment

[Rapidly kills fleas] [Kills fleas rapidly] [and] [may reduce flea allergy dermatitis (FAD)]
[Controls fleas within one (1) day of treatment]
[Flea infestation protection for [at least][a minimum of] fo ur (4) weeks]

[Three] [Four] [Six] [3] [4] [6] months' supply
LIQUID

READ THE LABEL AND INSERT BEFORE USING

FOR USE ONLY ON DOGS

Do not use on other animals

KEEP OUT OF REACH OF CHILDREN

WARNING

POISON

EYE IRRITANT

DOMESTIC

GUARANTEE: Imidacloprid 9.1%

REG. NO. 32016 P.C.P. ACT
[Three] [Four] [Six] 0.4 mL Tubes

Pro-Shield 10 Flea Adulticide will control fleas on dogs within one day of treatment and will continue to protect the dog from further flea infestation for at least four weeks. Reinfestation from the emergence of new fleas in the environment may continue to occur for six weeks or longer after treatment is initiated. Reapply every 4 weeks for continual pest control in situations where there is an ongoing risk of pest infestation. Treatment must only be repeated if pest problem reoccurs. If retreatment becomes necessary earlier than 4 weeks, do not retreat more than once weekly.

Shampooing of the dog may shorten the duration of flea protection, therefore, re-application following shampooing may be necessary.

For effective flea control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist an insecticidal premise treatment may be required.

The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with Pro-Shield 10 Flea Adulticide rapidly kills fleas and may reduce the incidence of this condition.

Pro-Shield 10 Flea Adulticide is safe to use concurrently with commonly used veterinary products. Consult your veterinarian before using on animals receiving drug or other pesticide treatment. For use on pregnant/lactating animals consult your veterinarian.

PRECAUTIONS:

HUMANS: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed, or absorbed through the skin. Causes eye irritation. DO NOT get in eyes. Avoid contact with skin and clothing. Wear rubber gloves when applying product. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Avoid contact with pets until dry. Do not contaminate water, feed or food by storage or disposal.

PETS: Do not use on animals under 12 weeks of age. Care should be taken to prevent exposure of other pets who may be in close physical contact with the treated animal. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. Smaller animals are more likely to experience an adverse reaction. Observe carefully. If signs persist, or become more severe, consult a veterinarian immediately.

FIRST AID:

HUMANS: IF IN EYES, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

STORAGE: Store container at room temperature.

DISPOSAL: Do not reuse empty container. Wrap empty tube and cap in newspaper and dispose with household garbage.

DIRECTIONS FOR USE:

Do NOT administer by mouth. Do NOT use on other animals. Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment. A veterinarian should be consulted when pest infestation continues to be a problem on the pet.

Treatment must only be repeated if pest problem reoccurs.

Do not use on puppies under 12 weeks of age.

Use only on dogs weighing 4.5 kg and under.

This product is packaged in single dosage tubes. This package contains [Three] [Four] [Six] [3] [4] [6] monthly 0.4 mL dosages. Treat your dog monthly in the manner described below. Do not apply a dosage higher than one 0.4 mL tube at a time.

Remove one applicator from the package.

Hold applicator tube in upright position.

Twist tip to break seal. The dog should be standing for easy application. Part the hair on the dog's back between the shoulder blades until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to completely empty the contents directly on the skin. Do not get this product in your pet's eyes or mouth.

Until Pro-Shield 10 Flea Adulticide dries, keep pet off furniture and away from hardwood floors.

Discard the empty tube as described in "DISPOSAL".

RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE.

NorVet Supplies Inc.

2325 Hurontario St., Suite 196

Mississauga, ON L5A 4KA

(905) 274-0009

(Tube label)

Pro-Shield 10 Flea Adulticide

READ [THE] LABEL [&] [AND] INSERT BEFORE USING

WARNING

POISON

EYE IRRITANT

DOMESTIC

EXTERNAL USE ONLY

GUARANTEE: IMIDACLOPRID 9.1%
REG NO 32016 PCP ACT

[NorVet Supplies Inc.
2325 Hurontario St., Suite 196
Mississauga, ON L5A 4KA]
[NorVet Supplies Inc.]
(905) 274-0009
0.4 mL

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.