

PMRA Approved Label  
2015-3956  
2015-08-25

# **CUEVA<sup>®</sup> RTU WITH QUICK CONNECT<sup>™</sup> SPRAYER**

COPPER FUNGICIDE

DOMESTIC

SOLUTION

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: Copper, present as copper octanoate and related copper compounds.....0.0178%

REG. NO. 31833 PCP ACT

NET CONTENTS 1 L - 6 L

W. Neudorff GmbH KG, An der Mühle 3,  
31860 Emmerthal, Germany

**Canadian Agent:**  
Neudorff North America  
PO Box 178  
Brentwood Bay, BC V8M 1R3  
250-652-5888

## PMRA Approved Label

2015-3956

2015-08-25

Dormant and growing season liquid copper fungicide. Controls many plant diseases using low concentrations of copper. Biodegradable - Decomposes to a form useful to plants & microbes. Do not apply within one day of harvest.

### DIRECTIONS FOR USE

DO NOT apply this product in a way that will contact adults, children or pets, either directly or through drift.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT apply to any body of water. Avoid application of this product when winds are gusty.

### USE AND STORAGE DIRECTIONS FOR THE QUICK CONNECT™ SPRAYER:

#### [ILLUSTRATIONS]

1. Pull trigger sprayer from bottle.
2. Pull hose connector and coil from trigger sprayer handle.
3. Insert hose connector into spout on cap until it "clicks" in place.
4. Flip up spout.
5. Adjust spray nozzle to deliver a wide spray for best coverage.
6. Aim sprayer nozzle at plants and spray to wet.
7. Storage: After use, flip spout down to the "flat" position to close. No need to disconnect trigger sprayer. Close nozzle on trigger sprayer. Snap trigger sprayer back into place on bottle.

### PESTS/SITES AND APPLICATION INSTRUCTIONS

GENERAL USE: For outdoor and greenhouse use only. Unless otherwise stated below, the standard application rate for the listed species is 1 L of CUEVA® RTU WITH QUICK CONNECT SPRAYER per 10 m<sup>2</sup>. Spray all plant surfaces thoroughly; if necessary, repeat sprays every 7 to 10 days unless otherwise noted below. For best control, begin treatment 2 weeks before disease normally appears or when weather forecasts predict a long period of wet weather. Alternatively, begin treatment when disease first appears and repeat at 7 to 10 day intervals unless otherwise noted below. For outdoor flowers and ornamentals, do not exceed 10 applications per year. If possible, time applications so that 12 hours of dry weather follow application.

CUEVA® RTU WITH QUICK CONNECT SPRAYER may cause some copper toxicity on some plant species. During excessive moisture and cold, leaf spots can appear on copper-sensitive crop species. If concerned about sensitivity of plants, apply to individual plants or small areas of plants and wait to determine if plant damage occurs prior to treatment of larger areas.

TURF		
Plant	Application Rate and Notes	Diseases Controlled
Turf (lawns)	Apply 1 L product/10 m <sup>2</sup>  Apply CUEVA® RTU WITH QUICK CONNECT SPRAYER when disease first appears, and repeat at 7 to 10 day intervals for a maximum of 15 applications. In frequently diseased areas, prune adjacent trees and shrubs to reduce turf shading and to improve air	Powdery mildew ( <i>Erysiphe graminis</i> )

PMRA Approved Label  
 2015-3956  
 2015-08-25

	movement.	
--	-----------	--

<b>ORNAMENTALS (OUTDOOR AND GREENHOUSE)</b>		
<b>Plant</b>	<b>Application Rate and Notes</b>	<b>Diseases Controlled or Suppressed</b>
Greenhouse and outdoor ornamental shrubs and flowering plants ( such as rose, hollyhock, hydrangea, crape myrtle)	Apply 1 L product/10 m <sup>2</sup> . Re-apply using 5-10 day intervals.  CUEVA® RTU WITH QUICK CONNECT SPRAYER may cause copper toxicity on some rose varieties. Copper toxicity appears as purple spots.  In damp cool conditions (below 18°C), phytotoxicity is likely to occur with the use of CUEVA® RTU WITH QUICK CONNECT SPRAYER	Corynespora leaf spot ( <i>Corynespora asiicola</i> ) – Suppression Powdery mildew ( <i>Erysiphe</i> spp.), Rust ( <i>Puccinia malvacearum</i> ) Bacterial blight ( <i>Pseudomonas syringae</i> ) Fire blight ( <i>Erwinia amylovora</i> ) Coryneum blight ( <i>Thyrostroma carpophilum</i> )  Black spot ( <i>Diplocarpon rosae</i> ) Powdery mildew ( <i>Podosphaera pannosa</i> var. <i>rosae</i> ) Rust ( <i>Phragmidium mucronatum</i> )
Rose		
Crape Myrtle		Cercospora leaf spot ( <i>Cercospora lythracearum</i> ) -Suppression

<b>VEGETABLES, FRUITS, NUTS, and BERRIES</b>		
<b>Plant</b>	<b>Application Rate and Notes</b>	<b>Diseases Controlled or Suppressed</b>
Root and Tuber Vegetables ( Garden beet, Celeriac, Potatoes, Sugar Beet)	Apply 1 L product/10 m <sup>2</sup> Re-apply using 5-10 day intervals.	Early blight ( <i>Alternaria solani</i> ) on potato Late blight ( <i>Phytophthora infestans</i> ) on potato Septoria leaf spot ( <i>Septoria lycopersici</i> ) on potato Septoria (late) blight ( <i>Septoria apiicola</i> ) on celeriac

PMRA Approved Label  
 2015-3956  
 2015-08-25

		Cercospora leaf spot ( <i>Cercospora beticola</i> ) on beet and sugar beet
Bulb Vegetables (Chives, Garlic, Leek, Onion, Shallot)	Apply 1 L product/10 m <sup>2</sup> Re-apply using 5-10 day intervals.	Downy mildew ( <i>Peronospora destructor</i> ) Botrytis leaf blight ( <i>Botrytis squamosa</i> ) Soft rot ( <i>Erwinia carotovora</i> subsp. <i>carotovora</i> )
Celery	Apply 1 L product/10 m <sup>2</sup> Re-apply using 5-10 day intervals.	Cercospora (early) blight ( <i>Cercospora apicola</i> ) Septoria (late) blight ( <i>Septoria apicola</i> )
Brassica Leafy Vegetables (Bok choy, Broccoli, Brussels sprouts, Cabbage, Cauliflower, Kale, Kohlrabi, Mustard, Pak-choi)	Apply 1 L product/10 m <sup>2</sup> Re-apply using 5-10 day intervals..  For Brussels sprouts, do not exceed 10 applications per year.	Black rot ( <i>Xanthomonas campestris</i> pv. <i>campestris</i> ) - Suppression
Legumes (Bean, Pea, Soybean)	Apply 1 L product/10 m <sup>2</sup> Re-apply using 5-10 day intervals.	Ascochyta blight ( <i>Ascochyta pisi</i> ) Halo blight ( <i>Pseudomonas syringae</i> pv. <i>phaseolicola</i> ) Common blight ( <i>Xanthomonas campestris</i> pv. <i>phaseoli</i> ) Brown spot ( <i>Pseudomonas syringae</i> pv. <i>syringae</i> ) Powdery mildew ( <i>Erysiphe</i> spp.) Rust ( <i>Uromyces appendiculatus</i> )
Fruiting Vegetables (Eggplant, Peppers, Tomatoes)	Apply 1 L product/10 m <sup>2</sup> Re-apply using 5-10 day intervals.	Early blight ( <i>Alternaria solani</i> ) Late blight ( <i>Phytophthora infestans</i> ) Septoria leaf spot ( <i>Septoria lycopersici</i> ) Bacterial speck ( <i>Pseudomonas syringae</i> ) Bacterial leaf spot ( <i>Xanthomonas campestris</i> pv. <i>vesicatoria</i> ) Bacterial canker ( <i>Clavibacter michiganensis</i> pv. <i>michiganensis</i> )
Cucurbits (Cucumbers, Cantaloupe, Melon, Squash, Pumpkin, Zucchini)	Apply 1 L product/10 m <sup>2</sup> Re-apply using 5-10 day intervals.	Powdery mildew ( <i>Podosphaera xanthii</i> ; <i>Erysiphe cichoracearum</i> ) Downy mildew ( <i>Pseudoperonospora cubensis</i> ) Alternaria leaf blight

PMRA Approved Label

2015-3956

2015-08-25

		<p>(<i>Alternaria cucumerina</i>)          Anthracnose (<i>Colletotrichum orbiculare</i>)          Angular leaf spot          (<i>Pseudomonas syringae</i> pv. <i>orbiculare</i>)          Bacterial wilt (<i>Erwinia tracheiphila</i>)          Septoria leaf spot (<i>Septoria cucurbitacearum</i>)</p>
<p>Pome Fruit Trees (Apple, Pear, Quince)</p>	<p>Apply 1 L product/10 m<sup>2</sup>          Re-apply using 5-10 day intervals.          Do not exceed 10 applications per year.</p> <p>For fireblight control, apply CUEVA® RTU WITH QUICK CONNECT SPRAYER in the dormant period, during bloom, or in-season cover spray applications.</p> <p><b>Do not apply to light-skinned apples after fruit has formed or russetting may occur.</b></p>	<p>Fire blight (<i>Erwinia amylovora</i>)          Scab (<i>Venturia Inaequalis</i>, <i>V. pirina</i>)</p>
<p>Stone Fruit Trees (Apricot, Cherry, Nectarine, Peach, Plum, Necta-plum)</p>	<p>Apply 1 L product/10 m<sup>2</sup>          Re-apply using 5-10 day intervals.</p> <p>For peaches, do not exceed 5 applications per year.</p> <p>For nectarines, do not exceed 10 applications per year.</p> <p>For bacterial canker, apply as a dormant spray as buds begin to swell, repeating at the bud burst stage, and weekly thereafter as needed, up to six sprays. In the fall spray again at 10% and 80% leaf fall. For brown rot blossom blight apply full cover spray at delayed dormant (bud swell), popcorn, full bloom and petal fall stages. During wet weather additional bloom sprays may be necessary.</p> <p>Apply to leaf and fruit spot, Coryneum blight, and peach leaf curl as a dormant spray in late fall during a period of dry weather.</p>	<p>Peach leaf curl (<i>Taphrina deformans</i>)          Bacterial spot (<i>Xanthomonas campestris</i> pv. <i>pruni</i>)          Coryneum blight (<i>Thyrostroma carpophilum</i>)          Bacterial canker (<i>Pseudomonas syringae</i> pv. <i>syringae</i>, <i>morsprunum</i>)          Brown rot (<i>Monilinia fructicola</i>)          Leaf and fruit spot (<i>Blumeriella jaapii</i>)</p>
<p>Small Fruits (Blackberry,</p>	<p>Apply 1 L product/10 m<sup>2</sup></p>	<p>Powdery mildew (<i>Erysiphe</i></p>

PMRA Approved Label

2015-3956

2015-08-25

Blueberry, Currant, Gooseberry, Grape, Raspberry, Strawberry)	<p>Apply at the start of flowering and continue every 7 to 10 days. Do not apply within 1 day of harvest.</p> <p>For strawberries, spray 1 month after planting (or before flowering on established plants) and twice more at 7 day intervals.</p> <p>For grapes, do not mix CUEVA® RTU WITH QUICK CONNECT SPRAYER with lime. Certain <i>Vitis vinifera</i> and French hybrid varieties may be sensitive to copper sprays resulting in marginal leaf burn. Make test sprays before spraying these varieties</p>	<p><i>necator</i>) on grape</p> <p>Powdery mildew (<i>Podosphaera aphanis</i>) on strawberry</p> <p>Downy mildew (<i>Plasmopara viticola</i>) on grape</p> <p>Rust (<i>Cronartium ribicola</i>) on currant and gooseberry</p> <p>Bacterial blight (<i>Pseudomonas syringae</i> pv. <i>Syringae</i>) on raspberry, blackberry and blueberry</p>
Filbert, Hazelnut, ,	<p>Apply 1 L product/10 m<sup>2</sup></p> <p>Re-apply using 5-10 day intervals.</p>	<p>Bacterial blight (<i>Xanthomonas campestris</i> pv. <i>corylina</i>)</p> <p>Eastern filbert blight (<i>Anisogramma anomala</i>)</p>
Walnut	<p>Apply 1 L product/10 m<sup>2</sup></p> <p>Re-apply using 5-10 day intervals.</p>	<p>Bacterial blight (<i>Xanthomonas campestris</i> pv. <i>juglandis</i>)</p>
Parsley	<p>Apply 1 L product/10 m<sup>2</sup></p> <p>Re-apply using 5-10 day intervals.</p>	<p>Septoria leaf spot (<i>Septoria petroselinii</i>)</p>

**GREENHOUSE FOOD CROPS**

Crop or Plant	Application Rate and Notes	Diseases Controlled or Suppressed
Celery	<p>Apply 1 L product/10 m<sup>2</sup></p> <p>Re-apply using 5-10 day intervals.</p>	<p>Cercospora (early) blight (<i>Cercospora apiicola</i>)</p> <p>Septoria (late) blight (<i>Septoria apiicola</i>)</p>
Legumes (Bean, Pea, Soybean)	<p>Apply 1 L product/10 m<sup>2</sup></p> <p>Re-apply using 5-10 day intervals.</p>	<p>Ascochyta blight (<i>Ascochyta pisi</i>)</p> <p>Halo blight (<i>Pseudomonas syringae</i> pv. <i>phaseolicola</i>)</p> <p>Common blight (<i>Xanthomonas campestris</i> pv. <i>phaseoli</i>)</p> <p>Brown spot (<i>Pseudomonas syringae</i> pv. <i>syringae</i>)</p> <p>Powdery mildew (<i>Erysiphe</i> spp.)</p> <p>Rust (<i>Uromyces appendiculatus</i>)</p>
Fruiting Vegetables (Eggplant, Pepper, Tomatoes)	<p>Apply 1 L product/10 m<sup>2</sup></p> <p>Re-apply using 5-10 day intervals.</p>	<p>Early blight (<i>Alternaria solani</i>)</p> <p>Late blight (<i>Phytophthora infestans</i>)</p> <p>Septoria leaf spot (<i>Septoria lycopersici</i>)</p> <p>Bacterial speck (<i>Pseudomonas syringae</i>)</p> <p>Bacterial leaf spot (<i>Xanthomonas</i></p>

PMRA Approved Label

2015-3956

2015-08-25

		<i>campestris</i> pv <i>vesicatoria</i> ) Bacterial canker ( <i>Clavibacter michiganensis</i> pv. <i>michiganensis</i> )
Cucurbits (Cucumbers, Cantaloupe, Melon, Squash, Pumpkin, Zucchini)	Apply 1 L product/10 m <sup>2</sup> Re-apply using 5-10 day intervals.	Powdery mildew ( <i>Podosphaera xanthii</i> ; <i>Erysiphe cichoracearum</i> ) Downy mildew ( <i>Pseudoperonospora cubensis</i> ) Alternaria leaf blight ( <i>Alternaria cucumerina</i> ) Anthracnose ( <i>Colletotrichum orbiculare</i> ) Angular leaf spot ( <i>Pseudomonas syringae</i> pv. <i>orbiculare</i> ) Bacterial wilt ( <i>Erwinia tracheiphila</i> ) Septoria leaf spot ( <i>Septoria cucurbitacearum</i> )

**PRECAUTIONS**

KEEP OUT OF REACH OF CHILDREN. Use only in well ventilated areas. For good hygiene practice, wear gloves when handling this product.

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

DO NOT allow adults, children or pets to enter the treated area until sprays have dried.

**TOXICOLOGICAL INFORMATION**

Treat symptomatically.

**ENVIRONMENTAL HAZARDS**

TOXIC to birds, small wild mammals and aquatic organisms. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

**FIRST AID**

**If on skin or clothing**, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If in eyes**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take the container label or product name and Pest Control Product Registration Number with you when seeking medical attention.

PMRA Approved Label

2015-3956

2015-08-25

#### **STORAGE**

Store in original container in a cool, dry, well ventilated area. To prevent contamination store this product away from food or feed. Protect from freezing.

#### **DISPOSAL/DECONTAMINATION**

DO NOT reuse the empty containers. Dispose in household garbage. Unused or partially used products should be disposed at provincially or municipally designated hazardous waste disposal sites.

#### **NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.



Cueva® is a registered trademark of W. Neudorff GmbH KG

- Quick Connect™ is a registered trademark of OMS Investments, Inc., used under license by Scotts Canada Ltd.