

APRON® ADVANCE

SEED TREATMENT

SUSPENSION

COMMERCIAL - AGRICULTURAL

A seed treatment product for control of certain seed-borne and soil-borne diseases of dry beans (including lupins and dry faba beans), dry peas, chickpeas and lentils.

ACTIVE INGREDIENTS:

Thiabendazole 150 g/L
Fludioxonil..... 25 g/L
Metalaxyl-M and S-isomer 20 g/L

Contains 1,2-benzisothiazolin-3-one at 0.015% as a preservative.

**READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

REGISTRATION NO: **30627**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **1 L to 1050 L**

ALL SEED TREATED WITH THIS PRODUCT MUST BE CONSPICUOUSLY COLOURED

For use with commercial seed treaters (facilities and mobile treaters).

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, ON N1G 4Z3
Telephone: 1-877-964-3682

LABEL

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a doctor or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

3.0 TOXICOLOGICAL INFORMATION

NOTE TO PHYSICIAN: Treat symptomatically.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Do not contaminate food, feed, or domestic water supplies.

Do not apply in a way that this product will contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area when seed is being treated or bagged.

For use with commercial seed treaters (facilities and mobile treaters). Treat seed in a well-ventilated area. Keep treated seeds out of reach of children and animals.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

For On-Farm Seed Treatment: Workers must wear cotton coveralls over a long-sleeved shirt and long pants, hat, chemical-resistant gloves, and a NIOSH/MSHA approved dust mask during mixing, loading, treating, clean-up, and maintenance of seed treatment equipment.

For Commercial Seed Treatment: Workers must wear chemical-resistant coveralls over a long-sleeved shirt and long pants, chemical-resistant gloves during mixing, loading, treating, clean-up, maintenance of seed treatment equipment, and bagging, sewing or stacking of bagged treated seed. Wear a NIOSH/MSHA approved dust mask when bagging or sewing bags of treated seed or when transferring seed to a storage bin.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash splashes from skin IMMEDIATELY with plenty of water. Remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. As soon as possible, wash thoroughly and change into clean clothing. After spraying, wash hands and shower thoroughly with soap and water.

Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. Do not reuse them.

Users should wash hands and exposed skin before eating, drinking, chewing gum, using tobacco, applying cosmetics or using the toilet.

While using product, do not eat, drink or use tobacco (including smoking).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Wear freshly laundered clothes daily.

Do not wear contaminated shoes.

6.0 ENVIRONMENTAL PRECAUTIONS

This product is TOXIC to fish and other aquatic organisms. Do not apply directly to aquatic habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, ditches and wetlands), estuaries or marine habitats.

Do not contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Treated seed is TOXIC to birds. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned up from the soil surface.

7.0 USE RESTRICTIONS

1. Do not use treated seed for food, feed or oil purposes.
2. Store away from food, feed, and out of the reach of children and animals.
3. Do not plant any crop other than dry beans, dry peas, chickpeas or lentils within 30 days to

fields in which treated seeds were planted.

4. All bags containing treated seed for sale or use in Canada must be labeled or tagged as follows: **“This seed has been treated with thiabendazole, fludioxonil and metalaxyl-M and S-isomer fungicides. Do not use for food, feed or oil purposes. Keep out of reach of children and animals. Store away from feeds and foodstuffs. When handling treated seed, wear a long-sleeved shirt, long pants, and chemical resistant gloves. Do not plant any crop other than dry beans, dry peas, chickpeas or lentils within 30 days to fields in which treated seeds were planted. For dry peas, do not graze treated crops or cut for forage or hay within 65 days of planting. For all other labelled crops, do not graze treated crops or cut for forage or hay within 30 days of planting. Treated seed is TOXIC to birds. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned up from the soil surface.”**

8.0 STORAGE

Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated area away from other pesticides, feed and foodstuffs, and out of the reach of children and animals.

Ideal storage temperature of bulk material is above freezing and below 30 °C.

9.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

For recyclable containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for future use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

APRON® ADVANCE

SEED TREATMENT

SUSPENSION

COMMERICAL - AGRICULTURAL

A seed treatment product for control of certain seed-borne and soil-borne diseases of dry beans (including lupins and dry faba beans), dry peas, chickpeas and lentils.

ACTIVE INGREDIENTS:

Thiabendazole 150 g/L
Fludioxonil..... 25 g/L
Metalaxyl-M and S-isomer 20 g/L

Contains 1,2-benzisothiazolin-3-one at 0.015% as a preservative.

**READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

REGISTRATION NO: **30627**
PEST CONTROL PRODUCTS ACT

ALL SEED TREATED WITH THIS PRODUCT MUST BE CONSPICUOUSLY COLOURED

For use with commercial seed treaters (facilities and mobile treaters).

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, ON N1G 4Z3
Telephone: 1-877-964-3682

PAMPHLET

TABLE OF CONTENTS

	<u>Section Number</u>
NOTICE TO USER	1.0
FIRST AID	2.0
TOXICOLOGICAL INFORMATION	3.0
PRECAUTIONS.....	4.0
PERSONAL PROTECTIVE EQUIPMENT (PPE).....	5.0
ENVIRONMENTAL PRECAUTIONS.....	6.0
USE RESTRICTIONS.....	7.0
STORAGE	8.0
DISPOSAL	9.0
PRODUCT INFORMATION	10.0
DIRECTIONS FOR USE	11.0
General Information	11.1
Seed Treatment and Inoculants.....	11.2
CROP USE DIRECTIONS	12.0
Pulses	12.1
Tank Mixes.....	12.2
RESISTANCE MANAGEMENT RECOMMENDATIONS	13.0

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a doctor or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

3.0 TOXICOLOGICAL INFORMATION

NOTE TO PHYSICIAN: Treat symptomatically.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

Do not contaminate food, feed, or domestic water supplies.

Do not apply in a way that this product will contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area when seed is being treated or bagged.

For use with commercial seed treaters (facilities and mobile treaters). Treat seed in a well-ventilated area. Keep treated seeds out of reach of children and animals.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

For On-Farm Seed Treatment: Workers must wear cotton coveralls over a long-sleeved shirt and long pants, hat, chemical-resistant gloves, and a NIOSH/MSHA approved dust mask during mixing, loading, treating, clean-up, and maintenance of seed treatment equipment.

For Commercial Seed Treatment: Workers must wear chemical-resistant coveralls over a long-sleeved shirt and long pants, chemical-resistant gloves during mixing, loading, treating, clean-up, maintenance of seed treatment equipment, and bagging, sewing or stacking of bagged treated seed. Wear a NIOSH/MSHA approved dust mask when bagging or sewing bags of treated seed or when transferring seed to a storage bin.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash splashes from skin IMMEDIATELY with plenty of water. Remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. As soon as possible, wash thoroughly and change into clean clothing. After spraying, wash hands and shower thoroughly with soap and water.

Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. Do not reuse them.

Users should wash hands and exposed skin before eating, drinking, chewing gum, using tobacco, applying cosmetics or using the toilet.

While using product, do not eat, drink or use tobacco (including smoking).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Wear freshly laundered clothes daily.

Do not wear contaminated shoes.

6.0 ENVIRONMENTAL PRECAUTIONS

This product is TOXIC to fish and other aquatic organisms.

Do not apply directly to aquatic habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, ditches and wetlands), estuaries or marine habitats.

Do not contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Treated seed is TOXIC to birds. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned up from the soil surface.

7.0 USE RESTRICTIONS

1. Do not use treated seed for food, feed or oil purposes.
2. Store away from food, feed, and out of the reach of children and animals.

3. Do not plant any crop other than dry beans, dry peas, chickpeas or lentils within 30 days to fields in which treated seeds were planted.
4. All bags containing treated seed for sale or use in Canada must be labeled or tagged as follows: **“This seed has been treated with thiabendazole, fludioxonil and metalaxyl-M and S-isomer fungicides. Do not use for food, feed or oil purposes. Keep out of reach of children and animals. Store away from feeds and foodstuffs. When handling treated seed, wear a long-sleeved shirt, long pants, and chemical resistant gloves. Do not plant any crop other than dry beans, dry peas, chickpeas or lentils within 30 days to fields in which treated seeds were planted. For dry peas, do not graze treated crops or cut for forage or hay within 65 days of planting. For all other labelled crops, do not graze treated crops or cut for forage or hay within 30 days of planting. Treated seed is TOXIC to birds. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned up from the soil surface.”**

8.0 STORAGE

Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated area away from other pesticides, feed and foodstuffs, and out of the reach of children and animals.

Ideal storage temperature of bulk material is above freezing and below 30 °C.

9.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

For recyclable containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for future use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

10.0 PRODUCT INFORMATION

APRON ADVANCE Seed Treatment is a seed treatment product containing the fungicides thiabendazole, fludioxonil, and metalaxyl-M and S-isomer. This product controls certain seed-borne and soil-borne diseases of dry beans, dry peas, chickpeas and lentils.

APRON ADVANCE Seed Treatment is active against a wide range of plant diseases caused by *Fusarium* spp., *Rhizoctonia* spp., *Pythium* spp., *Ascochyta* spp. and seed-borne *Colletotrichum* spp. and *Botrytis* spp.

Note: Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigour and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination, reduced emergence and/or reduction of seed and seedling vigour. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of the seed before committing the total seed lot to a selected seed treatment. Due to seed quality and seed storage conditions beyond the control of Syngenta Canada Inc. no claims are made to guarantee the germination of carry-over treated seed.

Note: Experience has shown that strains of fungus resistant to metalaxyl-M may develop. Failure to control the disease will likely result in crop damage and/or yield losses. Since the occurrence of resistance cannot be foreseen, Syngenta Canada Inc. accepts no responsibility for any loss of, or damage to, crops resulting from the failure of APRON ADVANCE Seed Treatment to control the resistant fungus strains. If disease appears in a treated field, consult the government extension specialist immediately.

11.0 DIRECTIONS FOR USE

11.1 General Information

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

APRON ADVANCE Seed Treatment is a seed treatment formulation for use in commercial seed treatment plants, and for on-farm treatment using auger treating only; do not use in hopper box or seed drill. The equipment must provide uniform coverage of APRON ADVANCE Seed Treatment on the seed. Uneven seed coverage may not give the desired level of disease control.

Thoroughly mix the recommended amount of APRON ADVANCE Seed Treatment with the required amount of water for the slurry treater and dilution rate to be used. Slurry volumes will vary depending on seed size. Follow the manufacturer's application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment. Allow the seed to dry before bagging, storing or seeding.

This product contains a pigment that will colour the treated seed; however, users are

responsible for ensuring that the treated seed, when dried and ready for bagging, storage or seeding has an unnatural colour. If the pigment contained in the formulation does not colour the seed adequately, additional colourant must be added to the mixture while treating the seed. Follow instructions on the colourant package for mixing. Regulations pertaining to colouration of treated seed enforced under the “Seeds Act” must be strictly adhered to when using this product.

11.2 Seed Treatment and Inoculants

APRON ADVANCE Seed Treatment is flexible with Rhizobium based inoculants. Depending on compatibility at application, inoculants can be applied as a tank mix through simultaneous or sequential applications. Please check with inoculant manufacturers for application and compatibility details prior to use. Consult the manufacturer of the application equipment planned to be used for suitability for this application and for instructions on operation and calibration of the equipment.

Seed treated with APRON ADVANCE Seed Treatment, or a combination of APRON ADVANCE Seed Treatment and seed inoculants, may not flow through a seed drill at the same rate as untreated seed. Recalibrate the seed drill before planting treated seed. Mixing with inoculants may increase drying time while treating, extending the processing time.

12.0 CROP USE DIRECTIONS

12.1 Pulses

Apply 100 mL of APRON ADVANCE Seed Treatment per 100 kg of seed to the crops listed below. At this rate, APRON ADVANCE Seed Treatment delivers 15 g of thiabendazole, 2.5 g of fludioxonil, and 2 g of metalaxyl-M and S-isomer per 100 kg of seed.

PULSES	
CROP	Dry Beans (including lupins and dry faba beans)
DISEASES CONTROLLED	Seed rot/pre-emergence damping-off, and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Pythium</i> spp. and <i>Rhizoctonia</i> spp. Seedling blight caused by <i>Pythium</i> spp. Anthracnose caused by seed-borne <i>Colletotrichum</i> spp.
RATE	100 mL / 100 kg of seed
APPLICATION METHOD	Seed Treatment
CROP	Chickpeas
DISEASES CONTROLLED	*Seed-borne <i>Ascochyta</i> blight caused by <i>Ascochyta rabiei</i> Seed rot/pre-emergence damping-off, and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Pythium</i> spp. and <i>Rhizoctonia</i> spp. Seedling blight caused by <i>Fusarium</i> spp. and <i>Pythium</i> spp. Seed rot and seedling blight caused by seed-borne <i>Botrytis</i> spp.
RATE	100 mL / 100 kg of seed
APPLICATION METHOD	Seed Treatment
CROP	Lentils
DISEASES CONTROLLED	*Seed-borne <i>Ascochyta</i> blight caused by <i>Ascochyta lentis</i> Seed rot/pre-emergence damping-off, post-emergence damping-off and seedling blight caused by <i>Fusarium</i> spp., <i>Pythium</i> spp. and <i>Rhizoctonia</i> spp. Seedling root rot caused by <i>Fusarium</i> spp.

	Seed rot and seedling blight caused by seed-borne <i>Botrytis</i> spp.
RATE	100 mL / 100 kg of seed
APPLICATION METHOD	Seed Treatment
CROP	Dry Peas
DISEASES CONTROLLED	*Seed-borne <i>Ascochyta</i> blight and foot rot caused by <i>Ascochyta pinodes</i> Seed rot/pre-emergence damping-off, post-emergence damping-off and seedling blight caused by <i>Fusarium</i> spp., <i>Pythium</i> spp. and <i>Rhizoctonia</i> spp.
RATE	100 ml/ 100 kg of seed
APPLICATION METHOD	Seed Treatment

**Ascochyta*-infected seed is an important vehicle for moving the disease into new areas of the field due to the ability of the pathogen to readily transmit from the infected seed to seedlings. APRON ADVANCE Seed Treatment will minimize the growth of fungus on the seed and prevent the movement of fungus in the field. APRON ADVANCE Seed Treatment is an important tool in an integrated approach to *Ascochyta* management in pulse crops.

12.2 Tank Mixes

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682 for information before applying any tank mix that is not specifically recommended on this label.

Syngenta Canada Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) of all potential tank mixes under all environmental conditions or for all crop varieties. Tank mixes that are not specifically listed on this label should be tested on a small area first, under local conditions and using standard practices, to confirm the tank mix is suitable for widespread application.

To determine the physical compatibility of this product with other products, use a jar test.

Always read and follow label directions including WALES mixing order.

13.0 RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that APRON ADVANCE Seed Treatment contains Group 1, 4 and 12 fungicides. Any fungal population may contain individuals naturally resistant to APRON ADVANCE Seed Treatment and other Group 1, 4 and 12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of APRON ADVANCE Seed Treatment or other Group 1, 4 and 12 fungicides with different groups that control the same pathogens.

Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.

Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for signs of resistance development. Notify Syngenta Canada Inc. if reduced sensitivity of pathogen to APRON ADVANCE Seed Treatment is suspected.

If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information or to report suspected resistance, contact company representatives at 1-87-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.