

2015-7108
2016-02-04

CARTON

**WILSON[®] PREDATOR[™]
RAT BAIT STATION
PRE-FILLED**

Ready To Use

Kills Norway Rats and Roof Rats

SOLID

DOMESTIC

REG. NO. 32153 PCP ACT

GUARANTEE: Chlorophacinone.....0.005%

Warning, contains the allergens peanuts, wheat, soy and fish

**KEEP OUT OF REACH OF CHILDREN
READ THE LABEL BEFORE USING**

**THIS BAIT STATION IS NOT TAMPER RESISTANT.
USE INDOORS ONLY IN AREAS INACCESSIBLE TO CHILDREN AND PETS.**

1 disposable bait station (contains 113 g of bait)

Sure-Gro IP Inc./Lic. Premier Tech Home & Garden Inc.
P.O. Box 21001, Brantford, ON N3R 7W9
1-800-268-2806

DIRECTIONS FOR USE / USE LIMITATIONS: Use **Wilson® Predator™ Rat Bait Station Pre-Filled** for the control of Norway rats (*Rattus norvegicus*) and Roof rats (*Rattus rattus*) **INDOORS IN LOCATIONS NOT ACCESSIBLE TO CHILDREN, PETS, DOMESTIC ANIMALS, OR NON-TARGET WILDLIFE. DO NOT USE THIS PRODUCT OUTDOORS.** DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Selection of Treatment Areas: Place bait in locations rats travel and will most likely find and consume the bait. Placement along walls, by gnawed openings, in corners and concealed places, between floors or walls, or in locations where rats or their signs have been seen. Remove as much alternative food as possible.

APPLICATION DIRECTIONS: Remove bait station from package and place in horizontal position with side against wall, exposing entrance. For Norway Rats (*Rattus norvegicus*) and Roof Rats (*Rattus rattus*): Norway rats usually inhabit the ground floor of structures and burrow in the ground. They will range approximately 15 to 100 m from their nesting area seeking food and water. Roof rats, however, are excellent climbers and usually infest the upper stories of structures. Hence, special attention must be paid to this species ability to climb by placing baits in the roof rats activity area. For both species, eliminate as much of the naturally occurring food as possible and place 1 to 5 bait stations (at intervals of 5-10 m) within the treatment area. Maintain an uninterrupted supply of fresh bait for at least 10 days, or until signs of rat activity cease.

Replace bait stations if contents are consumed or contaminated. Wearing rubber gloves, collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation occurs, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations as necessary.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Rubber gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.** This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

FIRST AID: If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **If on skin or clothing,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. **If swallowed,** for all cases of human ingestion, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **If pet or**

livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Vitamin K1 in the form of intramuscular or subcutaneous injections, or by oral ingestion is a suggested remedial treatment for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE AND DISPOSAL: Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals. Wrap used or damaged bait stations and discard in household garbage. DO NOT use damaged bait station.

RODENT DISPOSAL: Dead rodents should be securely wrapped and placed in closed containers and disposed of in the garbage. Otherwise, securely wrap carcasses and bury to a depth that will make them inaccessible to scavengers.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

For additional product information or in case of emergency, spills or fire, call toll-free 1-800-268-2806. www.pthomeandgarden.com

BAIT STATION

**WILSON® PREDATOR™
RAT BAIT STATION
PRE-FILLED**

DOMESTIC

REG. NO. 32153 PCP ACT

GUARANTEE: Chlorophacinone.....0.005%

Warning, contains the allergens peanuts, wheat, soy and fish

WARNING



POISON

READ THE ENTIRE LABEL ON THE OUTER PACKAGING BEFORE USE

113 g

Sure-Gro IP Inc./Lic. Premier Tech Home & Garden Inc.
P.O. Box 21001, Brantford, ON N3R 7W9
1-800-268-2806