

GROUP **7** FUNGICIDE

TRIVAPRO® B Fungicide
SOLATENOL® technology

FUNGICIDE

EMULSIFIABLE CONCENTRATE

COMMERCIAL - AGRICULTURAL

Fungicide for control or suppression of certain broad-spectrum diseases in Cereals, Corn, and Soybean

ACTIVE INGREDIENT:

Benzovindiflupyr..... 100 g/L

READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN



POISON
DANGER: CORROSIVE TO EYES & SKIN

REGISTRATION NO: **32185**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **0.5 – 1000 L**

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

Label

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote. Treat symptomatically. Contains petroleum distillates. Vomiting may cause aspiration pneumonia.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Keep unused product in original container tightly closed, locked up and away from food and feed.

CORROSIVE to the eye. **DO NOT** get in eyes. Causes irreversible eye damage. Harmful or fatal if swallowed. Harmful if inhaled. Avoid inhaling/breathing sprays. Severely irritating to the skin. **DO NOT** get on skin.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours

Apply only to agricultural crops when the potential for drift to areas of human habitation and activity, such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear coveralls over a long-sleeved shirt, long pants, chemical resistant gloves, socks, chemical-resistant footwear and protective eyewear (goggles or face shield) during mixing, loading, application, clean up and repair. Protective eyewear, chemical resistant gloves and coveralls are not required during application within a closed cab and/or cockpit.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

All users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Benzovindiflupyr is persistent and may carryover. It is recommended that any products containing benzovindiflupyr not be used in areas treated with this product during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

7.0 STORAGE

Store this product away from food or feed. Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated area out of the reach of children and animals. This product does not require heated storage.

8.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for cleanup of spills.

For returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

TRIVAPRO® and SOLATENOL® are trademarks of their respective companies.

GROUP **7** FUNGICIDE

TRIVAPRO® B Fungicide
SOLATENOL® echnology

FUNGICIDE

EMULSIFIABLE CONCENTRATE

COMMERCIAL – AGRICULTURAL

Fungicide for control or suppression of certain broad-spectrum diseases in Cereals, Corn, and Soybean

ACTIVE INGREDIENT:

Benzovindiflupyr..... 100 g/L

**READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**



**POISON
DANGER: CORROSIVE TO EYES & SKIN**

REGISTRATION NO: **32185**
PEST CONTROL PRODUCTS ACT

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

Pamphlet

Table of Contents

	<u>Section Number</u>
NOTICE TO USER	1.0
FIRST AID	2.0
TOXICOLOGICAL INFORMATION	3.0
PRECAUTIONS	4.0
PERSONAL PROTECTIVE EQUIPMENT (PPE)	5.0
ENVIRONMENTAL PRECAUTIONS	6.0
STORAGE	7.0
DISPOSAL	8.0
PRODUCT INFORMATION	9.0
General Use Precautions	9.1
DIRECTIONS FOR USE	10.0
General Information	10.1
Ground Application	10.2
Mixing Instructions	10.2.1
Spraying Instructions	10.2.2
Equipment Clean-Up	10.2.3
Aerial Application	10.3
Operator Precautions	10.3.1
Pilot Precautions	10.3.2
Mixing Instructions	10.3.3
Spraying Instructions	10.3.4
Equipment Clean-Up	10.3.5
Product Specific Precautions	10.4
Spray Buffer Zones	10.5
Rotational Crop Restrictions	10.6
CROP USE DIRECTIONS	11.0
Soybeans	11.1
Cereals	11.2
Corn	11.3
RESISTANCE MANAGEMENT RECOMMENDATIONS	12.0

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote. Treat symptomatically. Contains petroleum distillates. Vomiting may cause aspiration pneumonia.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Keep unused product in original container tightly closed, locked up and away from food and feed.

CORROSIVE to the eye. **DO NOT** get in eyes. Causes irreversible eye damage. Harmful or fatal if swallowed. Harmful if inhaled. Avoid inhaling/breathing sprays. Severely irritating to the skin. **DO NOT** get on skin.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and

sprayer settings.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear coveralls over a long-sleeved shirt, long pants, chemical resistant gloves, socks, chemical-resistant footwear and protective eyewear (goggles or face shield) during mixing, loading, application, clean up and repair. Protective eyewear, chemical resistant gloves and coveralls are not required during application within a closed cab and/or cockpit.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

All users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Benzovindiflupyr is persistent and may carryover. It is recommended that any products containing benzovindiflupyr not be used in areas treated with this product during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

7.0 STORAGE

Store this product away from food or feed. Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated area out of the reach of children and animals. This product does not require heated storage.

8.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for cleanup of spills.

For returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

9.0 PRODUCT INFORMATION

TRIVAPRO® B Fungicide is a broad-spectrum product and is recommended for the control or suppression of many important plant diseases. TRIVAPRO B Fungicide provides disease control of many rust, leaf spots and powdery mildews. TRIVAPRO B Fungicide is primarily applied as a foliar spray and can be used in block, or alternating spray programs with other crop protection products. All applications should be made according to the use directions listed in the label.

9.1 General Use Precautions

Do not apply TRIVAPRO B Fungicide through any type of irrigation equipment.

DO NOT USE IN GREENHOUSES (ALONE OR IN TANK MIXTURES).

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Apply the higher rate and shorter interval under conditions of high disease pressures.

Unless indicated otherwise in the use directions for specific crops, make no more than two sequential application of TRIVAPRO B Fungicide before alternating with a non-Group 7 fungicide registered for the same disease.

Not all members of the crop groups have been tested for efficacy and phytotoxicity at the recommended label rates, and should be used at the discretion of the user.

10.0 DIRECTIONS FOR USE

10.1 General Information

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Thorough coverage is necessary to provide good disease control. Applications using nozzles, sufficient water volume and pressure to provide thorough and uniform coverage generally provide the most effective disease control.

To avoid spray drift, do not apply when conditions favor drift beyond the target area. Avoid spray overlap, as crop injury may occur.

For more information on spray equipment and calibration, consult sprayer manufacturers and provincial/territorial recommendations.

Do not apply through any ultra-low volume (ULV) spray system.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Syngenta Canada Inc. at 1-87-SYNGENTA (1-877-964-3682) for information before applying any tank mix that is not specifically recommended on this label.

Syngenta Canada Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) of all potential tank-mixes under all environmental conditions or for all crop varieties. Tank mixes that are not specifically listed on this label should be tested on a small area first, under local conditions and using standard practices, to confirm the tank mix is suitable for widespread application.

To determine the physical compatibility of this product with other products, use a jar test.

Always read and follow label directions including WALES mixing order.

10.2 Ground Application

- Apply in a minimum of 150 L of water per hectare, unless specified otherwise.
- Do not apply through any ultra-low volume (ULV) spray system.
- Thorough coverage is necessary to provide good disease control.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

10.2.1 Mixing Instructions

TRIVAPRO B Fungicide is an emulsifiable concentrate (EC) formulation. Prepare only the amount of spray mixture that is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

TRIVAPRO B Fungicide Alone (No Tank Mix)

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add TRIVAPRO B Fungicide to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after TRIVAPRO B Fungicide has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

Mixing in the Spray Tank

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and TRIVAPRO B Fungicide to the spray tank.
- Allow TRIVAPRO B Fungicide to completely disperse.
- Spray the mixture with the agitator running.

10.2.2 Spraying Instructions

1. Water Volume: Apply in a minimum spray volume of 150 L/ha OR the volume given in the crop and pest specific instructions tabulated below, whichever is LARGER.
2. Sprayer Agitation: Use a jet agitator or liquid sparge tub which recirculates 10% or more of the tank per minute. **DO NOT** use an air sparger.
3. Pump: Screens should be used to protect the pump and prevent clogging. Use 16 mesh or *coarser* screens on the suction side of the pump. **DO NOT** place a screen in the recirculation line. Use 50 mesh or *coarser* screens between the pump and boom.
4. Spray Nozzles: 80° or 110° drift reducing flat fan (e.g. those with a pre-orifice or turbulence chamber) or air induction nozzles are recommended. Use 50 mesh nozzle screens. **DO**

NOT use flood type nozzles, controlled droplet application equipment, spray foils or hollow cone nozzles.

5. Pressure: As recommended by the nozzle manufacturer to achieve ASAE medium sized droplets.
6. Apply at uniform speed and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to avoid potential crop injury from over application.

10.2.3 Equipment Clean-Up

Before Spraying:

Prior to using TRIVAPRO B Fungicide, ensure that the spray tank, chem handling equipment, lines and filter are thoroughly cleaned.

After Spraying:

Thoroughly clean application equipment immediately after spraying. **DO NOT** let TRIVAPRO B Fungicide residue to dry within application equipment.

When using tank mixes, consult the tank-mix partner label for additional clean-up instructions.

The following recommendations are provided:

1. Drain and flush tank walls, boom and all hoses for ten minutes with a clean water/detergent mixture. Rinse with clean water. **DO NOT** clean application equipment near desirable vegetation, wells or other water sources.
2. Remove all nozzles and screens and wash separately.
3. Dispose of all rinsate in accordance with provincial/territorial regulations.

10.3 Aerial Application

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan. Apply at least 45 litres of water per hectare.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions, and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. When no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

10.3.1 Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

10.3.2 Pilot Precautions

Read and understand the entire label before using this product. Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides. Ensure uniform application and a uniform spray with minimum potential for drift. To avoid streaked, uneven or overlapped application, use appropriate marking technology. GPS based marking is preferred.

10.3.3 Mixing Instructions

Mixing this product directly in the aircraft hopper **IS NOT** recommended. The use of chemical handling or managing equipment to load the hopper **IS** recommended. This product **MAY BE** inducted into a hopper which is prefilled with water or when the product and water are mixed prior to entering the hopper. This product **MAY BE** batch mixed and pumped into the hopper. In all cases the chemical handling equipment and hopper interior must be clean prior to use.

NOTE: WG and DF formulations are preferentially batch mixed.

NOTE: SC, SN, and SL formulations may be inducted or batch mixed.

NOTE: EC formulations are preferentially batch mixed.

It is **NOT** recommended to combine solid (WG or DF) formulations with liquid tank mix partners within a single batch. Batch mix WG or DF formulations first, pump into the hopper, and then add liquid tank mix partners by induction or as an additional batch mix. When tank mixing multiple products, follow the mixing order outlined below:

1. Pump water into the hopper to at least $\frac{1}{4}$ to $\frac{1}{2}$ of the desired spray volume. Engage hopper circulation, if possible.
2. Thoroughly batch mix any WG or DF formulation mix partners and agitate to ensure complete mixing. Pump into the hopper
3. Induct or thoroughly batch mix any SE or SC formulations.
4. Thoroughly batch mix TRIVAPRO B Fungicide and any EC formulation mix partners. EC formulations may be added to the batch from Step 3, if desired.
5. Induct or thoroughly batch mix any solution (SN or SL) formulation mix partners. SN/SL formulations may be added to the batch from Step 3, if desired.
6. Pump batch mixed SC, EC, and/or SN/SL products into the hopper.
7. Finish filling the hopper with water.
8. If it was not possible to engage hopper agitation in Step 1, do so as soon as possible once

- airborne.
9. Spray the pesticide suspension the same day as mixing.
 10. **DO NOT** mix, load or clean equipment where there is a potential to contaminate wells or aquatic systems.

Mixing in the Spray Tank

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and TRIVAPRO B Fungicide to the spray tank.
- Allow TRIVAPRO B Fungicide to completely disperse.
- Spray the mixture with the agitator running.

10.3.4 Spraying Instructions

1. Water Volume: Apply in a minimum spray volume of 45 L/ha OR the volume given in the crop and pest specific instructions tabulated below, whichever is LARGER.
2. Spray Nozzles: Use only ASAE medium or coarse nozzles rated as delivering droplets of volume median diameter of 300 microns or greater.
3. Pressure: As recommended by the nozzle manufacturer to achieve ASAE medium sized droplets.
4. Ensure hopper agitation is engaged whenever possible during flight.

10.3.5 Equipment Clean-Up

Before Spraying:

Prior to using TRIVAPRO B Fungicide, ensure that the hopper, chem handling equipment, lines and filter are thoroughly cleaned.

After Spraying:

Thoroughly clean application equipment immediately after spraying. **DO NOT** let TRIVAPRO B Fungicide residue to dry within application equipment.

When using tank mixes, consult the tank-mix partner label for additional clean-up instructions.

The following recommendations are provided:

1. Drain and flush tank walls, boom and all hoses for ten minutes with a clean water/detergent mixture. Rinse with clean water. **DO NOT** clean application equipment near desirable vegetation, wells or other water sources.
2. Remove all nozzles and screens and wash separately.
3. Dispose of all rinsate in accordance with provincial/territorial regulations.

10.4 Product Specific Precautions

Follow recommendations provided by local disease monitoring services or provincial/territorial spray calendars regarding the appropriate application timing for protectant fungicides in your area. Cultural practices such as canopy management and removal of overwintered plant debris should be integrated with the use of fungicides to reduce disease incidence. Use the higher rate, and the shorter application interval under conditions of heavy infection pressure, on highly susceptible varieties, or when environmental conditions are favourable for disease development.

TRIVAPRO B Fungicide treatments should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance, plant residue management, crop rotation, and proper timing and placement of fertilizer and irrigation. Fungicide applications should begin prior to disease infection and continue throughout the season following a resistance management strategy (see **Resistance Management Recommendations**).

Read and understand the entire label before opening this product. If you have questions, call Syngenta Canada Inc. at 1-877-964-3682 or obtain technical advice from the distributor or your provincial/territorial agricultural representative. Application of this specific product must meet and/or conform to the following:

Volume: Apply the recommended rate in a minimum spray volume of 45 litres per hectare.

10.5 Spray Buffer Zones

A spray buffer zone is NOT required for uses with hand-held application equipment permitted on this label.

Use of low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage, soil drench or soil incorporation do not require a spray buffer zone.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Spray Buffer Zones (metres) Required for the Protection of:				
			Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial Habitat
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Soybeans, Cereals (wheat, barley, oats, rye, triticale), corn (field, sweet, pop and specialty)		10	1	1	1	1
Aerial	Soybeans, Cereals (wheat, barley, oats, rye, triticale), corn (field, sweet, pop and specialty)	Fixed wing	675	15	5	1	15
		Rotary wing	325	10	2	1	10

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

10.6 Rotational Crop Restrictions

Please see the following table for the crop rotational restrictions:

Rotational Crops	Planting Time From Last TRIVAPRO B Fungicide Application
Potatoes and tuberous and corm vegetable subgroup (CG 1C) Dried shelled pea and bean subgroup (CG 6C) Soybeans Fruiting vegetables crop group (CG 8-09) Cucurbit vegetables crop group (CG 9) Cereals (wheat, barley, oats, rye, triticale) Corn (field, sweet, pop and specialty) Rapeseed subgroup (CG 20A)	0 days
All other crops Intended for Food and Feed	6 months/180 days

11.0 CROP USE DIRECTIONS

11.1 Soybeans

SOYBEANS	
Diseases controlled at 300-750 mL product/ha	Asian soybean rust (<i>Phakopsora pachyrhizi</i>)
Diseases controlled at 500-750 mL product/ha	Septoria brown spot (<i>Septoria glycines</i>)
Diseases suppressed at 500-750 mL product/ha	Frogeye leaf spot (<i>Cercospora sojina</i>) Pod and stem blight (<i>Diaporthe phaseolorum</i>)
Application Timing/Instructions	Make the first application prior to disease establishment. A second application can be made 7-14 days after the first application, when disease pressure is severe or when agronomic or weather conditions are conducive to disease development or movement. If disease pressure is high, use the highest rate and shortest interval. For best results, sufficient water volume must be used to provide thorough coverage. A minimum spray volume of 100-200 L/ha for ground applications is recommended. For aerial applications, a minimum of 45 L/ha of water is recommended.
Specific Use Restrictions	<ol style="list-style-type: none"> 1. Do not apply more than 1.5 L/ha/season of TRIVAPRO B Fungicide. 2. Do not apply within 14 days of harvest for soybean grain (14-day PHI). 3. Soybean forage and hay may be fed or harvested 1 day (1-day PHI) after last application. 4. For best management practices when applying TRIVAPRO B Fungicide to soybean, apply in tank-mix or pre-mix with another fungicide from a different mode of action group that controls the same pathogen. 5. Apply a maximum of two TRIVAPRO B Fungicide or other SDHI containing sprays per soybean crop. Follow all precautions, restrictions and directions on the labels of fungicide products used in an alternation program.

11.2 Cereals

CEREALS	
Crops	Wheat
Diseases controlled at 300-750 mL product/ha	Stem rust (<i>Puccinia graminis</i>) Leaf rust (<i>P. recondita</i>) Stripe rust (<i>P. striiformis</i>)
Diseases controlled at 500-750 mL product/ha	Septoria leaf blotch (<i>Septoria tritici</i>) Tan spot (<i>Pyrenophora tritici-repentis</i>) Powdery mildew (<i>Erysiphe graminis</i>)
Crops	Barley
Diseases suppressed at 300 mL product/ha	Net blotch (<i>Drechslera teres</i>) Barley scald (<i>Rhynchosporium secalis</i>) – Under low disease pressure only.
Diseases controlled at 300-750 mL product/ha	Leaf rust (<i>Puccinia hordei</i>) Stem rust (<i>P. graminis</i>) Crown rust (<i>P. coronata</i>) Stripe rust (<i>P. striiformis</i>)
Diseases controlled at 500-750 mL product/ha	Septoria leaf blotch (<i>Septoria tritici</i>) Tan spot (<i>Pyrenophora tritici-repentis</i>) Net blotch (<i>Drechslera teres</i>) Barley scald (<i>Rhynchosporium secalis</i>) Powdery mildew (<i>Erysiphe graminis</i>)
Crops	Rye
Diseases controlled at 300-750 mL product/ha	Stem rust (<i>P. graminis</i>) Stripe rust (<i>P. striiformis</i>) Leaf rust (<i>Puccinia recondita</i>)
Diseases controlled at 500-750 mL product/ha	Septoria leaf blotch (<i>Septoria tritici</i>) Tan spot (<i>Pyrenophora tritici-repentis</i>) Powdery mildew (<i>Erysiphe graminis</i>)
Crops	Oats
Diseases suppressed at 300 mL product/ha	Net blotch (<i>Drechslera teres</i>) Scald (<i>Rhynchosporium secalis</i>) – Under low disease pressure only.
Diseases controlled at 300-750 mL product/ha	Stem rust (<i>Puccinia graminis</i>) Crown rust (<i>P. coronata</i>) Leaf rust (<i>P. recondita</i>)
Diseases controlled at 500-750 mL product/ha	Septoria leaf blotch (<i>Septoria tritici</i>) Net blotch (<i>Drechslera teres</i>) Scald (<i>Rhynchosporium secalis</i>) Powdery mildew (<i>Erysiphe graminis</i>)
Crops	Triticale
Diseases controlled at 300-750 mL product/ha	Leaf rust (<i>P. recondita</i>)
Diseases controlled at 500-750 mL product/ha	Septoria leaf spot (<i>Septoria tritici</i>) Tan spot (<i>Pyrenophora tritici-repentis</i>) Net blotch (<i>Drechslera teres</i>) Scald (<i>Rhynchosporium secalis</i>) Powdery mildew (<i>Erysiphe graminis</i>)

CEREALS	
Application Timing/Instructions	<p>Apply TRIVAPRO B Fungicide prior to disease development. Make applications no closer than 14 days apart.</p> <p>For early disease suppression, apply at beginning of tillering to stem elongation (BBCH 21-30 or Feekes 2-4), typically when herbicide treatments are applied.</p> <p>For disease control or suppression on the stem or leaves, apply at beginning of stem elongation to the end of flowering (BBCH 29-70 or Feekes 4-10.5.4). If disease pressure is high, use the highest rate. A minimum spray volume of 100 L/ha for ground applications is recommended. For aerial applications, a minimum of 45 L/ha of water is recommended.</p>
Specific Use Restrictions	<ol style="list-style-type: none"> 1. Do not apply more than 1.5 L/ha/season of TRIVAPRO B Fungicide. 2. For best management practices when applying TRIVAPRO B Fungicide to cereal crops, apply in tank-mix or pre-mix with another fungicide from a different mode of action group that controls the same pathogen. 3. Do not apply more than a total of 2 Group 7 applications per season on cereals. 4. Do not apply after Feekes 10.5.4 or BBCH 70 (full flower). 5. Forage and hay may be fed or harvested 7 days (7-day PHI) after last application.

11.3 Corn

CORN (FIELD, SWEET, POPCORN AND SPECIALTY INCLUDING ALL CULTIVARS AND/OR HYBRIDS OF THESE, INCLUDING SEED PRODUCTION)	
Diseases controlled at 300-750 mL product/ha	Rust (<i>Puccinia sorghi</i>)
Diseases controlled at 500-750 mL product/ha	Grey leaf spot (<i>Cercospora zeae-maydis</i>)
Application Timing/Instructions	<p>Begin applications prior to disease onset when conditions are conducive for disease. Make applications no closer than 7 days apart. A minimum spray volume of 200 L/ha for ground applications is recommended. For aerial applications, a minimum of 45 L/ha of water is recommended.</p>
Specific Use Restrictions	<ol style="list-style-type: none"> 1. Do not apply more than 1.5 L/ha/season of TRIVAPRO B Fungicide. 2. Do not apply within 7 days of harvest for corn grain and sweet corn (7 day-PHI). 3. Forage may be fed or harvested 7 days (7-day PHI) after last application. 4. Do not apply more than 2 consecutive applications before switching to a non-Group 7 fungicide. Follow all precautions, restrictions and directions on the labels of fungicide products used in an alternation program.

12.0 RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, TRIVAPRO B Fungicide contains a Group 7 fungicide in the succinyl dehydrogenase inhibitor (SDHI) class. Any fungal population may contain individuals naturally resistant to benzovindiflupyr and other Group 7 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of benzovindiflupyr or other Group 7 fungicides with different groups that control the same pathogens.

Apply no more than 2 sequential applications unless otherwise stated in the crop section.

DO NOT apply at rates lower than recommended on the label.

Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.

Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for sign of resistance development. Notify Syngenta Canada Inc. if reduced sensitivity of the pathogen to TRIVAPRO B Fungicide is suspected.

If disease continues to progress after treatment with this product, DO NOT increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, to which pathogen resistance has not developed.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information or to report suspected resistance, contact company representatives at 1-87-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.

Application Limitations and Pre-harvest Intervals

Crop*	Maximum Product Rate/ha/application (mL/ha)	Maximum total (L/ha/Season)	Pre-Harvest Interval (PHI) (days)	Minimum Re-treatment Interval (days)
Soybeans	750	1.5	14	7
Soybean forage and hay	750	1.5	1	7
Cereals	750	1.5	FK 10.5.4	14
Cereal forage and hay	750	1.5	7	14
Corn	750	1.5	7	7
Corn forage	750	1.5	7	7
Sweet corn	750	1.5	7	7

*For specific crops in a group and use directions, refer to the Specific Directions For Use.

TRIVAPRO® and SOLATENOL® are trademarks of a Syngenta Group Company.