

GROUP	4	HERBICIDE
--------------	----------	------------------

Albaugh

**MCPA Amine 600
Liquid
[Selective] Herbicide**

CONTAINS MCPA AMINE SOLUTION

For use in wheat, barley, rye, oats, flax, peas, corn and pastures.

COMMERCIAL - AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: MCPA, present as dimethylamine salt ... 600 g /L

REGISTRATION NO. 32686 PEST CONTROL PRODUCTS ACT

POISON

DANGER – CORROSIVE TO EYES

(1.0 to 450 Litres)

Albaugh, LLC
1525 NE 36th Street
Ankeny, IA 50021
USA
1-800-247-8013

For medical or treatment information from exposure to this product, call 1-888-347-6732
(7 days/week, 24-hr).

For 24-hour chemical spill, leak, fire, exposure, or accident response information, call
CHEMTREC toll free at 1-800-424-9300.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes and clothing. Wash concentrate from skin or eyes immediately. After use, wash hands and other exposed skin. Avoid breathing spray mist.

Do not use in residential areas, which are defined as sites where bystanders may be present during or after spraying, including homes, schools, parks, playgrounds, playing fields and public buildings.

PROTECTIVE CLOTHING AND EQUIPMENT:

Handling the concentrate (mixing and loading): Wear a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves. Rinse gloves before removal. Wear goggles or face shield during mixing/loading. Remove and wash contaminated clothing before reuse.

Handling the dilute spray solution (during application or repairing or cleaning equipment): Wear a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves. Rinse gloves before removal. Remove and wash contaminated clothing before reuse. Aerial applicators and applicators using a closed cab are not required to wear chemical-resistant gloves.

ENVIRONMENTAL PRECAUTIONS:

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Surface runoff:

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

Leaching:

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

FIRST AID:

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove

contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote. Employ supportive care. Treatment should be based on judgement of the physician in response to reactions of the patient. High concentrations of MCPA may cause severe irritation to the eyes. Symptoms of overexposure to MCPA could include slurred speech, twitching, jerking and spasms, drooling, low-blood pressure and unconsciousness. Treat symptomatically.

STORAGE:

Store this product away from food or feed. Store the container tightly closed away from seeds, fertilizer, plants and foodstuffs.

Store at temperatures above 0°C. If subjected to freezing temperatures, warm to at least 5°C and mix thoroughly. SHAKE WELL BEFORE USING.

DISPOSAL:

For Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site.

Contact your local distributor/dealer or municipality for the location of the nearest collection site.

Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or The provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.

GROUP	4	HERBICIDE
--------------	----------	------------------

Albaugh

**MCPA Amine 600
Liquid
[Selective] Herbicide**

CONTAINS MCPA AMINE SOLUTION

For use in wheat, barley, rye, oats, flax, peas, corn and pastures.

COMMERCIAL - AGRICULTURAL

READ THE LABEL AND BOOKLET BEFORE USING

ACTIVE INGREDIENT: MCPA, present as dimethylamine salt ... 600 g /L

REGISTRATION NO. 32686 PEST CONTROL PRODUCTS ACT

POISON

DANGER – CORROSIVE TO EYES

Albaugh, LLC
1525 NE 36th Street
Ankeny, IA 50021
USA
1-800-247-8013

For medical or treatment information from exposure to this product, call 1-888-347-6732
(7 days/week, 24-hr).

For 24-hour chemical spill, leak, fire, exposure, or accident response information, call
CHEMTREC toll free at 1-800-424-9300.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes and clothing. Wash concentrate from skin or eyes immediately. After use, wash hands and other exposed skin. Avoid breathing spray mist.

Do not use in residential areas, which are defined as sites where bystanders may be present during or after spraying, including homes, schools, parks, playgrounds, playing fields and public buildings.

PROTECTIVE CLOTHING AND EQUIPMENT:

Handling the concentrate (mixing and loading): wear a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves. Rinse gloves before removal. Wear goggles or face shield during mixing/loading. Remove and wash contaminated clothing before reuse.

Handling the dilute spray solution (during application or repairing or cleaning equipment): wear a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves. Rinse gloves before removal. Remove and wash contaminated clothing before reuse. Aerial applicators and applicators using a closed cab are not required to wear chemical-resistant gloves.

ENVIRONMENTAL PRECAUTIONS:

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Surface runoff:

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

Leaching:

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

FIRST AID:

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then

give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote. Employ supportive care. Treatment should be based on judgement of the physician in response to reactions of the patient. High concentrations of MCPA may cause severe irritation to the eyes. Symptoms of overexposure to MCPA could include slurred speech, twitching, jerking and spasms, drooling, low-blood pressure and unconsciousness. Treat symptomatically.

STORAGE:

Store this product away from food or feed. Store the container tightly closed away from seeds, fertilizer, plants and foodstuffs.

Store at temperatures above 0°C. If subjected to freezing temperatures, warm to at least 5°C and mix thoroughly. **SHAKE WELL BEFORE USING.**

DISPOSAL:

For Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site.

Contact your local distributor/dealer or municipality for the location of the nearest collection site.

Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsing to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. **DO NOT** reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.

DIRECTIONS FOR USE:

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, Albaugh MCPA Amine 600 Liquid Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Albaugh MCPA Amine 600 Liquid Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Albaugh MCPA Amine 600 Liquid Herbicide or other Group 4 herbicides within a growing season or among growing seasons with different herbicides groups that control the same weed in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Albaugh at 1-800-247-8013.

GENERAL USE PRECAUTIONS:

Albaugh MCPA Amine 600 Liquid Herbicide may cause damage to susceptible crop, ornamental, and other plants, even in minute quantities.

Avoid spray drift to any desirable vegetation. Coarse sprays are less likely to drift. Do not spray during periods of high winds.

- Do not permit lactating dairy animals to graze fields within 7 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.
- Do not harvest forage or cut hay within 7 days after application.

MIXING:

Shake well before using. To ensure even mixing, half fill the sprayer tank with clean water, add the required amount of Albaugh MCPA Amine 600 Liquid Herbicide and agitate thoroughly. Add the remainder of the water and agitate before spraying. If the spray solution has been left standing, agitate thoroughly before use.

EQUIPMENT AND SPRAY VOLUME:

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wingspan or rotorspan.

Spray Buffer Zones

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats, (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop	Spray Buffer Zones (metres) Required for the Protection of:					
		Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial Habitat	
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m		
Field Sprayer	All crops	1	1	1	1	4	
Aerial	Terrestrial Food and Feed Crops						
	Cereals, flax,	Fixed	1	0	1	0	60
		Rotary wing	1	0	1	0	50
	Pasture, rangeland	Fixed	5	1	1	1	100
		Rotary wing	4	1	1	1	80
	Legumes	Fixed	1	1	0	0	25
Rotary wing		1	1	0	0	25	
	Non-crop Uses						
	Uncropped land	Fixed	30	1	20	1	200
Rotary wing		20	1	10	1	100	

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

Generic Aerial Application Label Instructions

Apply only by fixed-wing or rotary aircraft equipment, which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices. No human flaggers are permitted.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Mixer/loader and applicator must be different individuals. Aircraft must be closed cab. Loading of premixed chemicals with a closed system is permitted. It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear long-sleeved shirt, long pants and coveralls, chemical resistant gloves and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-247-8013 or obtain technical advice from the distributor or your provincial agricultural representative.

Application of this specific product must meet and/or conform to the following:

Apply in 50 to 200 litres of water per hectare by ground equipment unless otherwise indicated. Higher water volumes will reduce the risk of crop injury.

Apply in a minimum of 30 litres of water per hectare by aerial equipment.

Use spray nozzles and pressures that deliver a medium to coarse quality spray.

Avoid generating fine, driftable droplets.

TANK MIXES:

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Albaugh LLC at 1-800-247-8013 for information before applying any tank mix that is not specifically recommended on this label.

CROPS, TIMING, RATES AND METHODS OF APPLICATION:

Use the rate required to control the weeds present.

Apply in warm weather when the crop and weeds are growing well, and the weeds are at a susceptible stage.

Application must be made before the crop canopy shields the weeds.

To minimize the risk of crop injury, do not exceed the recommended rate listed for the crop.

Higher rates may be required to control certain weed species.

Rates above those recommended for a crop may result in significant crop injury, and should only

be used when the risk of crop injury will be offset by the benefits of enhanced weed control.

Avoid application during drought conditions or during exceptionally hot weather.

CROP:

Spring Cereals (Wheat, Barley, Rye, Oats), not underseeded with legumes

APPLICATION TIME:

From the 2-leaf expanded to the early flag-leaf stage.

Do not apply after the flag leaf stage.

Treatment at the 3 to 4-leaf stage usually combines maximum crop tolerance with maximum weed susceptibility.

Do not apply more than one treatment per year.

METHOD:

Ground or aerial application.

RECOMMENDED RATE PER HECTARE:

Up to 1.05 L for wheat, barley and rye.

Up to 900 mL for oats.

WATER VOLUME PER HECTARE:

Minimum of 50 L (ground).

Minimum of 30 L (aerial).

Spraying with higher volumes of water (100-150 L) may reduce risk of crop injury.

CROP:

Winter Cereals (Winter Wheat, Fall Rye)

APPLICATION TIME:

In spring, from full tillering to the flag-leaf stage.

Do not apply during and after the flag-leaf stage.

Do not apply to seedling winter cereals in the fall.

Do not apply more than one treatment per year.

METHOD:

Ground or aerial application.

RECOMMENDED RATE PER HECTARE:

Up to 1.05 L.

WATER VOLUME PER HECTARE:

Minimum of 50 L (ground).

Minimum of 30 L (aerial).

Spraying with higher volumes of water (100-150 L) may reduce risk of crop injury.

CROP:

Spring Cereals, underseeded with alfalfa or bird's-foot trefoil

APPLICATION TIME:

From the 3-leaf expanded to the early flag-leaf stage (cereals).

From the 1st to 4th trifoliolate and not more than 10 cm tall for the seedling legumes.

Do not apply after the flag-leaf stage of the cereals.

METHOD:

Ground application only.

Do not apply more than one treatment per year.

RECOMMENDED RATE PER HECTARE:

Between 250 mL and 470 mL (Do not exceed 470 mL.)

WATER VOLUME PER HECTARE:

Spraying with higher volumes of water (100-150 L) may reduce risk of crop injury.

NOTES:

Some injury to the legumes may occur, especially if growing conditions are very favourable.

To minimize crop injury, spray in the evening and use higher water volumes.

Do NOT apply Albaugh MCPA Amine 600 liquid herbicide to sweet clover.

For cereals underseeded to clovers, the use of TROPOTOX[®] PLUS 400 is recommended.

TANK MIXTURE:

Albaugh MCPA Amine 600 liquid herbicide at 60 mL with EMBUTOX[®] Liquid Herbicide at 1.25 L per hectare may be applied.

CROP:

Flax (Do not apply to low linolenic acid varieties.)

APPLICATION TIME:

When flax is between 5 cm in height to before bud stage.

To minimize crop injury, spray at early growth stages, in the evening, and use higher water volumes.

Do not apply more than one treatment per year.

METHOD:

Ground or aerial application.

RECOMMENDED RATE PER HECTARE:

Up to 830 mL.

RECOMMENDED WATER VOLUME PER HECTARE:

Minimum of 50 L (ground).

Minimum of 30 L (aerial).

Spraying with higher volumes of water (100-150 L) may reduce risk of crop injury.

NOTES:

Caution: Early crop injury may be observed, but yield should not be affected.

CROP:

Field Corn

APPLICATION TIME:

Before corn reaches 15cm tall (leaf extended) and/or before the 6-leaf stage as an overall spray.

Do not apply more than one treatment per year. When hand-harvesting corn (field and sweet), follow a Re-entry Interval (REI) and Pre-harvest Interval (PHI) of 15 days.

METHOD:

Ground application only.

RECOMMENDED RATE PER HECTARE:

Up to 917 mL. Do not add oil or surfactant

RECOMMENDED WATER VOLUME PER HECTARE:

Minimum of 100 L.

NOTES:

Damage to the corn may result if an overall spray is applied beyond this stage.

CROP:

Peas (field and processing)

APPLICATION TIME:

When vines are 10 to 20 cm long but before flower buds are formed.

METHOD:

Ground application only.

RECOMMENDED RATE PER HECTARE:

Between 250 and 417 mL. (Do not exceed 417 mL.)

RECOMMENDED WATER VOLUME PER HECTARE:

150 - 200 L.

NOTES:

DO NOT apply to peas in Nova Scotia.

CROP:

Established Legumes and Pasture (dormant treatment)

APPLICATION TIME:

Apply BEFORE legumes start their growth in the spring.

Do not apply more than two treatments per year, with a minimum retreatment interval of 90 days.

METHOD:

Ground or aerial application.

RECOMMENDED RATE PER HECTARE:

Up to 1.8 L.

RECOMMENDED WATER VOLUME PER HECTARE:

Minimum of 50 L (ground).

Minimum of 30 L (aerial).

NOTES:

Do not graze livestock for at least 7 days after application.

Do NOT apply Albaugh MCPA Amine 600 liquid herbicide to sweet clover.

CROP:

Pasture (without legumes) and Uncropped Land (Rangeland) for tall buttercup control

APPLICATION TIME:

Apply when tall buttercup are in the pre-bud to early bud stage.

Do not apply more than two treatments per year, with a minimum retreatment interval of 90 days.

METHOD:

Ground or aerial application.

RECOMMENDED RATE PER HECTARE:

2.25 to 3.5 L for uncropped land and 2.25 to 2.9 L for pastures.

RECOMMENDED WATER VOLUME PER HECTARE:

Minimum of 50 L (ground).

Minimum of 30 L (aerial).

CROP:

Seedling and Established Red Clover for Seed – Western Canada Only.

APPLICATION TIME:

Seedling – 1st to 3rd trifoliate stage.

Established – breaking of dormancy in spring, up to 7.5 cm.

Do NOT graze or cut for green feed.

Do not apply more than two treatments per year, with a minimum retreatment interval of 21 days.

METHOD:

Ground application only.

RECOMMENDED RATE PER HECTARE:

470 mL.

RECOMMENDED WATER VOLUME PER HECTARE:

Spraying with higher volumes of water (100-150 L) may reduce risk of crop injury. Some crop injury may occur with low water volumes.

CROP:

Seedling Red Clover for Forage.

APPLICATION TIME:

Seedling – 1st to 3rd trifoliate stage.

Do NOT graze or cut for green feed in the first year.

Do not apply more than 1 treatment per year.

METHOD:

Ground application only.

RECOMMENDED RATE PER HECTARE:

470 mL.

RECOMMENDED WATER VOLUME PER HECTARE:

Spraying with higher volumes of water (100-150 L) may reduce risk of crop injury. Some crop injury may occur with low water volumes.

WEEDS CONTROLLED:

(DO NOT exceed the rates given for each specific crop identified in the CROPS, TIMING, RATES AND METHODS OF APPLICATION section)

SUSCEPTIBLE WEEDS (2-4 leaf stage):

Bluebur, burdock (before 4-leaf stage), cocklebur, false flax, flixweed, lamb's-quarters, mustards (except dog and tansy), prickly lettuce, ragweeds, redroot pigweed, Russian pigweed, shepherd's-purse, stinkweed (field pennycress), vetch, wild radish, wild sunflower.

RATES TO CONTROL SUSCEPTIBLE WEEDS:

Small seedlings, good growing conditions: 580 mL/ha.

Large weeds, dry or cold weather, heavy infestations: 920 mL/ha.

Resistance increases with age.

HARDER-TO-CONTROL WEEDS (2-4 leaf stage):

Weeds listed under susceptible plus annual sow-thistle, biennial wormwood, common chickweed, curled dock (before the 4-leaf stage), daisy fleabane, goat's-beard, groundsel, hairy galinsoga, hawkweed, heal-all, hempnettle (before the 4-leaf stage), knotweed (before the 4-leaf stage), kochia, oak-leaved goosefoot, peppergrass, pineapple weed, plantain, purslane, sheep sorrel, smartweed, stinging nettle, velvetleaf, wild radish, yellow rocket (before the 4-leaf stage).

RATES FOR HARDER-TO-CONTROL WEEDS:

Small seedlings, good growing conditions: 1.05 L/ha.

Large weeds, dry or cold weather, heavy infestations: 1.80 L/ha.
Resistance increases with age.

TOP GROWTH CONTROL ONLY:

Blue lettuce, bull thistle, burdock, buttercup, Canada thistle, chicory, curled dock, dandelion, dog mustard, field bindweed, gumweed, hedge bindweed, hoary cress, leafy spurge, mouse-eared chickweed, perennial sow-thistle, Tartary buckwheat, teasel, wild buckwheat, wormwood, yellow rocket.

RATES FOR TOP GROWTH CONTROL:

Small seedlings, good growing conditions: 1.05 L/ha.
Large weeds, dry or cold weather, heavy infestations: 1.80 L/ha.
Resistance increases with age.

CLEANING OF SPRAYER:

Thoroughly clean the herbicide sprayer before applying any other chemical on crops which may be damaged (see USE PRECAUTIONS). Wash the outside of the sprayer and drain the tank completely. Remove and clean filters, screens and nozzle tips separately. Fill the sprayer tank with clean water. With the agitator running, flush out the lines and boom, then drain. Fill the sprayer with clean water and for each 100 L, add 1 L household ammonia. Operate the pump and agitator for 15 minutes. If possible, let the solution remain in the tank and hoses overnight; recirculate and flush out the lines and boom, then drain. Rinse out twice with clean water, recirculating and draining each time. DO NOT contaminate irrigation or drinking water when cleaning the sprayer.

Embutox and Tropotox are registered trademarks of Nufarm Agriculture Inc.