1 2 2	<u>Advantage™ II Kitten</u> <u>Insert (English)</u>
3 4 5	<elanco logo=""> Advantage[™] II kitten</elanco>
5 6 7	Flea Adulticide/Larvicide/Ovicide
, 8 9	For cats 8 weeks and older weighing 2.3 kg or less
10	Once-A-Month Topical Flea Treatment
11 12	Liquid
13	
14	READ THE LABEL AND INSERT BEFORE USING
15 16	FOR USE ONLY ON CATS
17	
18	Do not use on other animals
19 20	
20 21	KEEP OUT OF REACH OF CHILDREN
21	
22	\vee
23	WARNING POISON
24 25	EYE IRRITANT
25 26	
27	DOMESTIC
28	
29	ACTIVE INGREDIENT: Imidacloprid 9.1% and Pyriproxyfen 0.46%
30 31	REG. NO. 31517 P.C.P. ACT
31 32	REG: NO. 31317 F.C.F. ACT
33	Contents: One to six 0.23 mL Tubes
34	
35	Elanco Canada Limited
36	1919 Minnesota Court, Suite 401
37 38	Mississauga, Ontario L5N 0C9 1-800-265-5475
38 39	1-000-200-0470
40	
41	

- 42 Advantage II kitten kills 98-100% of the existing fleas on cats within 1 day after
- 43 treatment.
- 44 Protection against further flea infestation can last up to 4 weeks.
- 45 Retreatment at monthly intervals is required if re-infestation is continuous or if flea
- 46 control is desired. Fleas, eggs, and larvae are killed following contact with skin debris
- 47 from an Advantage II kitten treated pet. Advantage II kitten provides multi-stage flea
- 48 control effectively breaking all flea life-cycle stages for quick and lasting control of flea
- 49 populations. Do not shampoo or immerse the animal in water immediately after
- 50 treatment. The animal can be shampooed, immersed in water or exposed to rain 3 days
- 51 after treatment. Advantage II kitten retains its ovicidal and larvicidal efficacy up to 4
- 52 weeks following a shampoo treatment, swimming or exposure to rain. The adulticidal
- effect may be shortened by repeated shampooing, water immersion or exposure to rain.
 Reapplication may be required after such activities. The successive feeding activity of
- 55 fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis
- 56 (FAD). Treatment of pets with Advantage II kitten rapidly kills fleas and may reduce the
- 57 incidence of this condition.
- 58 For effective flea control, treatment of the pet should be combined with sanitation of any
- area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after
- 60 use) and wash the pet's bedding, living quarters and surrounding areas. If pest
- 61 problems persist, an insecticidal premise treatment may be required. A veterinarian
- 62 should be consulted when pest infestation continues to be a problem on the pet.
- 63
- 64 PRECAUTIONS:
- 65
- 66 HUMANS: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed. Causes 67 eye irritation. DO NOT get in eyes. Avoid contact with skin or clothing. Wear rubber
- 68 gloves during application of this product. After handling or applying, wash hands (or any
- 69 other skin that came in contact with the product), with soap and water. Avoid contact
- 70 with pets until dry.
- 71
- PETS: Do not use on animals under 8 weeks of age. Care should be taken to prevent
 exposure of other pets who may be in close physical contact with the treated animal.
 Sensitivities may occur after using ANY pesticide product for pets. Smaller animals are
 more likely to experience an adverse reaction. Observe carefully. If signs persist, or
 become more severe, consult a veterinarian immediately.
- 77

Side Effects: Monitor your cat after application. Side effects may include: skin irritation such as scratching or hair loss at the application site, or changes in behaviour such as agitation or lethargy. Gastrointestinal effects such as drooling, vomiting or loss of appetite may also occur. If these or other side effects occur, consult your veterinarian or Elanco at 1-800-265-5475.

- 78
- 79
- 80

- 81 FIRST AID:
- 82

HUMANS: IF IN EYES, hold eye open and rinse slowly and gently with water for 15–20
minutes. Remove contact lenses, if present, after the first 5 minutes, then continue
rinsing eye. Call a poison control centre or doctor for treatment advice.

- 86 IF SWALLOWED, call a poison control centre or doctor immediately for treatment
- 87 advice. Have person sip a glass of water if able to swallow. Do not induce vomiting
- unless told to do so by a poison control centre or doctor. Do not give anything by mouth
- to an unconscious person. IF ON SKIN OR CLOTHING, take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control
- 91 centre or doctor for treatment advice.
- 92
- 93 PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large
- 94 amounts of water. If signs continue, consult a veterinarian immediately.
- 95 Take container, label or product name and Pest Control Product Registration Number
- 96 with you when seeking medical or veterinary attention.
- 97
- 98 TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.
- 99
- 100 STORAGE: Store container at room temperature.
- 101
- 102 DISPOSAL: Wrap empty tube and cap in newspaper and dispose in household
- 103 garbage.
- 104
- 105 DIRECTIONS FOR USE:
- 106 This product contains imidacloprid and pyriproxyfen. Do not apply another pest control
- 107 product such as a shampoo, collar, or powder that contains imidacloprid or pyriproxyfen
- 108 to the treated animal after applying Advantage II Kitten.
- 109
- 110 Do NOT administer by mouth.
- 111
- 112 Use only on cats 8 weeks of age and older weighing 2.3 kg or less. Do not use on cats 113 under 8 weeks of age.
- 114
- 115 This product is packaged in single dosage tubes. This package contains one to six
- 116 monthly 0.23 mL dosages. Do not apply a dosage higher than one 0.23 mL tube at one 117 time. Treat your cat monthly in the manner described below.
- 118 1. Use only on kittens and cats 8 weeks of age and older weighing 2.3 kg or less.
- 119 Do not use on cats under 8 weeks of age. Do not use on other animals.
- 120 2. Remove one applicator from the package.
- 121 3. Hold applicator tube in upright position. Pull cap off tube.
- 122 4. Turn cap around and place other end of cap back on tube.
- 123 5. Twist cap to break seal, then remove cap from tube.

- 6. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to completely empty the contents directly on the skin. Do not get this product in your pet's eyes or mouth. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product.
- 130 7. Discard the empty tube as described in "DISPOSAL".
- 131

Consult a veterinarian before using on sick, aged, pregnant or nursing animals or
 animals receiving drug or other pesticide treatment. A veterinarian should be consulted
 when pest infestation continues to be a problem on the pet.

- Repeat every month for continual pest control in situations where there is an ongoing risk of pest infestation.
- 137
- 138 Until Advantage II kitten dries, keep pet off furniture and away from hardwood floors.
- 139
- 140 Retain packaging information for future reference
- 141
- 142 Advantage, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.
- 143 ©2022 Elanco or its affiliates.
- 144

145 146	<u>Advantage™ II Kitten</u> <u>Carton (English)</u>
140 147	
147	<elanco logo=""> Advantage[™] II kitten</elanco>
148	Advantage in kitten
150	Flea Adulticide/Larvicide/Ovicide
151	
152	For cats 8 weeks and older weighing 2.3 kg or less
153	
154	Once-A-Month Topical Flea Treatment
155	
156	Liquid
157	READ THE LABEL AND INSERT BEFORE USING
158 159	READ THE LABEL AND INSERT BEFORE USING
160	FOR USE ONLY ON CATS
161	
162	Do not use on other animals
163	
164	KEEP OUT OF REACH OF CHILDREN
165	
166	V
167	WARNING POISON
168	
169	EYE IRRITANT
170	DOMESTIC
171	DOMESTIC
172	ACTIVE INGREDIENT: Imidacloprid 9.1% and Pyriproxyfen 0.46%
173 174	ACTIVE INGREDIENT. IIIIdaciophu 9.1% and Fynproxyten 0.40%
174	REG. NO. 31517 P.C.P. ACT
176	
177	Contents: One to six 0.23 mL Tubes
178	
179	Elanco Canada Limited
180	1919 Minnesota Court, Suite 401
181	Mississauga, Ontario L5N 0C9
182	1-800-265-5475
183	
184	
185	

- 186 Advantage II kitten kills 98-100% of the existing fleas on cats within 1 day after
- 187 treatment.
- 188 Protection against further flea infestation can last up to 4 weeks.
- 189 Retreatment at monthly intervals is required if re-infestation is continuous or if flea
- 190 control is desired. Fleas, eggs, and larvae are killed following contact with skin debris
- 191 from an Advantage II kitten treated pet. Advantage II kitten provides multi-stage flea
- 192 control effectively breaking all flea life-cycle stages for quick and lasting control of flea
- 193 populations. Do not shampoo or immerse the animal in water immediately after
- 194 treatment. The animal can be shampooed, immersed in water or exposed to rain 3 days
- after treatment. Advantage II kitten retains its ovicidal and larvicidal efficacy up to 4
- 196 weeks following a shampoo treatment, swimming or exposure to rain. The adulticidal
- 197 effect may be shortened by repeated shampooing, water immersion or exposure to rain.
- 198 Reapplication may be required after such activities.
- 199
- 200 PRECAUTIONS:
- 201 HUMANS: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed. Causes
- 202 eye irritation. DO NOT get in eyes. Avoid contact with skin or clothing. Wear rubber
- 203 gloves during application of this product. After handling or applying, wash hands (or any
- other skin that came in contact with the product), with soap and water. Avoid contact
- with pets until dry.
- 206 PETS: Do not use on animals under 8 weeks of age. Care should be taken to prevent
- 207 exposure of other pets who may be in close physical contact with the treated animal.
- 208 Sensitivities may occur after using ANY pesticide product for pets. Smaller animals are
- 209 more likely to experience an adverse reaction. Observe carefully. If signs persist, or
- 210 become more severe, consult a veterinarian immediately.
- 211

Side Effects: Monitor your cat after application. Side effects may include: skin irritation such as scratching or hair loss at the application site, or changes in behaviour such as agitation or lethargy. Gastrointestinal effects such as drooling, vomiting or loss of appetite may also occur. If these or other side effects occur, consult your veterinarian or Elanco at 1-800-265-5475.

- 212
- 213 FIRST AID:
- HUMANS: IF IN EYES, hold eye open and rinse slowly and gently with water for 15–20
- 215 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue 216 rinsing eve
- 216 rinsing eye.
- 217 Call a poison control centre or doctor for treatment advice.
- 218 IF SWALLOWED, call a poison control centre or doctor immediately for treatment
- advice. Have person sip a glass of water if able to swallow. Do not induce vomiting
- unless told to do so by a poison control centre or doctor. Do not give anything by mouth
- to an unconscious person. IF ON SKIN OR CLOTHING, take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control
- 223 centre or doctor for treatment advice. PETS: If signs of sensitivity occur, bathe the pet

- with mild soap and rinse with large amounts of water. If signs continue, consult a
- 225 veterinarian immediately.
- Take container, label or product name and Pest Control Product Registration Number with you when seeking medical or veterinary attention.
- 229 TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.
- 230

228

- 231 STORAGE: Store container at room temperature.
- 232
- DISPOSAL: Wrap empty tube and cap in newspaper and dispose in household garbage.
- 235

236 237	<u>Advantage™ II Kitten</u> <u>Tube (English)</u>
238	<elanco logo=""></elanco>
239 240	Advantage [™] II kitten
240	READ THE LABEL AND INSERT BEFORE USING
242	
243	\vee
244	WARNING POISON
245	
246	EYE IRRITANT
247	
248	DOMESTIC
249	
250	EXTERNAL USE ONLY
251	
252	ACTIVE INGREDIENT: Imidacloprid 9.1% and Pyriproxyfen 0.46%
253	
254	REG. NO. 31517 P.C.P. ACT
255	Flance Canada Limitad
256	Elanco Canada Limited
257	Mississauga, Ontario L5N 0C9
258	
259 260	0.23 mL
260 261	0.23 IIIL
201	