

2017-6981
2018-01-18

PRINCIPAL DISPLAY PANEL

GROUP

2

HERBICIDE

PP-2525 TOSS-N-GO® HERBICIDE

Soluble Granule

FOR SALE FOR USE ON WHEAT (SPRING, DURUM AND WINTER), SPRING BARLEY AND
OATS

AGRICULTURAL

REGISTRATION NO. 31733 PEST CONTROL PRODUCTS ACT

GUARANTEE: Thifensulfuron-methyl 25%
Tribenuron-methyl 25%

Warning: Contains the allergen milk, sulfites

POTENTIAL SKIN SENSITIZER

NET CONTENTS: 121.5 – 972 grams
(1 x 121.5 gram or 4 x 121.5 gram or 8 x 121.5 gram water soluble bags)

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104
1-833-362-7722

DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.

PRECAUTIONS:

- POTENTIAL SKIN SENSITIZER
- KEEP OUT OF REACH OF CHILDREN.
- AVOID BREATHING SPRAY MIST. USE WITH ADEQUATE VENTILATION.
- AVOID CONTACT WITH SKIN, EYES AND CLOTHING. WEAR GOGGLES OR FACE SHIELD, LONG-SLEEVED SHIRT, LONG PANTS, SOCKS, COVERALLS, AND CHEMICAL RESISTANT GLOVES AND FOOTWEAR DURING MIXING, LOADING, APPLICATION CLEAN UP AND REPAIR. CHEMICAL RESISTANT GLOVES ARE NOT REQUIRED WHILE OPERATING GROUNDBOOM SPRAYERS
- DO NOT CONTAMINATE ANY BODY OF WATER.
- MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. CAUSES EYE AND SKIN IRRITATION.
- WHEN USING, DO NOT EAT, DRINK OR SMOKE. WASH SKIN THOROUGHLY WITH SOAP AND WATER AFTER HANDLING.
- REMOVE CONTAMINATED CLOTHING IMMEDIATELY AFTER USE. STORE AND WASH CONTAMINATED CLOTHING SEPARATELY FROM HOUSEHOLD LAUNDRY BEFORE REUSE.
- DO NOT SELL WATER SOLUBLE BAGS SEPERATELY
- DO NOT USE HUMAN FLAGGERS
- DO NOT ENTER OR ALLOW WORKER RE-ENTRY FOR 12 HOURS AFTER APPLICATION.

FIRST AID:**IF IN EYES:**

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or the product name and Pest Control Product Registration Number with you when seeking medical attention.

IN CASE OF A MEDICAL EMERGENCY, CALL TOLL FREE, DAY OR NIGHT: 1-800-331-3148

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

DISPOSAL:

If this is a cardboard container:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

If this is a refillable container:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

If this is a returnable container:

Do not reuse this container for any other purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For all package types:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

Water Soluble Bag Label

PP-2525 TOSS-N-GO® HERBICIDE

KEEP OUT OF REACH OF CHILDREN

CAUTION: SEE DIRECTIONS FOR USE
FOR COMPLETE PRECAUTIONARY STATEMENTS

GUARANTEE:	Thifensulfuron methyl	25%
	Tribenuron methyl	25%

SOLUBLE GRANULES

NET CONTENTS: 486 GRAMS

THIS BAG TREATS 4 HECTARES (10 ACRES)

PP-2525 TOSS-N-GO® HERBICIDE

Soluble Granule

FOR SALE FOR USE ON WHEAT (SPRING, DURUM AND WINTER), SPRING BARLEY AND OATS .

AGRICULTURAL

REGISTRATION NO. 31733 PEST CONTROL PRODUCTS ACT

GUARANTEE:	Thifensulfuron-methyl	25%
	Tribenuron-methyl	25%

Warning: Contains the allergen milk, sulfites

POTENTIAL SKIN SENSITIZER

READ THE LABEL AND THIS ATTACHED BOOKLET BEFORE USING

FMC Corporation

2929 Walnut Street

Philadelphia, PA 19104

1-833-362-7722

DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.

PRECAUTIONS:

- POTENTIAL SKIN SENSITIZER
- KEEP OUT OF REACH OF CHILDREN.
- AVOID BREATHING SPRAY MIST. USE WITH ADEQUATE VENTILATION.
- AVOID CONTACT WITH SKIN, EYES AND CLOTHING. WEAR GOGGLES OR FACE SHIELD, LONG-SLEEVED SHIRT, LONG PANTS, SOCKS, COVERALLS, AND CHEMICAL RESISTANT GLOVES AND FOOTWEAR DURING MIXING, LOADING, APPLICATION CLEAN UP AND REPAIR. CHEMICAL RESISTANT GLOVES ARE NOT REQUIRED WHILE OPERATING GROUNDBOOM SPRAYERS.
- DO NOT CONTAMINATE ANY BODY OF WATER.
- MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. CAUSES EYE AND SKIN IRRITATION.
- WHEN USING, DO NOT EAT, DRINK OR SMOKE. WASH SKIN THOROUGHLY WITH SOAP AND WATER AFTER HANDLING.
- REMOVE CONTAMINATED CLOTHING IMMEDIATELY AFTER USE. STORE AND WASH CONTAMINATED CLOTHING SEPARATELY FROM HOUSEHOLD LAUNDRY BEFORE REUSE.
- DO NOT SELL WATER SOLUBLE BAGS SEPARATELY
- DO NOT USE HUMAN FLAGGERS
- DO NOT ENTER OR ALLOW WORKER RE-ENTRY FOR 12 HOURS AFTER APPLICATION.

FIRST AID:**IF IN EYES:**

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or the product name and Pest Control Product Registration Number with you when seeking medical attention.

IN CASE OF A MEDICAL EMERGENCY, CALL TOLL FREE, DAY OR NIGHT: 1-800-331-3148

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS:

- Overspray or drift to important wildlife habitats such as shelterbelts, wetlands, sloughs or dry slough borders and woodlots should be avoided. Observe buffer zones specified under DIRECTIONS FOR USE.
- Do not remove soil from treated fields for use on lawns, in gardens or for back fill.
- Do not contaminate irrigation water. Do not apply to land where tail water will be used to irrigate cropland.
- Do not apply more than 30 g per hectare of PP-2525 Toss-N-Go® Herbicide per crop year.
- TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted or fine textured such as clay).
- Avoid application of this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

TABLE OF CONTENTS:

.....	1
PRINCIPAL DISPLAY PANEL	1
PRECAUTIONS:	2
FIRST AID:	2
<i>IF IN EYES:</i>	2
<i>IF ON SKIN OR CLOTHING:</i>	2
<i>IF SWALLOWED:</i>	2
<i>IF INHALED:</i>	2
TOXICOLOGICAL INFORMATION:.....	2
STORAGE:	3
DISPOSAL:	3
NOTICE TO USER:	3
DFU BOOKLET	5
PRECAUTIONS:	6
FIRST AID.....	6
<i>IF IN EYES:</i>	6
<i>IF ON SKIN OR CLOTHING:</i>	6
<i>IF SWALLOWED:</i>	6
<i>IF INHALED:</i>	6
TOXICOLOGICAL INFORMATION:.....	7
ENVIRONMENTAL PRECAUTIONS:	7
PRODUCT DESCRIPTION:	10
<i>Minimum Crop Rotation Guidelines:</i>	11
GRAZING/HARVEST RESTRICTIONS:.....	11
VARIETY RESTRICTIONS:.....	11
RAINFASTNESS INFORMATION:	11
APPLICATION INFORMATION:	11
APPLICATION PRECAUTIONS:.....	11
APPLICATION INSTRUCTIONS:	12
DIRECTIONS FOR USE:	13
BUFFER ZONES:	13
WHEAT (SPRING, DURUM AND WINTER) AND SPRING BARLEY	15
<i>PP-2525 Toss-N-Go® HERBICIDE</i>	15
<i>PP-2525 Toss-N-Go® Herbicide + MCPA ESTER</i>	16
<i>PP-2525 Toss-N-Go® Herbicide + 2,4-D ESTER</i>	16
WHEAT (SPRING, DURUM) AND SPRING BARLEY	17
<i>PP-2525 Toss-N-Go® Herbicide + Perimeter™ Herbicide OR Perimeter™ Mega PrecisionPac® Herbicide</i>	17
<i>PP-2525 Toss-N-Go® Herbicide + Perimeter™ II Herbicide</i>	17
<i>PP-2525 Toss-N-Go® Herbicide + PUMA*¹²⁰ SUPER</i>	18
<i>PP-2525 Toss-N-Go® Herbicide + PUMA*¹²⁰ SUPER + 2,4-D Ester</i>	18
<i>PP-2525 Toss-N-Go® Herbicide + BANVEL* II or BANVEL* Dry EG Herbicide</i>	19
<i>PP-2525 Toss-N-Go® Herbicide + 2,4-D ESTER + BANVEL* II or BANVEL* Dry EG Herbicide</i>	19
SPRING WHEAT (EXCLUDING DURUM) AND SPRING BARLEY.....	21
<i>PP-2525 Toss-N-Go® HERBICIDE + AXIAL* 100EC Herbicide</i>	21
FOR WEED CONTROL IN SPRING WHEAT AND DURUM WHEAT	22
<i>PP-2525 Toss-N-Go® Herbicide + HARMONY® GRASS 128EC Herbicide</i>	22
<i>PP-2525 Toss-N-Go® Herbicide + HORIZON* NG Herbicide</i>	22
<i>PP-2525 Toss-N-Go® Herbicide + Horizon* 240 EC Herbicide Tank Mix</i>	23
<i>PP-2525 Toss-N-Go® Herbicide + SIMPLICITY*</i>	23
FOR WEED CONTROL IN SPRING WHEAT ONLY.....	24
<i>PP-2525 Toss-N-Go® Herbicide+ 2,4-D ESTER + EVEREST* 70 DF OR EVEREST* SOLUPAK 70</i> .	24

MIXING INFORMATION:.....	25
<i>MIXING PRECAUTIONS:</i>	25
<i>MIXING DIRECTIONS:</i>	25
<i>SPECIAL MIXING INSTRUCTIONS FOR CHEMICAL HANDLERS</i>	25
EQUIPMENT CLEANING INFORMATION:	26
<i>CLEANING PRECAUTIONS:</i>	26
<i>CLEANING INSTRUCTIONS:</i>	26
RESISTANCE-MANAGEMENT RECOMMENDATIONS:	27
STORAGE:	27
DISPOSAL:	28
NOTICE TO USER:	28

PRODUCT DESCRIPTION:

PP-2525 Toss-N-Go® Herbicide is recommended for selective post-emergent control or suppression of broadleaf weeds in wheat (spring, durum and winter), spring barley and oats not underseeded to legumes or grasses, and in certain grasses for forage or seed production.

PP-2525 Toss-N-Go® Herbicide is a 50% active ingredient herbicide formulated as a water soluble granule. It is non-corrosive, non-flammable, non-volatile and does not freeze.

PP-2525 Toss-N-Go® Herbicide is to be mixed with water and a recommended adjuvant and applied as a uniform broadcast spray. Recommended adjuvants are: AGRAL* 90, AG-SURF* and CITOWETT* PLUS. Apply early postemergence, to the main flush of actively growing broadleaf weeds.

Warm, moist growing conditions promote active weed growth and enhance the activity of PP-2525 Toss-N-Go® Herbicide by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and regrowth may occur. For best results, ensure thorough spray coverage of target weeds. See "DIRECTIONS FOR USE" section of this label for complete use details.

PP-2525 Toss-N-Go® Herbicide stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

IMPORTANT INFORMATION: READ BEFORE USING

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

- Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, recreational grounds or similar areas.
- Prevent drift of spray to desirable plants.
- Do not contaminate any body of water, including irrigation water that may be used on other crops.
- Carefully observe sprayer clean-up instructions, as spray tank residues may damage crops other than wheat and spring barley.

REPLANTING TO OTHER CROPS:

Minimum Crop Rotation Guidelines:

Minimum Interval	Crop
2 months	Canola, Flax, Lentils, Alfalfa
Next crop year	All crops

GRAZING/HARVEST RESTRICTIONS:

Wheat, spring barley or oats may not be grazed or fed to livestock within 7 days of application.

VARIETY RESTRICTIONS:

Limit use of PP-2525 Toss-N-Go® Herbicide to currently licensed varieties of wheat (spring, durum and winter), spring barley and oats. Consult FMC Corporation prior to use on unlicensed varieties for crop tolerance information. **DO NOT USE ON LEGER SPRING BARLEY OR BELVEDERE WHEAT VARIETIES.**

RAINFALL INFORMATION:

Up to 25 mm of rain beginning 1 hour or more after spraying will not reduce the effectiveness of PP-2525 Toss-N-Go® Herbicide.

APPLICATION INFORMATION:

APPLICATION PRECAUTIONS:

- Do not apply to irrigated land where tail water will be used to irrigate cropland.
- Overspray or drift to important wildlife habitats such as shelterbelts, wetlands, sloughs or dry slough borders and woodlots should be avoided. Observe buffer zones specified under DIRECTIONS FOR USE.
- Apply with an adjuvant registered for use with PP-2525 Toss-N-Go® Herbicide. Use only water as a carrier. Other carriers may accelerate breakdown of PP-2525 Toss-N-Go® Herbicide and reduce its effectiveness.
- Do not apply to crops underseeded to legumes or grasses as injury to forage may result.
- Application to crops that are stressed by severe weather conditions, frost, low fertility, water logged soil (soil at or near field capacity), disease or insect damage may result in crop injury. Drought, disease or insect damage following application may also result in crop injury.
- Under certain conditions, such as heavy rainfall, prolonged cool weather, frost or wide fluctuations in day/night temperatures, temporary lightening in crop colour, and occasionally, a slight reduction in crop height may occur.
- Do not apply more than 30 grams/hectare of PP-2525 Toss-N-Go® Herbicide per year.
- Apply only when the potential for drift to areas of human habitation or areas of human activity (houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature, application equipment and spray settings.

APPLICATION INSTRUCTIONS:

GROUND APPLICATION

Apply the spray mixture uniformly with properly calibrated ground equipment. Ensure thorough coverage and a uniform spray pattern. Flat fan nozzles are recommended (minimum 55 litres spray volume per hectare). Use 50 mesh or larger filter screens (metal or nylon). A spray pressure of 210-275 kPa is recommended. Avoid overlapping and shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over application.

AERIAL APPLICATION:

Apply only by fixed-wing or rotary aircraft equipment, which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label. Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment. Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions:

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions:

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted. It is desirable that the pilot have communication capabilities at each treatment site at the time of application. The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions:

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-833-362-7722 or obtain technical advice from the distributor or your provincial agricultural representative.

Application of this specific product must meet and/or conform to the following:

Apply a minimum of 25 L of spray volume per hectare.

Apply a maximum of 50 L of spray volume per hectare.

DIRECTIONS FOR USE:

- PP-2525 Toss-N-Go[®] Herbicide is formulated as a soluble granule sealed within a water-soluble package (Toss-N-Go[®] bag). Drop intact Toss-N-Go[®] bag directly into spray tank. The Toss-N-Go[®] bag and pesticide will dissolve readily in water.
- Do not allow the Toss-N-Go[®] bag to become wet prior to use. Do not handle individual Toss-N-Go[®] bag with wet hands or wet gloves as this may cause breakage. Do not remove Toss-N-Go[®] bag from over-wrap container except for immediate use.
- Do not open Toss-N-Go[®] bag or attempt to remove the contents from them. The contents are not designed to be re-measured or subdivided.
- Do not use opened or punctured Toss-N-Go[®] bag.
- If broken Toss-N-Go[®] bag is found when the over-wrap container is opened, avoid contact with, and inhalation of the product. Handle with chemical-resistant gloves and dispose of broken Toss-N-Go[®] bag according to the DISPOSAL section.

Field Sprayer Application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

Buffer zones:

Use of the following spray methods or equipment **DO NOT** require a buffer zone: hand-held or backpack sprayer, inter-row hooded sprayer, spot treatment, soil drench, and soil incorporation.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, pastures, rangelands, riparian areas and shrub lands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop		Buffer Zones (metres) Required for the Protection of:		
			Aquatic Habitats of Depths:		Terrestrial Habitats
			Less than 1 m	Greater than 1 m	
Field sprayer*	Wheat (spring, durum and winter), spring barley, oats		1	0	15
Aerial	Wheat (spring, durum and winter), spring barley, oats	Fixed wing	1	0	125
		Rotary wing	1	0	100

* For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.

NOTE: Applicators may recalculate a site-specific buffer zone by combining information on current weather conditions and spray configuration for the following applications: all airblast applications, and for field and aerial applications which specify the following droplet size category wording on the product label: 'DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium or coarse classification.' To access the Buffer Zone Calculator, please visit the Pest Management Regulatory Agency web site.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

WHEAT (SPRING, DURUM AND WINTER) AND SPRING BARLEY

PP-2525 Toss-N-Go® HERBICIDE	
LOCATION: No restrictions	
CROP (S): Spring wheat (including durum), winter wheat and spring barley	
TIMING: Post-emergent application from the 2-leaf stage to just before the flag leaf (shot blade) stage of wheat (including durum), winter wheat and barley, prior to head emergence, but before the crop canopy will prevent thorough coverage of target weeds. Do not apply after head has emerged. For best results, apply to young, actively growing weeds that are less than 10 cm in height or diameter. Thorough coverage of target weeds is essential.	
APPLICATION RATES	
PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
Adjuvants: Agral 90, Ag-Surf or Citowett * Plus	2 litres/1000 litres spray solution
MIXING: Mixing order is PP-2525 Toss-N-Go® Herbicide, and then add adjuvant while maintaining agitation. Antifoaming agent may be required. Refer to MIXING DIRECTIONS section for general instructions.	
Weeds Controlled	Weeds Suppressed
Annual smartweed (green smartweed, lady's thumb)	Canada thistle ²
Common chickweed ⁴	
Cow cockle	
Flixweed	
Hemp-nettle	
Kochia ³	
Lamb's-quarters	
Narrow-leaved hawk's beard	
Night flowering catchfly	
Redroot Pigweed	
Russian thistle	
Sow thistle (perennial)	
Stinkweed	
Volunteer Rapeseed ¹	
Wild Buckwheat	
Wild mustard	

Notes:

1. PP-2525 Toss-N-Go® Herbicide alone will not control imazethapyr tolerant canola varieties (e.g. Clearfield* varieties or other varieties with the Pursuit Smart* trait)
2. Apply when the majority of thistles have emerged and are actively growing. For best top growth control, apply before bud stage when thistles are no larger than 15 cm in height and before crop canopy prevents thorough coverage of weeds. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.
3. PP-2525 Toss-N-Go® Herbicide alone will not control Group 2 resistant Kochia.
4. Apply when chickweed is small (1-6 leaf) and actively growing, but before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.

PP-2525 Toss-N-Go® Herbicide + MCPA ESTER	
LOCATION: No restrictions	
CROP (S): Spring wheat (including durum), winter wheat, spring barley and oats	
TIMING: Post-emergent application from full 3-leaf to just before flag leaf (shot blade) growth stage of crop.	
TANK MIX COMPONENTS AND APPLICATION RATES	
PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
MCPA Ester 500	0.56 L/ha
Or	Or
MCPA Ester 600	0.47 L/ha
MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is dissolved before adding MCPA Ester. Then add MCPA Ester while continuing agitation. Refer to MIXING DIRECTIONS section for general directions. If antifoam agent is required, add last. Consult the MCPA Ester label for additional application instructions and use precautions.	
Weeds Controlled	
Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide alone, as well as white cockle, dandelion (spring and fall rosettes, up to 15 cm in diameter), scentless chamomile and volunteer imazethapyr tolerant canola varieties (e.g. Clearfield* canola or other varieties with the Pursuit* Smart trait). Apply when volunteer canola is in the 2-4 leaf stage.	

PP-2525 Toss-N-Go® Herbicide + 2,4-D ESTER	
LOCATION: No restriction	
CROP (S): Spring wheat (including durum), winter wheat and spring barley	
TIMING: Post-emergent application from full 3-leaf to just before flag leaf (shot blade) growth stage of crop.	
TANK MIX COMPONENTS AND APPLICATION RATES	
PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
2,4-D Ester 700LV	0.6 L/ha (or 420 g/ha acid equivalent)
Adjuvants: AGRAL 90, AG-SURF or CITOWETT PLUS	2 litres/1000 litres of spray solution
MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is dissolved before adding 2,4-D. Then add 2,4-D while continuing agitation, then adjuvant. Refer to MIXING DIRECTIONS section for general directions. If antifoam agent is required, add last. Consult the 2,4-D label for additional application instructions and use precautions.	
Weeds Controlled	
Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide alone, as well as weeds controlled by 2,4-D Ester.	

WHEAT (SPRING, DURUM) AND SPRING BARLEY

PP-2525 Toss-N-Go® Herbicide + PERIMETER™ HERBICIDE OR PERIMETER™ MEGA PRECISIONPAC® HERBICIDE	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP (S): Spring wheat (including durum) and spring barley	
TIMING: Post-emergent application from 2-leaf to initiation of stem elongation of crop.	
TANK MIX COMPONENTS AND APPLICATION RATES	
PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
PERIMETER™ Herbicide or PERIMETER™ MEGA PRECISIONPAC® HERBICIDE	0.3 L/ha for cleavers and kochia (including Group 2 resistant biotypes) 0.4 L/ha for control of weeds listed above plus volunteer flax and stork's bill and round-leaved mallow
Adjuvants: Agral 90, Ag-Surf or Citowett * Plus	2 litres/1000 litres spray solution
MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is dissolved before adding PERIMETER™ Herbicide or PERIMETER™ MEGA PRECISIONPAC® HERBICIDE. Then add PERIMETER™ Herbicide or PERIMETER™ MEGA PRECISIONPAC® HERBICIDE while continuing agitation, then adjuvant. Refer to MIXING DIRECTIONS section for general directions. If antifoam agent is required, add last. Consult the PERIMETER™ Herbicide or PERIMETER™ MEGA PRECISIONPAC® HERBICIDE label for additional application instructions and use precautions.	
Weeds Controlled	
Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide alone, as well as cleavers (1-4 whorl), kochia (including Group 2 resistant biotypes; seedling to 8-leaf stage) stork's bill (1-6 leaf) and volunteer flax (up to 12 cm in height) and round-leaved mallow (1-5leaf).	

PP-2525 Toss-N-Go® Herbicide + PERIMETER™ II HERBICIDE	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP (S): Spring wheat (including durum) and spring barley	
TIMING: Post-emergent application from 2-leaf to initiation of stem elongation of crop.	
TANK MIX COMPONENTS AND APPLICATION RATES	
PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
PERIMETER™ II Herbicide	0.155 L/ha for control of cleavers and kochia (including Group 2 resistant biotypes) 0.21 L/ha for control of weeds listed above plus volunteer flax and stork's bill (1-6 leaf) and round-leaved mallow
Adjuvants: Agral 90, Ag-Surf or Citowett * Plus	2 litres/1000 litres spray solution
MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is dissolved before adding PERIMETER™ II Herbicide. Then add PERIMETER™ II Herbicide while continuing agitation, then adjuvant. Refer to MIXING DIRECTIONS section for general directions. If antifoam agent is required, add last. Consult the PERIMETER™ II Herbicide label for additional application instructions and use precautions.	

Weeds Controlled

Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide alone, as well as cleavers (1-4 whorl) and kochia (including Group 2 resistant biotypes; seedling to 8-leaf stage), volunteer flax (up to 12 cm in height) and stork's bill (1-6 leaf) and round-leaved mallow (1-5 leaf).

PP-2525 Toss-N-Go® Herbicide + PUMA*¹²⁰ SUPER

LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.

CROP (S): Spring wheat (including durum) and spring barley

TIMING: Post-emergent application from 2 to 6-leaf plus 3 tillers growth stage of spring and durum wheat; 2 to 5-leaf plus 2 tillers growth stage of spring barley. For annual grass weeds, apply from 1 to 6-leaf stage, up to emergence of third tiller.
For green foxtail control, apply at the 1-6 leaf (up to emergence of the 3rd tiller) stage.

TANK MIX COMPONENTS AND APPLICATION RATES

PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
PUMA* ¹²⁰ SUPER	385 ml/hectare for green foxtail only 770 ml/hectare for wild oats, barnyard grass, green foxtail and yellow foxtail.

No adjuvant required.

MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is completely dissolved before adding Puma*¹²⁰ Super. Then add Puma*¹²⁰ Super. Refer to MIXING DIRECTIONS section for general directions. Consult the Puma*¹²⁰ Super label for additional information on appropriate weed staging, application instructions, environmental and use precautions.

WEEDS CONTROLLED

Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide alone, as well as wild oats, green foxtail, barnyard grass and yellow foxtail.

PP-2525 Toss-N-Go® Herbicide + PUMA*¹²⁰ SUPER + 2,4-D Ester

LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.

CROP (S): Spring wheat (including durum) and spring barley

TIMING: Apply this tank mix from the full 3-leaf stage to the 6-leaf + 3 tiller stage of wheat (including durum) and the full 3-leaf stage to the 5-leaf + 2 tiller stage of barley.

For green foxtail control, apply at the 1-6 leaf (up to emergence of the 3rd tiller) stage.

TANK MIX COMPONENTS AND APPLICATION RATES

PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
PUMA* ¹²⁰ SUPER	385 ml/hectare for green foxtail only

Choose one of the following:

2,4-D Ester LV700

0.6 L/ha (or 420 g/ha acid equivalent)

No adjuvant required.

MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is completely dissolved before adding Puma*¹²⁰ Super. Then add Puma*¹²⁰ Super, and, with reduced agitation, add 2,4-D Ester. Refer to MIXING DIRECTIONS section for general directions. Consult the Puma*¹²⁰ Super label for additional information on appropriate weed staging, application instructions, environmental and use precautions.

WEEDS CONTROLLED

Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide plus 2,4-D Ester, and green foxtail.

PP-2525 Toss-N-Go® Herbicide + BANVEL* II or BANVEL* Dry EG Herbicide

LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.

CROP (S): Spring wheat (including durum) and spring barley

TIMING: Apply the tank mix from the 2-5 leaf stage wheat (including durum) and barley, and to kochia plants up to the 8-leaf stage. Expect superior results when kochia is small. Use the higher rate of Banvel II or Banvel Dry EG Herbicide when the majority of Kochia plants are at or near the 8-leaf stage, or when growing conditions are not favourable.

TANK MIX COMPONENTS AND APPLICATION RATES

PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
Banvel* II Herbicide or	110 – 145 mL/ha or
Banvel*Dry EG Herbicide	76 - 100 g/ha
Adjuvants: AGRAL [®] 90, AG-SURF [®] or CITOWETT [®] PLUS	2 litres/1000 litres of spray solution

MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is dissolved before adding Banvel II or Banvel Dry EG Herbicide. Refer to MIXING DIRECTIONS section for general directions.

WEEDS CONTROLLED

Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide plus Group 2 resistance KOCHIA.

PP-2525 Toss-N-Go® Herbicide + 2,4-D ESTER + BANVEL* II or BANVEL* Dry EG Herbicide

LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.

CROP (S): Spring wheat (including durum) and spring barley

TIMING: Apply the tank mix from the 3-5 leaf stage of wheat (including durum) and barley, and to kochia plants up to the 8-leaf stage. Expect superior results when kochia is small. Use the higher rate of Banvel* II or Banvel* Dry EG Herbicide when the majority of Kochia plants are at or near the 8-leaf stage, or when growing conditions are not favourable.

TANK MIX COMPONENTS AND APPLICATION RATES

PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
Banvel* II Herbicide or	110 – 145 mL/ha or
Banvel* Dry EG Herbicide	76 - 100 g/ha

2,4-D Ester LV700	0.6 L/ha (or 420 g/ha acid equivalent)
Adjuvants: AGRAL 90, AG-SURF or CITOWETT PLUS	2 litres/1000 litres of spray solution
MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is dissolved before adding Banvel* II or Banvel* Dry EG Herbicide. Add 2,4-D Ester last. Refer to MIXING DIRECTIONS section for general directions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide plus 2,4-D Ester, plus Group 2 resistance KOCHIA.	

SPRING WHEAT (EXCLUDING DURUM) AND SPRING BARLEY

PP-2525 Toss-N-Go® HERBICIDE + AXIAL* 100EC Herbicide	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): Spring wheat (excluding durum) and spring barley only	
TIMING: Post-emergent application from 1 leaf to flag leaf. For annual grass weeds, apply from 1 to 6-leaf stage, up to emergence of 4 th tiller.	
TANK MIX COMPONENTS AND APPLICATION RATES	
PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
AXIAL* 100 EC Herbicide	600 mL/hectare
ADIGOR* Adjuvant	700 mL/hectare
MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is completely dissolved. Then add Axial* 100 EC Herbicide with reduced agitation, and finally either Adigor* adjuvant. DO NOT USE ADDITIONAL NON-IONIC SURFACTANT. Refer to MIXING DIRECTIONS section for general instructions. Consult the Axial* 100EC Herbicide label for application instructions and use precautions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide alone, plus wild oats, volunteer oats, green foxtail, yellow foxtail, volunteer canary seed and proso millet.	
NOTES: Do not harvest within 60 days for grain and straw and 30 days for hay. Do not graze treated crop for 7 days after application. Consult AXIAL* 100 EC Herbicide label for additional information on appropriate weed staging, application instructions, environmental and use precautions.	

FOR WEED CONTROL IN SPRING WHEAT AND DURUM WHEAT

PP-2525 Toss-N-Go® Herbicide + HARMONY® GRASS 128EC Herbicide	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP: Wheat (spring and durum) only, not underseeded to legumes	
TIMING: Post-emergent application from 2-leaf to flag leaf growth stage of the crop. Apply to actively growing wild oats from the 1 to 6-leaf stage, prior to emergence of the 4 th tiller and to green foxtail from the 1 to 5-leaf stage, prior to emergence of the 3 rd tiller.	
TANK MIX COMPONENTS AND APPLICATION RATES	
PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
HARMONY® GRASS 128EC Herbicide	0.438 L/Ha
MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is completely dissolved. Reduce agitation, and then add HARMONY® GRASS 128EC Herbicide. Refer to MIXING DIRECTIONS section for general instructions. Use minimum volume of 55 L/ha. Consult the HARMONY® GRASS 128EC Herbicide label for additional application instructions and use precautions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide alone, green foxtail and wild oats.	
NOTES: Do not permit lactating dairy animals to graze fields within 7 days after application.	

PP-2525 Toss-N-Go® Herbicide + HORIZON* NG Herbicide	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP: Wheat (spring and durum) only, not underseeded to legumes	
TIMING: Post-emergent application from 2-leaf to flag leaf growth stage of the crop. Apply to actively growing wild oats from the 1 to 6-leaf stage, prior to emergence of the 4 th tiller and to green foxtail from the 1 to 5-leaf stage, prior to emergence of the 3 rd tiller.	
TANK MIX COMPONENTS AND APPLICATION RATES	
PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
HORIZON* NG Herbicide	0.930 L/Ha
No adjuvant required.	
MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is completely dissolved. Reduce agitation, and then add HORIZON* NG Herbicide. Refer to MIXING DIRECTIONS section for general instructions. Use minimum volume of 55 L/ha. Consult the HORIZON* NG Herbicide label for additional application instructions and use precautions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide alone, green foxtail and wild oats.	
NOTES: Do not permit lactating dairy animals to graze fields within 7 days after application.	

PP-2525 Toss-N-Go® Herbicide + HORIZON* 240 EC Herbicide Tank Mix	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP: Wheat (spring and durum) only, not underseeded to legumes	
TIMING: Post-emergent application from 2-leaf to flag leaf growth stage of the crop. Apply to actively growing wild oats from the 1 to 6-leaf stage, prior to emergence of the 4 th tiller and to green foxtail from the 1 to 5-leaf stage, prior to emergence of the 3 rd tiller.	
TANK MIX COMPONENTS AND APPLICATION RATES	
PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
HORIZON* 240 EC Herbicide	0.230 L/Ha
Adjuvant: SCORE*	0.8 L/Ha
MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is completely dissolved. Reduce agitation, and then add Horizon* 240 EC Herbicide. Then add Score* adjuvant. Refer to MIXING DIRECTIONS section for general instructions. Use minimum volume of 55 L/ha. Consult the Horizon 240 EC Herbicide label for additional application instructions and use precautions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide alone, green foxtail and wild oats.	
NOTES: Do not permit lactating dairy animals to graze fields within 7 days after application.	

PP-2525 Toss-N-Go® Herbicide + SIMPLICITY*	
LOCATION: For use in the Prairie Provinces and Peace River Region of British Columbia only.	
CROP(S): Spring wheat and durum wheat.	
TIMING: Post-emergent application to spring wheat and durum wheat from the 3-leaf stage to the first node stage.	
TANK MIX COMPONENTS AND APPLICATION RATES	
PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
SIMPLICITY*	500 mL/hectare
Adjuvants: AGRAL* 90, AG-SURF* or CITOWETT* PLUS	2.5 litres/1000 litres of spray solution
MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is completely dissolved. Reduce agitation, and then add SIMPLICITY*, followed by adjuvant. Refer to MIXING DIRECTIONS section for general instructions. Consult the SIMPLICITY* label for application instructions and use precautions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide alone, plus weeds listed on the SIMPLICITY* label.	

FOR WEED CONTROL IN SPRING WHEAT ONLY

PP-2525 Toss-N-Go® Herbicide+ 2,4-D ESTER + EVEREST* 70 DF OR EVEREST* SOLUPAK 70	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP (S): Spring wheat only	
TIMING: Apply to spring wheat in the 3 to 4 leaf stage, plus 2 tillers; to wild oats and green foxtail in the 1-6 leaf stage (1-4 leaves on main stem, plus 2 tillers).	
TANK MIX COMPONENTS AND APPLICATION RATES	
PP-2525 Toss-N-Go® Herbicide	30 grams/hectare
EVEREST* 70 DF or EVEREST* SOLUPAK 70	43 Grams/Ha
2,4-D Ester LV700	0.6 L/ha (or 420 g/ha acid equivalent)
Non-Ionic Surfactant	2.5 L/1000 L (0.25 % v/v)
MIXING: Ensure that PP-2525 Toss-N-Go® Herbicide is completely dissolved. Add Everest Solupak 70 DF or Everest 70 DF while continuing agitation until these products are completely mixed. Reduce agitation, and add 2,4-D, and then adjuvant. Refer to MIXING DIRECTIONS section for general instructions. Consult the 2,4-D Amine or Ester and Everest Solupak 70 DF or Everest 70 DF labels for additional information on rotational cropping restrictions, application instructions, use precautions and other restrictions.	
WEEDS CONTROLLED Broadleaf weeds controlled or suppressed by PP-2525 Toss-N-Go® Herbicide + 2,4-D, plus wild oats and green foxtail.	
Note: Observe minimum interval to harvest of 80 days after treatment. For black, dark brown and grey-wooded soil zones only, field peas may be grown the year following application in fields where precipitation has been normal or above normal (10 year average) during the growing season, and where organic matter content is above 4% and pH below 7.5. Re-crop spring wheat only in brown soil zone. Consult the EVEREST* 70 DF or EVEREST* SOLUPAK 70 labels for additional information on rotational crop restrictions, application instructions, use precautions and other restrictions.	

MIXING INFORMATION:

MIXING PRECAUTIONS:

If the sprayer has been left standing for a period of time agitate thoroughly to re-suspend the tank mix partner before resuming spraying. Spray preparation should be used within 24 hours or product degradation may occur.

NOTE: Do not mix the soluble bags with liquid fertilizers, substances that contain boron or substances that release free chlorine. Mixing the soluble bags with any of these compounds will result in an insoluble substance in the tank.

MIXING DIRECTIONS:

Select a spray volume that will ensure thorough coverage and a uniform spray pattern. Use a minimum of 55 L of spray volume per hectare for ground application, and between 25 to 50 L of spray volume per hectare for aerial application. If tank mixing with other herbicides, always consult the label of the tank mix partner(s) for minimum spray volume requirements and apply the tank mixture using a water volume recommended for all products.

1. Always start with a clean and empty sprayer tank.
2. Fill the tank to 2/3 full of **clean** water.
3. With the agitator running, add the required amount of PP-2525 Toss-N-Go® Herbicide. Continue to agitate for a minimum of 5 minutes to ensure that PP-2525 Toss-N-Go® Herbicide is **completely** dissolved.
4. Make sure there are no un-dissolved granules, and then add the required amount of the tank mix partner. If the tank mix partner is an Emulsifiable Concentrate (EC), reduce agitation to avoid inducing an invert emulsion.
5. Add the rest of the water.
6. If required for the tank mixture, add the appropriate adjuvant. If an antifoam agent is required, add last.
7. Refer to the tank mix sections of this booklet for mixing order and other mixing instructions.
8. For repeat tank loads, reduce the material remaining in the tank to 10% of the original volume **or less** before proceeding with step 1, because remaining chemicals may prevent PP-2525 Toss-N-Go® Herbicide granules from completely dissolving or dispersing. If this is not possible, pre-slurry PP-2525 Toss-N-Go® Herbicide in a small amount (5-10 L) of water before adding to the tank.

SPECIAL MIXING INSTRUCTIONS FOR CHEMICAL HANDLERS

1. Ensure the chemical handler is clean (rinsed with ammonia).
2. For best results, add PP-2525 Toss-N-Go® Herbicide to the top of the sprayer tank.

3. If a chemical handler is used, make sure that all of the water soluble bags and granules of PP-2525 Toss-N-Go® Herbicide are completely dissolved/dispersed and injected into the main tank with agitation before adding other chemicals.
4. Low water volumes in some chemical handlers can cause the water to become saturated with PP-2525 Toss-N-Go® Herbicide, leaving un-dissolved granules. In that case, rinse the chemical handler with clean water before adding any other chemical.
5. Always dissolve the PP-2525 Toss-N-Go® Herbicide granules with agitation in the main tank before adding a tank mix partner.

NOTE: Follow the clean-out procedure recommended for sprayer tanks for the chemical handler.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact FMC Corporation at 1-833-362-7722 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by FMC Corporation.

EQUIPMENT CLEANING INFORMATION:

CLEANING PRECAUTIONS:

CAUTION: Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour, which may cause eye, nose, throat, and lung irritation. Do not clean equipment in an enclosed area.

DO NOT CLEAN SPRAYER NEAR WELL OR WATER SOURCE OR NEAR TO DESIRABLE VEGETATION.

CLEANING INSTRUCTIONS:

To avoid subsequent injury to crops other than wheat, spring barley or oats, immediately after spraying and prior to spraying other crops, thoroughly remove all traces of PP-2525 Toss-N-Go® Herbicide from mixing and spray equipment as follows:

1. Drain tank and flush tank, boom and hoses with clean water. If boom is equipped with multiple nozzle bodies, be sure to rotate through all nozzles to ensure clean water reaches all parts of these assemblies.
2. Visually inspect tank to ensure removal of all visible herbicide residues. If necessary, repeat Step 1.
3. Fill the tank with clean water, and then add 1 litre household AMMONIA (containing minimum of 3% ammonia) or equivalent amount of a sprayer tank cleaner containing ammonia, per 100 litres of water. Ammonia will not neutralize the herbicide, but helps dissolve dried-on herbicide deposits. If an emulsifiable concentrate has been tank mixed with PP-2525 Toss-N-Go® Herbicide, use of a wetting agent in the cleaning process will assist in removing oily residue that may trap PP-2525 Toss-N-Go® Herbicide on the tank and hoses.
4. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation.

5. Drain the tank.
6. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
7. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the hoses and boom.

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, PP-2525 Toss-N-Go® Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to PP-2525 Toss-N-Go® Herbicide and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of PP-2525 Toss-N-Go® Herbicide or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your local FMC Corporation representative or the FMC Corporation hotline at 1-833-362-7722 for further information.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

DISPOSAL:

If this is a cardboard container:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

If this is a refillable container:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

If this is a returnable container:

Do not reuse this container for any other purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For all package types:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

©/TM

* All other products mentioned are trademarks of their respective companies.