

2024-08-20
2023-4301

(Container)



with RINSKOR™ ACTIVE

GF-3206 Herbicide

| | | |
|-------|---|-----------|
| GROUP | 4 | HERBICIDE |
|-------|---|-----------|

For control of susceptible terrestrial weeds and certain woody plants, including invasive and noxious weeds, on hazelnut, rangeland, permanent grass pastures, non-cropland areas (such as roadsides, railroad and utility rights-of way), non-irrigation ditch banks, natural areas and grazed areas in and around these use sites.

COMMERCIAL

EMULSIFIABLE CONCENTRATE

READ THIS LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: Florpyrauxifen 19.91 g/L
(present as benzyl ester)

Contains the allergen soy

REGISTRATION NO. 34731 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1 L - bulk

Corteva Agriscience Canada Company

Suite 240, 115 Quarry Park Rd. SE
Calgary, AB
T2C 5G9
1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

PERSONAL PROTECTIVE EQUIPMENT

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab and/or cockpit.

OPERATOR USE PRECAUTIONS

- Wash hands before eating, drinking, using tobacco or using the washroom.
- If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing.
- Remove personal protective equipment immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.

AERIAL APPLICATION PRECAUTIONS

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. The pilot is allowed to load premixed chemicals with a closed system.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

RESTRICTED ENTRY INTERVALS

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours following application on agricultural areas.

For industrial and other non-crop areas, DO NOT enter or allow worker entry into treated areas until sprays have dried.

SPRAY DRIFT PRECAUTIONS

Apply only when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

STORAGE

Store this product away from food or feed.

DISPOSAL

Recyclable containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

Returnable containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

(Booklet)



with RINSKOR™ ACTIVE

GF-3206 Herbicide

| | | |
|-------|---|-----------|
| GROUP | 4 | HERBICIDE |
|-------|---|-----------|

For control of susceptible terrestrial weeds and certain woody plants, including invasive and noxious weeds, on hazelnut, rangeland, permanent grass pastures, non-cropland areas (such as roadsides, railroad and utility rights-of way), non-irrigation ditch banks, natural areas and grazed areas in and around these use sites.

COMMERCIAL

EMULSIFIABLE CONCENTRATE

READ THE LABEL AND THIS BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: Florpyrauxifen 19.91 g/L
(present as benzyl ester)

Contains the allergen soy

REGISTRATION NO. 34731 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 0.1 L - bulk

Corteva Agriscience Canada Company

Suite 240, 115 Quarry Park Rd. SE

Calgary, AB

T2C 5G9

1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

PERSONAL PROTECTIVE EQUIPMENT

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab and/or cockpit.

OPERATOR USE PRECAUTIONS

- Wash hands before eating, drinking, using tobacco or using the washroom.
- If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing.
- Remove personal protective equipment immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.

AERIAL APPLICATION PRECAUTIONS

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. The pilot is allowed to load premixed chemicals with a closed system.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

RESTRICTED ENTRY INTERVALS

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours following application on agricultural areas.

For industrial and other non-crop areas, DO NOT enter or allow worker entry into treated areas until sprays have dried.

SPRAY DRIFT PRECAUTIONS

Apply only when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

STORAGE

Store this product away from food or feed.

DISPOSAL

Recyclable containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

Returnable containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

GENERAL INFORMATION

GF-3206 is used for control of susceptible terrestrial weeds including invasive and noxious weeds and certain woody plants in hazelnut, on rangeland, permanent grass pastures, Conservation Reserve Program (CRP) acres, noncropland areas (such as roadsides, railroad and utility rights-of way), non-irrigation ditch banks, natural areas and grazed areas in and around these use sites.

GF-3206 is a broadleaf weed herbicide used for the control of broadleaf weeds in pastures and hayfields and is selective to most grasses and white clover. GF-3206 may be applied post-emergence as a broadcast spray or as a spot application to control weeds listed on this label. Best weed control results are obtained when spray volume is sufficient to provide uniform coverage of treated plants. For optimum uptake and translocation of the herbicide, avoid mowing, haying, shredding, burning or soil disturbance in treated areas for at least 7 days following application.

Applications should be avoided under conditions of drought or other environmental stresses. **Read all precaution statements before using this product.** For more information, contact your local Corteva Agriscience representative.

For best results, apply GF-3206 Herbicide to young weeds that are actively growing at time of application.

GENERAL USE PRECAUTIONS

GF-3206 Herbicide is highly active against broadleaf plants. This product is recommended for use on areas where loss of broadleaf forage plants, including legumes, can be tolerated.

Poor weed control may result from application of GF-3206 made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, prior herbicide applications, or soils with high salt content.

Do not apply where runoff or irrigation water may flow directly onto agricultural land. Do not use treated water for any form of irrigation.

Only one application per growing season is permitted.

GF-3206 Herbicide cannot be applied on domestic or commercial turf grass.

Tank Mixtures

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Corteva Agriscience Canada Company at 1-800-667-3852 or www.corteva.ca for information before applying any tank mix that is not specifically recommended on this label.

DIRECTIONS FOR USE

READ ALL DIRECTIONS CAREFULLY BEFORE APPLYING. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

RANGELAND, PERMANENT PASTURE, INDUSTRIAL AND OTHER NON-CROP AREAS OF CANADA

APPLICATION METHODS

GF-3206 Herbicide can be applied post-emergence as a broadcast (ground or aerial) spray or as a spot/strip application to control weeds and woody plants. Post emergence applications should be made before bud stage or early flowering, unless otherwise specified. Best results are obtained when the spray volume is sufficient to provide uniform coverage of treated plants. For optimum uptake and translocation of the herbicide, avoid mowing, haying, shredding or disturbing treated areas for at least 7 days following application.

Applications of GF-3206 Herbicide may be made once grasses are well-established (have developed a good secondary root system and show good vigour).

Individual Plant or Spot/Strip Applications

Use the following mixing chart to mix GF-3206 Herbicide with enough water to make the correct amount of spray solution. Use equipment designed for individual plant or spot/strip applications, such as a backpack sprayer or hose and handgun, to spray susceptible invasive plants. Thoroughly and uniformly wet the foliage of all target plants but not to the point of runoff. Direct spray away from aquatic habitats and non-target terrestrial plants. DO NOT apply this product directly to fresh water habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands), estuaries or marine habitats.

Mixing Chart

| GF-3206 Herbicide | Surfactant (at 0.5%v/v)* | Spray Solution (water and herbicide) |
|-------------------|--------------------------|--------------------------------------|
| 2.0-20 mL | 50 mL | 10 litres |
| 20- 200 mL | 500 mL | 100 litres |
| 200- 2000 mL | 5000 mL | 1000 litres |

*Use a surfactant such as Gateway Adjuvant, MSO or NIS (such as Ag-Surf, Agral 90 or Citowett Plus)

Use the higher rate in the rate range when growing conditions are less than favourable or when the plant foliage is tall and dense.

Preharvest/Grazing Intervals

The following restrictions for livestock or lactating dairy animals grazing in treated areas must be followed:

- Forage: Grazing/feeding is permitted on the day of application (0 days)
- Hay: Allow 7 days after application to cut or feed hay.
- Allow 3 days of grazing on an untreated pasture (or feed untreated hay) before transferring livestock to areas where sensitive broadleaf crops may be grown.

GF-3206 HERBICIDE ALONE

Apply GF-3206 Herbicide at a rate of 200 – 2400 mL/ha. Apply when targeted undesirable vegetation is actively growing. Only weeds and brush present at the time of application will be controlled. Apply with a minimum of 100 L spray volume/ha, that will assure uniform coverage. For better coverage use 200 L spray volume/ha for ground application. Use a surfactant, such as Gateway Adjuvant or an NIS at 0.25-0.5% v/v or methylated seed oil (MSO) at 0.5-1% v/v .Use the greater amount for harder to control weeds. See surfactant label for a full list of recommended rates.

Weeds Controlled or Suppressed by GF-3206 Herbicide Alone

| Rate of GF-3206 Herbicide | Weeds |
|---------------------------|--|
| 200 mL/ha | Canada fleabane ¹ velvetleaf ² common waterhemp ³ |
| 400 mL/ha | cleavers ⁴ common lamb's quarter ⁵ annual sowthistle ⁶ wild chervil ⁷ . wild parsnip wild caraway tall buttercup |

| | |
|-------------------|---|
| 800 mL/ha | dandelion ⁸ shepherd's purse ⁹ kochia ¹⁰ wild buckwheat ¹¹ |
| 1200 mL/ha | common barnyard grass ¹² |
| 1600 mL/ha | redroot pigweed ¹³ |
| 2400 mL/ha | tansy* |

¹Canada fleabane: Bolting up to 20 cm

²Velvetleaf: Up to 3 leaf and 8 cm

³Common waterhemp: Up to 3 leaf and 8 cm

⁴Cleavers: Up to 10 cm:

⁵Common lamb's quarter: Prior to budding stage

⁶Annual sowthistle: Up to 3 cm

⁷Wild chervil: Up to budding stage and 40 cm

⁸Dandelion: Rosette stage to beginning flowering, 35 plants/m² or less

⁹Shepherd's purse: Up to flowering stage

¹⁰Kochia: Bolting up to 10 cm <165 plants/m²

¹²Wild buckwheat

¹²Common barnyard grass: Vegetation stage up to 20 cm

¹³Redroot pigweed: Up to 6 leaf

*Suppression

CROP USES - HAZELNUT

GF-3206 Herbicide can be used to control labeled weeds in hazelnut at up to 400 mL/ha.

TOTAL VEGETATION CONTROL TANK-MIXES

Tank-Mix Combinations – GF-3206 Herbicide plus Arsenal Herbicide or glyphosate herbicides (present as isopropylamine salt, diammonium salt, potassium salt or dimethylamine salt, registered for use on non-cropland areas)

GF-3206 Herbicide can be tank-mixed with either Arsenal Herbicide or glyphosate herbicides (registered for use on non-cropland areas) to broaden the spectrum of broadleaf weeds and invasive plants controlled. These tank-mixes will provide control of most grass and broadleaf species. Use these tank-mixes on non-crop industrial sites such as railroad, electrical, highway, telephone and pipeline rights-of-way.

GF-3206 Herbicide plus Arsenal Herbicide: Apply 3.0 L/ha of Arsenal Herbicide tank mixed with 0.2 – 2.4 L of GF-3206 Herbicide per hectare.

GF-3206 Herbicide plus glyphosate herbicides (present as isopropylamine salt, diammonium salt, potassium salt or dimethylamine salt, registered for use on non-cropland areas): Apply 810 - 4320 g a.e. per hectare of glyphosate herbicide (present as isopropylamine salt, diammonium salt, potassium salt or dimethylamine salt, registered for use on non-cropland areas) tank mixed with 0.2 – 2.4 L of GF-3206 Herbicide per hectare.

MIXING METHODS

Mixing with Water

1. Fill the spray tank 3/4 full of clean water
2. Add the required amount of GF-3206 Herbicide with the agitation running. Pre-slurrying with water may be necessary where there is little or no agitation or an injection system is being used or where herbicide is first added to a tank other than the spray tank.

3. If tank mixing, add the required amount of the tank-mix partner with the agitation running.
4. Add the surfactant.
5. Add antifoaming agent, if required.

SPRAYER CLEAN-OUT INSTRUCTIONS

To avoid injury to desirable plants, thoroughly clean equipment used to apply this product before re-use or using it to apply other chemicals.

1. Immediately after spraying, completely drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. First rinse:
 - Spray the inside of tank with clean water and fill the sprayer with at least one tenth of the spray tank volume.
 - Agitate and circulate for 15 minutes, and flush through booms and hoses.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.
3. Second rinse:
 - Fill the tank with clean water.
 - Add All Clear Spray Tank Decontaminator, or Clean-Out Spray Tank Cleaner, or 1 L of household ammonia (containing a minimum of 3 % ammonia) per 100 L of water, or similar tank cleaning agent as per manufacturer's recommendations while filling the tank with clean water.
 - Agitate and then flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full. Allow to stand for 15 minutes with agitation. Flush the solution out of the spray tank through the spray booms. Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - If possible, let the solution stand in the sprayer tank and booms for an extended period of time, overnight if possible.
 - After flushing the boom and hoses, drain tank completely.
 - Remove nozzles and screens and clean separately with a cleaning agent or an ammonia solution (100 mL in 10 L water).
4. Third rinse:
 - Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.

Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty odour which may cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

INVASIVE PLANT MANAGEMENT

Invasive Plant Management Practices

GF-3206 Herbicide can be an important component of integrated vegetation management programs designed to restore desired plant communities. To maximize and extend the benefits of weed control provided by GF-3206 Herbicide, it is important that other vegetation management practices, including proper grazing management, fertilization, prescribed fire, mechanical mowing, hand-pulling, bio control and other methods, be used in appropriate sequences and combinations to further alleviate the adverse effects of invasive plants on desirable plant species and to promote development of desired plant communities. Consult local agriculture and land management resource specialists and/or invasive plant councils. These organizations can provide guidance on best management practices and the development of integrated vegetation management programs. Rapid response to the early detection of

new invasive plants is a key invasive plant management strategy. Containment, eradication and control are essential for the management of spreading and established invasive plants.

Rangeland and Pasture Areas

Make only one application per year for control of invasive plants. For best results, invasive plants should be treated when they are actively growing and under conditions favourable for growth. GF-3206 Herbicide provides control of emerged seedlings of susceptible plants following application.

To reduce the spread of invasive plants, it is important to treat identified invasive plants in the target area while observing required spray buffer zones for broadcast applications. Spray when the wind is blowing from the opposite direction to complete spraying the targeted plant population.

Individual Plant or Spot/Strip Applications

An important component of Invasive Plant Management programs is early detection and eradication. The instructions for individual plant or spot or strip applications can be used to stop the spread of invasive plants. This is particularly important in riparian areas, where spot application is preferable to broadcast treatment.

Field Sprayer Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Aerial Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) very coarse classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

Use GF-3206 Herbicide alone, or tank mixed as a broadcast treatment by air to control listed broadleaf weeds and woody plants. Apply GF-2306 Herbicide at the recommended rates in a minimum spray volume of 30 L/ha by air. For better coverage use 50 litres/ha. Follow all instructions under Mixing Methods for surfactant requirements and order of mixing.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate swath marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-667-3852 or obtain technical advice from the distributor or your provincial agricultural representative.

SPRAY DRIFT PRECAUTIONS AND SPRAY BUFFER ZONES

Do not apply to any body of water. Take care to minimize the incidental overspray along the shoreline when applying to terrestrial plants at the water's edge or to water in areas where surface water is present.

Avoid spray drift at the application site. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. Users are responsible for considering all these factors when making decisions.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt).

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label, or
- low-clearance hooded or shielded sprayers that prevent spray contact with crop or foliage.

For application to rights-of-way, spray buffer zones for protection of sensitive terrestrial habitats are not required; however, the best available application strategies which minimize off-site drift, including meteorological conditions (e.g., wind direction, low wind speed) and spray equipment (e.g., coarse droplet sizes, minimizing height above canopy), should be used. Applicators must, however, observe the specified spray buffer zones for protection of sensitive aquatic habitats.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

| Method of application | Crop | Spray Buffer Zones (metres) Required for the Protection of: | | | |
|-----------------------|--|---|------------------|----------------------|-----|
| | | Freshwater Habitat of Depths: | | Terrestrial Habitat: | |
| | | Less than 1 m | Greater than 1 m | | |
| Field sprayer | rangeland, permanent grass pastures, non-irrigation ditch banks, natural areas and grazed areas in and around these use sites* | 80 | 35 | 15 | |
| | hazelnut | 15 | 10 | 3 | |
| Aerial | rangeland, permanent grass pastures, non-irrigation ditch banks, natural areas and grazed areas in and around these use sites* | Fixed wing | 800 | 800 | 575 |
| | | Rotary wing | 800 | 625 | 275 |

* Spray buffer zones for the protection of terrestrial habitats are not required for use on rights-of-way including railroad ballast, rail and hydro rights-of-way, utility easements, roads, and training grounds and firing ranges on military bases.

When tank-mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank-mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank-mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website. Spray buffer zones for aerial applications of this product CANNOT be modified.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, GF-3206 Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to GF-3206 Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Development of plant populations resistant to this herbicide mode of action is usually not a problem on rangeland, permanent grass pastures, Conservation Reserve Program (CRP), or non-cropland sites since these sites receive infrequent pesticide applications.
- Similar looking biotypes of a given weed species occurring in a treated area may vary in their susceptibility to a herbicide. Application of a herbicide below its labeled rate may allow more tolerant weeds to survive and a shift to more tolerant biotypes within the treated area.
- Where possible, rotate the use of GF-3206 Herbicide or other Group 4 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Corteva Agriscience Canada Company at 1-800-667-3852 or at www.corteva.ca.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

™Trademark of Corteva Agriscience and its affiliated companies.