

Booklet

GROUP	2	HERBICIDE
-------	---	-----------

FBN IMAZAMOX WDG

Water Dispersible Granular Herbicide in
water-soluble bags

COMMERCIAL (AGRICULTURAL)

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND
PEACE RIVER AREA OF BRITISH COLUMBIA ONLY

ACTIVE INGREDIENT: Imazamox..... 70%

Warning, contains the allergen sulfites

REGISTRATION NO. 34786 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1 kg - Bulk

WARNING - EYE IRRITANT

FOR EMERGENCY MEDICAL ASSISTANCE (HUMAN OR ANIMAL) CONTACT ROCKY
MOUNTAIN POISON CONTROL AT 1-866-767-5041.

FOR CHEMICAL EMERGENCY ASSISTANCE (SPILL, LEAK, FIRE OR ACCIDENT)
CONTACT CHEMTREC AT 1-800-424-9300 (NORTH AMERICA) OR 1-703-527-3887
(INTERNATIONAL).

READ THE LABEL AND THIS BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

DO NOT SELL INDIVIDUAL WATER-SOLUBLE BAGS SEPARATELY

Farmer's Business Network Canada, Inc.
PO Box 5607
High River, Alberta
Canada T1V 1M7
1-844-200-FARM (3276)

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DO NOT APPLY BY AIR.

Causes eye irritation. DO NOT get in eyes.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

Store above 5°C.

Store in a cool, dry place away from children, animals, food, feed or fertilizers.

Keep unused water-soluble bags in resealed original container.

Keep packages dry at all times.

Keep packages out of rain.

Keep from freezing.

DISPOSAL

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer and the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a

vegetative filter strip between the treated area and the edge of the water body.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

Field Sprayer Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply by air.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

Spray Buffer Zones

A spray buffer zone is NOT required for uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands).

Method of Application	Crop	Spray Buffer zones (metres) Required for the Protection of Terrestrial Habitat
Field sprayer	All crops	1

When tank mixes are permitted, consult the labels of the tank mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

DO NOT apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, coulees, prairie potholes, creeks, marshes, streams, reservoirs, ditches and wetlands), estuaries or marine habitats.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

TANK MIXING

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer

Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Farmer's Business Network Canada, Inc. at 1-844-200-FARM (3276) for information before applying any tank mix that is not specifically recommended on this label.

BROADLEAF WEED AND GRASS CONTROL

FBN Imazamox WDG at 21 g/ha (15 g a.e./ha) will control only:	
Lamb's-quarters*	Volunteer spring wheat
Stinkweed	(non- Clearfield tolerant wheat)
Wild mustard	
FBN Imazamox WDG at 29 g/ha (20 g a.e./ha) will control:	
Barnyard grass	Volunteer barley
Cleavers*	Volunteer canary seed
Cow cockle	Volunteer canola (non- Clearfield
Green foxtail	tolerant canola only)
Green smartweed	Volunteer durum wheat
Japanese brome grass*	Volunteer spring wheat (non- Clearfield
Kochia*	tolerant wheat)
Lamb's-quarters	Volunteer tame oats
Persian darnel	Wild buckwheat*
Redroot pigweed	Wild mustard
Round-leaved mallow*	Wild oats
Russian thistle	Yellow foxtail
Shepherd's purse	
Stinkweed	

* = suppression only

Apply in 100 L/ha of water with MERGE® Adjuvant at a rate of 0.5% v/v (e.g. 5 L of MERGE Adjuvant per 1000 L spray solution).

FOR FIELD PEAS

DIRECTIONS FOR USE (Prairie Provinces and Peace River Area of British Columbia only)

FBN Imazamox WDG is a selective herbicide that can be applied as an early postemergence treatment in field peas.

When FBN Imazamox WDG is applied early post-emergence, absorption may occur through both the roots and foliage. Susceptible weeds stop growing and eventually die.

Apply using ground equipment only. DO NOT APPLY BY AIR.

DO NOT apply more than once per year.

DO NOT graze the treated field peas or cut for hay within 20 days of application.

Pre-harvest Interval: After 60 days, field peas can be harvested.

TIMING OF APPLICATION

Application should be made from the 1 to 6 true leaf stage and after weeds have emerged. Apply when broadleaf weeds are from the cotyledon to 4-leaf stage and when grassy weeds are at the 1 to 4 true leaf or early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.

Using Water-Soluble Packages Dissolved Directly in Spray tanks:

Water-Soluble Packages (WSPs) are designed to dissolve in water. Agitation may be used, if necessary, to help dissolve the WSP. Failure to follow handling and mixing instructions can increase your exposure to the pesticide products in WSPs.

Handling Instructions

Follow these steps when handling pesticide products in WSPs.

1. Mix in spray tank only.
2. Handle WSP(s) in a manner that protects package from breakage and/or unintended release of contents. If package is broken, put on a minimum of coveralls, chemical-resistant gloves, chemical-resistant footwear, and a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested and then continue with mixing instructions.
3. Keep the WSP(s) in outer packaging until just before use.
4. Keep the WSP dry prior to adding to the spray tank.
5. Handle with dry gloves and according to the label instructions for PPE.
6. Keep WSP intact. Do not cut or puncture WSP.
7. Reseal the WSP outer packaging to protect any unused WSP(s).

Mixing Instructions

Follow the steps below when mixing this product, including if tank mixed with other pesticide products. If being tank mixed, the mixing directions 1 through 9 below take precedence over the mixing directions of the other tank mix products. All other directions for use of all tank mixed products should be followed provided they do not conflict. Do not tank mix this product with products that prohibit tank mixing or have conflicting mixing directions.

1. If a basket or strainer is present in the tank hatch, remove prior to adding the WSP to the tank.
2. Fill tank with water to approximately one-third to one-half of the desired final volume of spray.
3. Stop adding water and stop any agitation.
4. Place intact/unopened WSP(s) into the tank.
5. Do not spray water from a hose or fill pipe to break or dissolve the WSP(s).
6. Start mechanical and recirculation agitation from the bottom of tank without using any overhead recirculation, if possible. If overhead recirculation cannot be turned off, close the hatch before starting agitation.

7. Dissolving the WSP(s) may take up to 5 minutes or longer, depending on water temperature, water hardness and intensity of agitation.
8. Stop agitation before tank lid is opened.
9. Open the lid to the tank, exercising caution to avoid contact with dusts or spray mix, to verify that the WSPs have fully dissolved and the contents have been thoroughly mixed into the solution.
10. Do not add other allowed products or complete filling the tank until the bags have fully dissolved and pesticide is thoroughly mixed.
11. Once the WSP have fully dissolved and any other products have been added to the tank, resume filling the tank with water to the desired level, close the tank lid, and resume agitation.
12. Use the spray solution when mixing is complete.
13. Maintain agitation of the diluted pesticide mix during transport and application.
14. It is unlawful to use any registered pesticide, including WSPs, in a manner inconsistent with its label.

Tank Mix Pre-harvest / Grazing Interval

Do not harvest the treated crop within 60 days of application.

MIXING INSTRUCTIONS

1. Use 100 L/ha of water.
2. Use a 50-mesh (or coarser) filter screen.
3. Fill the spray tank with three-quarters of the required amount of clean water, start agitation and continue agitation throughout the entire mixing and spraying procedure.
4. Add the required amount of FBN Imazamox WDG soluble bag(s) directly into the sprayer through the tank opening.
5. Agitate for at least ten minutes to dissolve the herbicide. After the herbicide is dissolved, continue agitation.
6. If required, add the required amount of MERGE Adjuvant. If excess foaming occurs, a silicone anti-foaming agent may be added (e.g. HALT®).
7. Complete filling the tank to the desired level with water. If agitation is stopped for more than 5 minutes, re-suspend spray solution by full agitation prior to commencing spraying again.
8. Between loads of FBN Imazamox WDG, check in-line and nozzle screens and rinse clean if necessary.
9. Upon completion of spraying FBN Imazamox WDG, thoroughly flush tank, boom, hoses and in-line and nozzle screens with clean water to avoid possible injury to other crops.

FOLLOW CROPPING

Winter wheat can be planted 3 months after treatment (3 MAT) as a rotational crop.

Initial crop injury to non-**Clearfield** canola may be observed. Avoid spray overlap as yield reduction may result. The following crops may be grown safely the year following an application:

Canary seed
Field corn
Field peas
Clearfield Canola (canola varieties with the **Clearfield** trait)
Non-**Clearfield** canola
Lentils
Spring wheat
Durum wheat
Spring barley
Sunflower
Tame oats
Flax
Chickpeas
Clearfield Sunflowers (sunflower varieties with the **Clearfield** trait)

The following crop may be grown safely two years following an application:

Mustard (condiment type only)

There are insufficient data for other follow crops. Conduct a field bioassay (a test strip grown to maturity) the year before growing any crop other than those listed above.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, FBN Imazamox WDG is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to FBN Imazamox WDG and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of FBN Imazamox WDG or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance

development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact FBN at 1-844-200-FARM (3276) or at www.fbn.com.

Clearfield is a registered trade-mark of BASF Agrochemical Products B.V., used with permission by BASF Canada Inc.

MERGE is a registered trade-mark of BASF Canada Inc.

®All other products listed are registered trademarks of their respective companies.

FBN IMAZAMOX WDG

Water Dispersible Granular
Herbicide in water-soluble bags

COMMERCIAL
(AGRICULTURAL)

FOR SALE FOR USE IN THE PRAIRIE PROVINCES
AND PEACE RIVER AREA OF BRITISH COLUMBIA
ONLY

ACTIVE INGREDIENT: Imazamox 70%

Warning, contains the allergen sulfites

REGISTRATION NO. 34786 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1 kg - Bulk

WARNING - EYE IRRITANT

FOR EMERGENCY MEDICAL ASSISTANCE (HUMAN OR ANIMAL) CONTACT
ROCKY MOUNTAIN POISON CONTROL AT 1-866-767-5041.
FOR CHEMICAL EMERGENCY ASSISTANCE (SPILL, LEAK, FIRE OR
ACCIDENT) CONTACT CHEMTREC AT 1-800-424-9300 (NORTH AMERICA) OR
1-703-527-3887 (INTERNATIONAL).

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

DO NOT SELL INDIVIDUAL WATER-SOLUBLE BAGS
SEPARATELY

Farmer's Business Network Canada, Inc.
PO Box 5607
High River, Alberta
Canada T1V 1M7
1-844-200-FARM (3276)

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DO NOT APPLY BY AIR.

Causes eye irritation. DO NOT get in eyes.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control

centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

Store above 5°C.

Store in a cool, dry place away from children, animals, food, feed or fertilizers.

Keep unused water-soluble bags in resealed original container.

Keep packages dry at all times.

Keep packages out of rain.

Keep from freezing.

DISPOSAL

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer and the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

Pouch Label

FBN IMAZAMOX WDG
70% Water Dispersible Granular Herbicide
COMMERCIAL (AGRICULTURAL)
NET CONTENTS: 1 kg - Bulk
ACTIVE INGREDIENT:
Imazamox ... 70% active
Warning, contains the allergen sulfites
WARNING - EYE IRRITANT
KEEP OUT OF REACH OF CHILDREN