

2022-09-21
2022-4030
[Sleeve]

GROUP

4

6

HERBICIDES

**IPCO EMIT
HERBICIDE**

Emulsifiable Concentrate

For use as a selective postemergent control of annual and perennial broadleaf weeds in wheat, barley, oats, bromegrass, annual canary grass and canary seed.

COMMERCIAL (AGRICULTURAL)

READ THE LABEL & ATTACHED BOOKLET BEFORE USING



WARNING

POISON

**POTENTIAL SKIN SENSITIZER
EYE & SKIN IRRITANT**

KEEP OUT OF REACH OF CHILDREN

REGISTRATION NO.: 34600 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENTS:

Fluroxypyr, present as 1-methylheptyl ester... 80 g a.e. /L
Bromoxynil, present as octanoate ester ... 200 g/L
MCPA, present as 2-ethylhexyl ester ... 200 g a.e. /L

Interprovincial Cooperative Limited,
P.O. Box 1050, Saskatoon, Saskatchewan S7K 3M9
1-204-233-3461

Net Contents: (1 – Bulk) Litres

IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING, CALL 1-888-CAN-UTEC
(226-8832)

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN.

HARMFUL OR FATAL IF SWALLOWED.

May irritate eyes. Avoid contact with eyes. Causes skin irritation. DO NOT get on skin. Wash concentrate from skin or eyes immediately.

DO NOT APPLY USING AERIAL APPLICATION EQUIPMENT

Avoid breathing spray mist. After use wash hands and other exposed skin.

Avoid contamination of ponds, streams, rivers and other water sources.

Avoid spray drift onto crops other than those recommended.

Store the container tightly closed away from seeds, feeds, fertilizer, plant and foodstuffs.

DO NOT use in residential areas. Residential areas are defined as any use site where bystanders or the public, including children, could be exposed during or after application. This includes areas in and around homes, schools, public buildings, or any other areas where they could be exposed.

Apply only when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PROTECTIVE CLOTHING AND EQUIPMENT:

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Rinse gloves before removal. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during application. Gloves and protective eyewear are not required during application within a closed cab.

After using this product, remove clothing and launder separately and promptly and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry. Wear freshly laundered clothing and clean protective equipment daily.

Only handlers wearing personal protective equipment may be in the area being treated during application.

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them.

Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL PRECAUTIONS:

TOXIC to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE. This product contains active ingredients and aromatic petroleum distillates, which are toxic to aquatic organisms.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product will harm other broadleaved plants in the vicinity of the treatment area. Do not directly spray or allow the spray to drift onto ornamentals or gardens.

LEACHING

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

FIRST AID:

If swallowed, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Contains petroleum distillates - vomiting may cause aspiration pneumonia. High concentrations of MCPA may cause severe irritation to the eyes. Symptoms of overexposure to MCPA could include slurred speech, twitching, jerking and spasms, drooling, low-blood pressure and unconsciousness. Treat symptomatically.

STORAGE:

Store this product away from food or feed. Store in original containers in a secure, dry, heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL:

For Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container and is to be disposed or at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is not container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

EMERGENCY TELEPHONE NUMBERS:

IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING, CALL 1-888-CAN-UTEC (226-8832)

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.

[Booklet]

GROUP	4	6	HERBICIDES
-------	---	---	------------

**IPCO EMIT
HERBICIDE**

Emulsifiable Concentrate

For use as a selective postemergent control of annual and perennial broadleaf weeds in wheat, barley, oats, bromegrass, annual canary grass and canary seed.

COMMERCIAL (AGRICULTURAL)

READ THE LABEL & BOOKLET BEFORE USING



WARNING

POISON

**POTENTIAL SKIN SENSITIZER
EYE & SKIN IRRITANT**

KEEP OUT OF REACH OF CHILDREN

REGISTRATION NO.: 34600 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENTS:

Fluroxypyr, present as 1-methylheptyl ester... 80 g a.e. /L

Bromoxynil, present as octanoate ester ... 200 g/L

MCPA, present as 2-ethylhexyl ester ... 200 g a.e. /L

Interprovincial Cooperative Limited,
P.O. Box 1050, Saskatoon, Saskatchewan S7K 3M9
1-204-233-3461

IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING, CALL 1-888-CAN-UTEC
(226-8832)

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN.

HARMFUL OR FATAL IF SWALLOWED.

May irritate eyes. Avoid contact with eyes. Causes skin irritation. DO NOT get on skin. Wash concentrate from skin or eyes immediately.

DO NOT APPLY USING AERIAL APPLICATION EQUIPMENT

Avoid breathing spray mist. After use wash hands and other exposed skin.

Avoid contamination of ponds, streams, rivers and other water sources.

Avoid spray drift onto crops other than those recommended.

Store the container tightly closed away from seeds, feeds, fertilizer, plant and foodstuffs.

DO NOT use in residential areas. Residential areas are defined as any use site where bystanders or the public, including children, could be exposed during or after application. This includes areas in and around homes, schools, public buildings, or any other areas where they could be exposed.

Apply only when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PROTECTIVE CLOTHING AND EQUIPMENT:

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Rinse gloves before removal. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during application. Gloves and protective eyewear are not required during application within a closed cab.

After using this product, remove clothing and launder separately and promptly and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry. Wear freshly laundered clothing and clean protective equipment daily.

Only handlers wearing personal protective equipment may be in the area being treated during application.

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them.

Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL PRECAUTIONS:

TOXIC to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE. This product contains active ingredients and aromatic petroleum distillates, which are toxic to aquatic organisms.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product will harm other broadleaved plants in the vicinity of the treatment area. Do not directly spray or allow the spray to drift onto ornamentals or gardens.

LEACHING

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

FIRST AID:

If swallowed, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Contains petroleum distillates - vomiting may cause aspiration pneumonia. High concentrations of MCPA may cause severe irritation to the eyes. Symptoms of overexposure to MCPA could include slurred speech, twitching, jerking and spasms, drooling, low-blood pressure and unconsciousness. Treat symptomatically.

STORAGE:

Store this product away from food or feed. Store in original containers in a secure, dry, heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL:

For Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container and is to be disposed or at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.

2. Make the empty, rinsed container unsuitable for further use.

If there is not container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

EMERGENCY TELEPHONE NUMBERS:

IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING, CALL 1-888-CAN-UTEC (226-8832)

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.

GENERAL INFORMATION:

IPCO EMIT provides selective post-emergence control of annual and perennial broadleaf weeds in wheat (spring, durum and winter), barley, oats, brome grass, annual canary grass and canary seed.

DIRECTIONS FOR USE:

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation/drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT apply by air.

Rotational Crop Restrictions: Field previously treated with IPCO EMIT can be seeded the following year to barley, canola, flax, forage grasses, lentils, mustard, oats, peas, rye and wheat or fields can be summerfallowed.

RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, IPCO EMIT is a Group 4 and 6 herbicide. Any weed population may contain or develop plants naturally resistant to IPCO EMIT and other Group 4 and 6 herbicides.

The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of IPCO EMIT or other Group 4 and 6 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information and to report suspected resistance, contact company representatives at 1-204-233-3461

PRODUCT SPECIFIC PRECAUTIONS:

Read and understand the entire label before opening this product. If you have questions, call the manufacturer or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

GROUND SPRAYER:

Wash the tank and clean all filters, screens and tips. Select nozzle tips to apply the recommended volume per hectare. Flood jet type tips are not recommended. Adjust boom height to ensure uniform coverage of the weeds. Ensure that all tips are in good condition and spraying the same volume.

Field sprayer application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural

Engineers (ASAE S572.1) coarse classification. Boom height must be 60 cm or less above the crop or ground.

SPRAY BUFFER ZONES:

A spray buffer zone is NOT required for uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop	Spray Buffer Zones (metres) Required for the Protection of:				
		Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial habitat
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Wheat, barley, oats and canary seed	1	1	1	1	1

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

MIXING:

Half fill the tank with clean water. Add the required amount of IPCO EMIT and agitate thoroughly. Add any tankmix partners. Fill the tank and agitate again before use.

CROPS

- **WHEAT (spring, durum and winter)**
- **BARLEY**
- **OATS**
- **CANARY SEED**
- **BROMEGRASS FOR SEED (See minor uses section)**
- **ANNUAL CANARYGRASS (See minor uses section)**

APPLICATION RATE: 0.625 – 1.25 L/ha

Do not apply more than one treatment per year.

APPLICATION RESTRICTIONS:

- Do not use treated crops for grazing livestock or green feed until 30 days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.
- Do not harvest the treated crop within 60 days after application.
- DO NOT graze, cut or feed treated brome grass to livestock.

SPRAY VOLUME AND PRESSURE: For ground application, apply in 50 to 100 L of water per hectare at a pressure of 275 kPa. For best results when there is a heavy crop canopy, or when the majority of the weeds are at advanced stage of growth, a volume rate of 100 L/ha is recommended.

TIMING: Wheat (spring and durum), barley and oats may be treated from the 2-leaf stage until the early flag leaf stage. Canary seed may be treated from the 3 to 5 leaf stage. Application before the 2-leaf stage may injure the crop. Winter wheat; apply in the fall to well established crops in the 3-leaf stage or larger, OR, apply in the spring at full tillering to the shot-blade stage. Do not apply when the crop is in the flag-leaf stage.

Apply in good growing conditions. Application must be made before the crop shields the weeds. Only weeds that have emerged at the time application will be controlled. Apply when weeds are in the seedling stage (up to 6 leaf stage) and actively growing.

WEEDS CONTROLLED:**Section 1 - IPCO EMIT ALONE**

IPCO EMIT Rate	Broadleaf Weeds Controlled or Suppressed	
0.625 L/ha	kochia (up to 5 cm tall)	wild buckwheat (suppression)
	lamb's quarters	wild mustard
1.25 L/ha	Weeds listed above plus:	
	American nightshade (up to 4-leaf)	redroot pigweed (suppression)
	ball mustard (up to 4-leaf)	Russian thistle (up to 4-leaf, < 5 cm tall)
	bluebur (up to 4-leaf)	scentless chamomile (up to 4-leaf)

Canada thistle (top growth control only)	shepherd's purse (up to 4-leaf)
chickweed	stinkweed
cleavers (up to 6 whorls)	stork's bill (suppression)
cocklebur (up to 4-leaf)	tartary buckwheat
common buckwheat	triazine resistant pigweed (up to 4-leaf)
common groundsel	velvetleaf (up to 4-leaf)
common ragweed	volunteer canola (up to 4-leaf)
cow cockle (up to 4-leaf)	volunteer flax
flixweed (up to 4-leaf)	volunteer rapeseed (up to 4-leaf)
green smartweed (up to 4-leaf)	volunteer sunflower (up to 4-leaf)
hemp-nettle	wild buckwheat
lady's thumb (up to 4-leaf)	wormseed mustard
night flowering catchfly (up to 4-leaf)	perennial sow thistle (top growth control only)
pale smartweed (up to 4-leaf)	

NOTE: IPCO EMIT activity is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24 °C. Reduced activity will occur when temperatures are below 8 °C or above 27 °C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, e.g. drought, heat or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.

Section 2 - IPCO EMIT in Tank Mix with other herbicides

TANK MIXTURES:

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

IPCO EMIT can be tank mixed with the following herbicides:

1. Nufarm Tralkoxydim Liquid or Achieve Liquid Herbicide (for wheat and barley only)
2. Cordon or Puma Advance Herbicide (for spring wheat, durum wheat and spring barley only)
3. Signal Herbicide or Nufarm Clodinafop Herbicide (for spring wheat and durum wheat only)
4. Axial 100EC Herbicide (for spring wheat and barley only)
5. Everest 2.0 Herbicide (for spring wheat and durum wheat only)
6. Varro Herbicide (for wheat only)
7. Simplicity (for spring wheat and durum wheat only)
8. IPCO EMIT plus Nufarm Boost Herbicide – for suppression of Narrow-leaved Hawk's beard

IPCO EMIT + Nufarm Boost Herbicide (wheat (spring, durum and winter), barley and oats only)	
Weeds Controlled:	See Section 1, plus suppression of Narrow-leaved Hawk's beard
Crop Treatment Stage:	From the 2-leaf to early flag-leaf stage.
Application Method:	Ground application only. DO NOT APPLY USING AERIAL APPLICATION EQUIPMENT. Apply in at least 100 litres of water per hectare.
Application Rate:	0.625 -1.25 L/ha IPCO EMIT + 6.7 g/ha Nufarm Boost Herbicide + surfactant as required on the Nufarm Boost Herbicide label directions.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact company representatives at 1-204-233-3461 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

®,™ Products are trademarks of their respective companies.

<p>NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE APPLICATIONS:</p> <p>The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Interprovincial Cooperative Ltd. under the User Requested Minor Use Label Expansion program. For these uses, Interprovincial Cooperative Ltd. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application</p>
<p>BROMEGRASS, HYBRID, MEADOW, SMOOTH (SEEDLING AND ESTABLISHED) GROWN FOR SEED</p> <p>DIRECTIONS FOR USE</p> <p>ALWAYS REFER TO THE PRODUCT LABEL FOR FURTHER INFORMATION ON WEEDS CONTROLLED, APPLICATION DIRECTIONS, AND USE PRECAUTIONS.</p> <p>Apply this product at the 3 to 6 leaf stage for a newly established stand and up to the first node for an established stand. Apply when weeds are at the growth stages listed on the product label. Use a single application of 0.625 to 1.25 L/ha.</p>
<p>ANNUAL CANARYGRASS</p> <p>DIRECTIONS FOR USE</p> <p>ALWAYS REFER TO THE PRODUCT LABEL FOR FURTHER INFORMATION ON WEEDS CONTROLLED, APPLICATION DIRECTIONS, AND USE PRECAUTIONS.</p>

ANNUAL CANARYGRASS

Rate: Apply 625 mL to 1.25 L of IPCO EMIT in 50-100 L spray volume per hectare.

Timing: Annual canarygrass may be treated from the 3 to 5 leaf stage. Application before the 2-leaf stage may injure the crop. Application must be made before the crop shields the weeds. Only weeds emerged at the time of application will be controlled. Apply when weeds are in the seedling stage (up to 6 leaf stage) and actively growing.

Do not apply more than one treatment per year. Do not harvest treated crop within 60 days of after application. Annual canarygrass seeds can be harvested for human consumption. Annual canarygrass is NOT to be cut for feed or grazed.