

2023-3830

2023-10-30

Cyclone PLUS

Bactericide and Fungicide

For suppression of powdery mildew on greenhouse cucumbers/ field cucurbits – Crop subgroup 9B (including cucumbers, squashes and pumpkins), on cannabis (marihuana) produced commercially indoors and field-grown, on greenhouse and outdoor roses and greenhouse and outdoor ornamentals including cut flowers, on grapes and on hemp; bacterial canker on greenhouse and field tomatoes; black spot on greenhouse and outdoor roses; fire blight on apples.

COMMERCIAL

Solution

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

ACTIVE INGREDIENTS:

Citric Acid..... 1.41%
Lactic Acid..... 2.54%

Warning, contains the allergen milk

REGISTRATION NUMBER: 34762
PEST CONTROL PRODUCTS ACT

DANGER – EYE AND SKIN IRRITANT

NET CONTENTS : 500 mL -1000L

Registered by:
AEF Global Inc.
440, 3e avenue, suite 100, Lévis, QC, G6W 5M6
(418) 838-4441
Toll free help-line: 1-866-622-3222
www.aefglobal.com

2023-3830

2023-10-30

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Severely irritating to the eye. **DO NOT** get in eyes. May irritate the skin. Avoid contact with skin. Avoid breathing mist. Wear a long-sleeved shirt, long pants, protective eyewear (goggles), chemical resistant gloves, socks and shoes when handling, mixing, loading or applying, and during all clean-up and repair activities. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours or until sprays have dried, unless wearing appropriate personal protective equipment including long-sleeved shirt, long pants, chemical resistant gloves, socks and shoes.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

2023-3830

2023-10-30

This product may cause injury to other plants if not mixed and applied properly. Follow instructions for mixing and applying carefully.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

To ensure efficiency, postpone application if possible when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

DIRECTIONS FOR USE

Shake well before use. Foliar treatment for crops listed below.

Crop	Disease	Rate/Concentration	Interval of application
Greenhouse Tomatoes	Bacterial canker (<i>Clavibacter michiganensis</i>)	2.4% (v/v) (e.g. 12 L in 500 L water/ha)	5 – 10 days
Field Tomatoes	Bacterial canker (<i>Clavibacter michiganensis</i>)	2.4% (v/v) (e.g. 12 L in 500 L water/ha)	5 – 10 days
Greenhouse Cucumbers	Powdery mildew (<i>Podosphaera xanthii</i> ; <i>Erysiphe cichoracearum</i> , syn. <i>Golovinomyces cichoracearum</i>)	1.2% - 2.0 % (v/v)	7 – 10 days
Field Cucurbits: Crop Subgroup 9B* (including squashes, pumpkins and cucumbers)	Powdery mildew (<i>Podosphaera xanthii</i>)	1.7% – 3.2% (v/v) (e.g. 12 L to 16L in 500 to 700 L water/ha)	7 – 10 days
Greenhouse and Outdoor Roses	Black spot (<i>Diplocarpon rosae</i>)	1.5 – 2.5% (v/v), spray volume of 800 L/ha	5 – 10 days
	Powdery mildew (<i>Sphaerotheca pannosa</i>)		7 – 10 days
Greenhouse and Outdoor ornamentals including cut flowers	Powdery Mildew (<i>Podosphaera fusca</i> syn. <i>Podosphaera xanthii</i> ; <i>Golovinomyces cichoracearum</i> syn. <i>Erysiphe cichoracearum</i>)	2.0% (v/v) (20 L in 1000 L water), spray volume of 1000 L/ha	7 days
Cannabis (marihuana) produced commercially indoors and field grown)	Powdery mildew (<i>Sphaerotheca macularis</i>)	1.2% (v/v) (12 L in 1000 L water/ha)	7 - 10 days
Hemp	Powdery mildew (<i>Sphaerotheca macularis</i>)	1.2% (v/v) (12 L/1000 L water), spray volume of 1000 L/ha	7 -10 days

2023-3830

2023-10-30

Apples	Fire blight (<i>Erwinia amylovora</i>)	1.4% (v/v) (14 L/1000 L water)	7 – 10 days Apply before wetting events when conditions are conducive to infection.
Grapes	Powdery mildew (<i>Uncinula necator</i>)	1.7 – 2.0%	7 – 10 days

* Crop Subgroup 9B includes the following crops: Chayote (fruit); Chinese waxgourd (Chinese preserving melon); Cucumber; Gherkin; Gourd, edible (*Lagenaria* spp.) (includes hyotan, cucuzza), (*Luffa acutangula*, *L. cylindrica*) (includes hechima and Chinese okra); *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); Pumpkin; Squash, summer (*Cucurbita pepo* var. melopepo) (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); Squash, winter (*Cucurbita maxima*, *C. moschata*) (includes butternut squash, calabaza, hubbard squash); (*C. mixta*, *C. pepo*) (includes acorn squash, spaghetti squash).

Where applicable, under conditions of moderate to high disease pressure, use the higher rate and/or shorter application interval. For maximum effectiveness, apply prior to, or at the early stages of disease development. Apply in sufficient volume to ensure full coverage for effective control. Both sides of leaves should be covered with the product.

Use nozzles that will provide excellent coverage of the plant foliage (ex: hollow cone nozzle). The spray quality should be very fine.

Begin applications when conditions are conducive to disease development.

The pre-harvest interval (PHI) is 0 (zero) day.

Cyclone PLUS may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact AEF Global Inc. technical services at servtechserv@aefglobal.com for information before applying any tank mix that is not specifically recommended on this label.

Do not mix with alkaline products. This product contains weak acids and could react with products containing significant amounts of carbonates. It is not recommended to tank mix Cyclone PLUS with copper-based products unless you have conducted a small scale phytotox test. AEF Global recommends conducting a jar test with any product mixes before any tank mix.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT apply by air.

2023-3830
2023-10-30

DO NOT allow effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.

STORAGE

Store this product away from food and feed. Keep in a cool place, away from direct sun rays. After extended storage, shake or stir the product thoroughly to assure a uniform suspension. Keep from freezing. Store in original container out of reach of children.

DISPOSAL

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.