TRICO® GARDEN

REPELLENT FOR PREVENTING FEEDING DAMAGE CAUSED BY DEER

EMULSIFIABLE CONCENTRATE

DOMESTIC

ACTIVE INGREDIENT:	
Sheep fat6	.4%

Contains Bronopol at 0.022%, 2-Methyl-4-isothiazolin-3-one at 0.0007% and 5-Chloro-2-methyl-4-isothiazolin-3-one at 0.0021% as preservatives

REGISTRATION NO. 34464 PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 0.25 L to 10 L

Lot/batch No.:

Kwizda Agro GmbH Universitätsring 6 A-1010, Vienna Austria

Ron W. Kobylnyk 2E - 9851 Second Street Sidney, B.C. Canada V8L 3C7

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

May cause mild eye and skin irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid breathing spray mists.

Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Wear rubber gloves and protective eyewear when handling, mixing/loading or applying the product, and during all clean-up/repair activities.

DO NOT allow children, adults or pets to enter treated areas until sprays have dried.

Do not apply this product in a way that will contact other people/domestic pets, either directly or through drift.

Do not contaminate food or any body of water.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

Store this product away from food or feed.

DISPOSAL:

DO NOT reuse the empty containers. Dispose in household garbage. Unused or partially used products should be disposed at provincially or municipally designated hazardous waste disposal sites.

PRODUCT INFORMATION

TRICO® GARDEN is a scent and taste repellent that prevents damage caused by deer that feed on outdoor ornamental plants (including flowers, shrubs and trees), apple trees and grapevines.

TRICO® GARDEN provides effective control for up to 28 days, depending on deer pressure and time of year.

Drying period depends on weather conditions. Under low humidity conditions, drying is expected to take a maximum of 6 hours. Once dried, TRICO® GARDEN provides excellent rain resistance.

DIRECTIONS FOR USE

Limitations:

- DO NOT apply to any body of water.
- Avoid application of this product when winds are gusty.

For apple trees and grapevines, apply as foliar spray only before flowering.

- Use freshly mixed spray solution each time product is applied.
- DO NOT apply undiluted. DO NOT apply more than 8 ml of 1:3 spray solution (containing 2 ml TRICO® GARDEN concentrate) or more than 12 ml of 1:5 of spray solution (containing 2 ml TRICO® GARDEN concentrate) per m².
- Apply only at temperatures between 2° C and 30 ° C.
- Protect product from freezing. Freezing makes TRICO® GARDEN unusable and must be disposed in accordance with the DISPOSAL section of this label.
- Avoid application in direct sunlight on hot days to avoid damage to leaves. It is recommended to apply in early morning or late afternoon.

- DO NOT apply more than 6 applications of TRICO® GARDEN per year (4 in spring/summer and 2 in autumn/fall).
- DO NOT apply other pest control products (including deer repellents) or foliar fertilizers one week before or one week after application of TRICO® GARDEN.

Mixing and Dilution Instructions

Shake the container thoroughly before use. Use 1 part of TRICO® GARDEN (e.g., 250 ml) to 3 parts of water (e.g., 750 ml) for heavy browsing pressure, and 1 part of TRICO® GARDEN (e.g., 250 ml) to 5 parts of water (e.g., 1.25 L) for moderate to light browsing pressure.

Application Instructions:

For use on outdoor ornamentals (including flowers, trees and shrubs), apple trees and grapevines to prevent feeding damage by deer.

Use a hand-held, backpack or garden sprayer with a flat fan or cone nozzle.

Apply spray on dry plant leaves, stems, flowers, tree bark and branches that need protection until plants look wet. For low growing plants, spray soil around plant base as well. In case of fresh deer damage or rapid plant growth, re-apply after a minimum of 28 days.

TRICO® GARDEN at the labeled application concentration and rates has been tested and found not to damage the following ornamental plants: black pine, common hawthorn, daisy, dogwood, hawthorn, hornbeam, silver fir, larkspur, maple, forsythia, geranium, willow, laburnum, lilac, cedar, Norway spruce, oak, Oriental spruce, rose, salvia, pansy and weigela. TRICO® GARDEN has not been tested on all plant species or varieties. Before large-scale treatments, it is recommended to test the product on a small area of that plant type and observe for at least 24 hours for any plant damage. Use with caution on open blooms and sensitive varieties such as impatiens, fuchsia, and begonias. A white residue may be visible on the plant after application.

[for 250 ml size only]

Degree of	Application Rate
Browsing	
Heavy	1:3 dilution:
Browsing	Contents of 250 ml bottle mixed with 0.75 L of water makes 1 L of spray solution.
_	This amount of spray solution covers 125 m ² .
Light to	1:5 dilution:
Moderate	Contents of 250 ml bottle mixed with 1.25 L of water makes 1.5 L of spray
Browsing	solution. This amount of spray solution covers 250 m ² .

[for 500 ml size only]

Degree of Browsing	Application Rate
Heavy	1:3 dilution:

Browsing	Contents of 500 ml bottle mixed with 1.5 L of water makes 2 L of spray solution. This amount of spray solution covers 250 m ² .
Light to	1:5 dilution:
Moderate	Contents of 500 ml bottle mixed with 2.5 L of water makes 3 L of spray solution.
Browsing	This amount of spray solution covers 500 m ² .

[for 1 L size only]

Degree of	Application Rate
Browsing	
Heavy	1:3 dilution:
Browsing	Contents of 1 L bottle mixed with 3 L of water makes 4 L of spray solution. This
	amount of spray solution covers 500 m ² .
Light to	1:5 dilution:
Moderate	Contents of 1 L bottle mixed with 5 L of water makes 6 L of spray solution. This
Browsing	amount of spray solution covers 1000 m ² .

[for 2 L size only]

Degree of	Application Rate
Browsing	
Heavy	1:3 dilution:
Browsing	Contents of 2 L bottle mixed with 6 L of water makes 8 L of spray solution. This
	amount of spray solution covers 1000 m ² .
Light to	1:5 dilution:
Moderate	Contents of 2 L bottle mixed with 10 L of water makes 12 L of spray solution. This
Browsing	amount of spray solution covers 2000 m ² .

[for 4 L size only]

Degree of	Application Rate
Browsing	
Heavy	1:3 dilution:
Browsing	Contents of 4 L bottle mixed with 12 L of water makes 16 L of spray solution. This
_	amount of spray solution covers 2000 m ² .
Light to	1:5 dilution:
Moderate	Contents of 4 L bottle mixed with 20 L of water makes 24 L of spray solution. This
Browsing	amount of spray solution covers 4000 m ² .

[for 10 L size only]

Degree of	Application Rate
Browsing	

Heavy	1:3 dilution:
Browsing	Contents of 10 L bottle mixed with 30 L of water makes 40 L of spray solution. This
	amount of spray solution covers 5,000 m ² .
Light to	1:5 dilution:
Moderate	Contents of 10 L bottle mixed with 50 L of water makes 60 L of spray solution. This
Browsing	amount of spray solution covers 10,000 m ² .