FIESTA Lawn Weed Killer Granule

READY TO USE GRANULE HERBICIDE COMMERCIAL

Selective granular weed killer

ACTIVE INGREDIENT: Iron (present as ferric sodium EDTA trihydrate) ... 0.91%

REG. NO. 34624 P.C.P. ACT

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS 1 - 1000 kg

Registrant:

W. Neudorff GmbH KG
An der Muhle 3, 31860 Emmerthal, Germany
www.neudorffpro.com

Canadian Agent:

Neudorff North America PO Box 178 Brentwood Bay, BC V8M 1R3 Telephone: (250)652-5888 For use on residential, commercial, industrial and municipal lawns and turf (on rights of way, landscaping, non-crop areas, golf courses, parks, cemeteries and athletic fields) consisting of one or more of Kentucky bluegrass, perennial ryegrass, and fescue (tall fescue, chewings fescue, and/or creeping fescue) for the pre- and post-emergent control of dandelion (*Taraxacum officinale*), English daisy (*Bellis perennis*), false dandelion (*Hypochaeris radicata*), white clover (*Trifolium repens*), black medic (*Medicago lupulina*), bull thistle (*Cirsium vulgare*), Canada thistle (*Cirsium arvense*), common chickweed (*Stellaria media*), creeping buttercup (*Ranunculus repens*), slender speedwell (*Veronica filiformis*), narrow-leaved plantain (*Plantago lanceolata*), dove's-foot geranium (*Geranium molle*), wild geranium (*Geranium robertianum*), lawn burweed (*Soliva pterosperma*), moss and algae, and for the suppression of broad-leaved plantain (*Plantago major*). For pre-emergent control of crabgrass.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes or clothing. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, and socks and shoes when applying the product and during all clean-up and repair activities. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before reuse.

FIRST AID:

If swallowed:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Rate of Application	Weeds Controlled	(Optional column) Area Treated
50 g /m²	black medic (<i>Medicago lupulina</i>), slender speedwell (<i>Veronica filiformis</i>), wild geranium (<i>Geranium robertianum</i>) and moss. Weed seedlings of: false dandelion (<i>Hypochaeris radicata</i>), English daisy (<i>Bellis perennis</i>), dandelion (<i>Taraxacum officinale</i>) or white clover (<i>Trifolium repens</i>).	20 m²/ kg
80 g /m²	dandelion (<i>Taraxacum officinale</i>), English daisy (<i>Bellis perennis</i>), false dandelion (<i>Hypochaeris radicata</i>), white clover (<i>Trifolium repens</i>), bull thistle (<i>Cirsium vulgare</i>), common chickweed (<i>Stellaria media</i>), narrow-leaved plantain (<i>Plantago lanceolata</i>), dove's foot geranium (<i>Geranium molle</i>), lawn burweed (<i>Soliva pterosperma</i>), and algae. Pre-emergent crabgrass control	12.5 m²/ kg
110 g /m²	broad-leaved plantain (<i>Plantago major</i>) (suppression only), creeping buttercup (<i>Ranunculus repens</i>) and Canada thistle (<i>Cirsium arvense</i>). For Canada thistle, best results are obtained when weeds are less than 15 cm tall and two applications may be required for control.	9 m²/ kg

One kg of product will treat between 9 and 20 m² (97- 215 ft²). Ensure lawn is wet from dew or water lawn lightly before application if temperature is below 25°C. If temperature is above 25°C do not pre-wet grass and apply product in the late afternoon or evening. Any standard dry spreader may be used. Uniform coverage is important.

Repeat treatment every 21-28 days up to 4 times per area per year.

Visible signs of control may be seen within several hours after application. Susceptible weeds, moss or algae will turn brown or black. Occasionally a darkening of the grass leaf blades can occur after treatment, however the grass will recover within a few days to several weeks. May stain thatch (dead grass under lawn); this does not affect the health of the grass. If turf is damaged, irrigate 24 hours after application, followed by mowing to remove the turf tip burn can help lawn recover faster.

Use Precautions

- Do not apply to drought stressed grass; ensure that the lawn is well watered prior to application.
- Do not apply to bentgrass. If unsure of grass sensitivity, test apply a small area of your lawn.
- Do not apply to newly seeded areas until four or more weeks after grass seedlings have emerged.

- Avoid applying to desirable plants. Any desirable plants or structures accidentally contacted should be rinsed immediately with water.
- Do not apply if rainfall is expected within 24 hours of application.
- Do not irrigate within 24 hours of application.
- Do not apply more than 4 times per area per year.

STORAGE: Store this product away from food or feed.

DISPOSAL

- 1. Thoroughly empty the contents of the container into the application device.
- 2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
- 3. Make the empty container unsuitable for further use.
- 4. Dispose of the container in accordance with provincial requirements.
- 5. For information on disposal of unused, unwanted product contact the manufacturer or provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Canadian Patent No. 2,477,493

Manufactured for: W. Neudorff GmbH KG An der Muhle 3, 31860 Emmerthal, Germany Canadian Office: Neudorff North America

PO Box 178, Brentwood Bay, BC, V8M 1R3 250-652-5888

 $\mbox{FIESTA}^{\mbox{\scriptsize 8}}$ is a registered trademark of W. Neudorff GmbH KG

Pull here to open and press to reseal.

Optional marketing statements: SELECTIVE BROADLEAF WEED KILLER

No unpleasant odor.

Made with iron.

See result within hours

Ready-to-use. No mixing necessary.

(1 kg) Will treat up to XXXm² (xxxx sq. ft.)

Pre- and post-emergent herbicide

Pre- and post-emergent weed control

Kills weeds and stops weed seeds from germinating

Kills weeds

Kills weeds and prevents new weeds from germinating

Kills the weeds you see, prevents new ones coming back

Kills the weeds you see, and the (seeds)(weeds) you don't

Granular herbicide for weed control in lawns