

2019-2707
2023-03-30

Dry Fiesta Broadleaf Weed Killer for Lawns RTU
GRANULE
HERBICIDE
DOMESTIC
READY TO USE

A selective granular herbicide for weed, moss, and algae control in lawns

ACTIVE INGREDIENT: Iron (present as ferric sodium EDTA trihydrate ... 0.91%

REG. NO. 34625 P.C.P. ACT

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS 1 - 6 kg

Registrant:

W. Neudorff GmbH KG
An der Muhle 3,
31860 Emmerthal, Germany
www.neudorffpro.com

Canadian Agent:

Neudorff North America
PO Box 178
Brentwood Bay, BC
V8M 1R3
Telephone: (250)652-5888

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes or clothing. Avoid breathing/inhaling dust. Wear gloves during mixing, loading, and repair/clean-up activities. Wear work shoes or boots when applying this product with a spreader.

FIRST AID:

If swallowed:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes:

Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

PESTS/SITES:

For pre- and post- emergent control of the following lawn weeds: Dandelion, English daisy, False dandelion, White clover, Black medic, Bull thistle, Canada thistle, Common chickweed, Creeping buttercup, Slender speedwell, Narrow-leaved plantain, Broad-leaved plantain (suppression only), Dove's-foot geranium, Wild geranium, Lawn burweed, Moss, and Algae. For pre-emergent control of crabgrass.

For use on lawn turf consisting of one or more of Kentucky bluegrass, perennial ryegrass, and fescue (tall fescue, chewing fescue, and/or creeping fescue).

DIRECTIONS FOR USE:

DO NOT apply to any body of water.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Post-Emergent Control: Apply at a rate of 50-110 g/m². The standard rate for this product is 80 g/m². Use the higher rate (110 g/m²) on the following tough perennial weeds: broad-leaved plantain (suppression only), creeping buttercup, and Canada thistle. For Canada thistle, best results are obtained when weeds are less than 15 cm tall. Use the lower rate (50 g/m²) on moss and the following easy to control weeds: black medic, slender

speedwell, wild geranium or on weed seedlings of false dandelion, English daisy, dandelion or white clover.

How to Apply:

Contents may have settled during shipment; invert bag several times to ensure product is mixed.

Ensure lawn is wet from dew, or water lawn lightly before application if temperature is below 25°C. If temperature is above 25°C do not pre-wet grass and apply product in the late afternoon or evening. For best results, use when temperatures are between 10°C and 30°C, and apply 2-3 days after lawn is mowed.

Use rotary or handheld spreader set to the correct application rate. See table below.

Uniform coverage is important. Uneven application may cause turf damage. If using a handheld spreader, hold spreader 30cm above the ground. If using a walk behind spreader, walk at a steady pace slightly overlapping each pass.

Reapply after 3-4 weeks for best results.

Pre-Emergent Control: For pre-emergent weed control, apply when soil temperatures are 10–15 °C. For most home lawns, the standard application rate of 80 g/m² will effectively prevent new weeds from emerging. If the lawn has a history of heavy weed infestations, applying 110 g/m² may improve control. Reapply after 3 – 4 weeks for best results.

| Weed Species | Rate | Scotts® Rotary Spreaders (All Models) | Scotts® Elite Spreader | Scotts® Hand-Held Spreader (All models) | Scotts® Wizz® Year-Round Spreader |
|------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------|-------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------|
| moss, black medic, slender speedwell, wild geranium or on weed seedlings of false dandelion, English daisy, dandelion or white clover. | 50 g/m ² | Set dial to xxxxx setting. To apply product begin walking, then squeeze and hold trigger. | Set dial to xxxxx setting. To apply product begin walking, then squeeze and hold trigger. | Adjust spreader selector setting to xxxxx. Pull trigger and turn crank clockwise while walking at normal speed across area to be covered with product | Set dial to xxxxx setting. To apply product, begin walking at a steady pace and squeeze trigger. |
| Dandelion, Bull thistle, Common chickweed, Narrow-leaved plantain, Dove's-foot geranium, Lawn burweed, and Algae. Pre-emergent control of crabgrass. | 80 g/m ² | Set dial to XXX setting. To apply product begin walking, then squeeze and hold trigger. | Set dial to xxxxx setting. To apply product begin walking, then squeeze and hold trigger. | Adjust spreader selector setting to xxxxx. Pull trigger and turn crank clockwise while walking at normal speed across area to be covered with product | Set dial to xxxxx setting. To apply product, begin walking at a steady pace and squeeze trigger. |
| broad-leaved plantain (suppression only), creeping buttercup, and Canada thistle | 110 g/m ² | Set dial to xxxxx setting. To apply product begin walking, then squeeze and hold trigger. | Set dial to xxxxx setting. To apply product begin walking, then squeeze and hold trigger. | Adjust spreader selector setting to xxxxx. Pull trigger and turn crank clockwise while walking at normal speed across area to be covered with product | Set dial to xxxxx setting. To apply product, begin walking at a steady pace and squeeze trigger. |

What to expect:

Visible signs of control may be seen within several hours after application.

Susceptible weeds, moss or algae will turn brown or black. Occasionally a darkening of the grass leaf blades can occur after treatment; however the grass will recover within a few days to several weeks. If turf is damaged, irrigation 24 hours after application, followed by mowing to remove the turf tip burn can help lawn recover faster.

May stain thatch (dead grass under lawn); this does not affect the health of the grass.

Use Precautions:

- Do not apply to drought stressed grass; ensure that the lawn is well watered prior to application, but do not apply when grass is wet if daytime temperatures are above 25°C.
- Do not apply when the daytime temperature will exceed 30°C.
- Do not apply in windy conditions
- Do not apply to bent grass. If unsure of grass sensitivity, test apply to a small area of your lawn.
- Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk or street back onto the treated area.
- Avoid contact with desirable plants. Any desirable plants or structures accidentally contacted should be rinsed immediately with water.
- Do not apply if rainfall is expected within 24 hours of application.
- Do not irrigate within 24 hours of application.
- Do not apply more than 4 times per area per year.

STORAGE:

Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals. Store unused material, tightly closed, in original container only, away from open flame. Store this product away from food or feed.

DISPOSAL:

Do not reuse the empty containers. Dispose in household garbage. Unused or partially used products should be disposed at provincially or municipally designated hazardous waste disposal sites.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

[optional marketing statements:

SELECTIVE BROADLEAF HERBICIDE

People and pets can enter treated area after product is applied.

No unpleasant odor.

Made with iron.

See results within hours

Visible results within hours

KILLS WEEDS, NOT LAWNS

Ready-to-use

(1 kg) Will treat up to xxx m² (xxxx sq. ft)

Second application may be required

Pre- and post-emergent herbicide

Pre- and post-emergent weed control

Kills weeds and stops weed seeds from germinating

Kills weeds

Kills weeds and prevents new weeds from germinating

Kills the weeds you see, prevents new ones coming back

Kills the weeds you see, and the (seeds)(weeds) you don't