

2021-6643
2023-07-19

GROUPS	1	4	7	12	U17	FUNGICIDES
---------------	----------	----------	----------	-----------	------------	-------------------

VIBRANCE® TOTAL

SEED TREATMENT FUNGICIDE

SUSPENSION

COMMERCIAL

A seed treatment product for control of seed- and soil-borne diseases in labeled crops.

ACTIVE INGREDIENTS:

Thiabendazole	4.30%
Sedaxane.....	1.43%
Metalaxyl-M and S-isomer	1.07%
Fludioxonil.....	0.72%
Picarbutrazox.....	0.72%

Contains 1,2-benzisothiazolin-3-one at 0.0097%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00058%, 2-methyl-4-isothiazolin-3-one at 0.00019%, and 2-bromo-2-nitropropane-1,3-diol at 0.016% as preservatives.

**READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

Warning, contains the allergen soy

REGISTRATION NO: **34890**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **1 L - 1050 L**

On-farm or commercial seed treatment (facilities and mobile treaters) with open or closed transfer treatment equipment is permitted.

ALL SEED TREATED WITH THIS PRODUCT MUST BE CONSPICUOUSLY COLOURED

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, ON N1G 4Z3
Telephone: 1-877-964-3682

Label

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

1.0 FIRST AID

IF POISONING IS SUSPECTED, IMMEDIATELY contact a physician or a poison control centre. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

2.0 TOXICOLOGICAL INFORMATION

Treat symptomatically.

3.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS. May irritate eyes. Avoid contact with eyes. Wash hands and face after handling and before eating, using tobacco (including smoking), applying cosmetics, or using the toilet. Avoid contamination of feed and foodstuffs.

Do not contaminate food, feed or domestic water supplies.

Apply only in a way that this product will not contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area when seed is being treated or bagged.

Treat seed in a well-ventilated area.

Keep treated seeds out of reach of children and animals.

NOTE: In case of accident, take the product label to the emergency facility or physician.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

4.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Seed Type	Tasks	PPE/ Engineering Controls
For Commercial Seed Treatment (including facility workers and mobile treaters)		
<i>All seed types</i>	Mixing, loading, treating, calibrating, bagging, sewing seed bags, stacking and any other activities involving handling treated seeds.	Open or closed transfer. During mixing, loading, treating and calibrating, bagging, sewing seed bags, stacking and any other activities involving handling treated seeds wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested
	Cleaning and repairing	During cleaning and repairing, wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested.
For On-Farm Seed Treatment		
<i>All seed types</i>	Mixing, loading, treating, calibrating, cleaning, repairing and any other activities involving planting and handling of treated seeds	Open or closed transfer. During mixing, loading, treating, calibrating, cleaning, repairing and any other activities involving planting and handling of treated seeds, wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks, and shoes. In addition, wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested when transferring treated seed to the planter/seedler.
For Planting and Handling Treated Seed		
<i>All seed types</i>	Planting	During planting and handling of treated seeds and any other activities involving calibrating, clean-up, and repair wear chemical resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, and shoes. In addition, wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested when transferring treated seed to the planter/seedler. Gloves are not required within a closed cab.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

5.0 ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms.

Toxic to birds and small wild mammals. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

6.0 USE RESTRICTIONS

1. Do not use treated seed for food, feed or oil purposes.
2. Store away from food and feed.
3. Do not plant any crop other than cereals, soybeans, members of Crop Subgroup 6C (Dried shelled peas and beans, chickpeas, lentils and faba beans) or members of Crop Subgroup 20A (Canola/rapeseed subgroup), within 60 days to fields in which treated seeds were planted.
4. Do not graze or feed livestock on treated area for 60 days after planting dried shelled pea and bean crops, chickpeas, lentils and faba beans.
5. **All containers or packages containing treated seed for sale or use in Canada must be labeled or tagged as follows:** "This seed has been treated with the fungicides thiabendazole, sedaxane, metalaxyl-M and S-isomer, fludioxonil, and picarbutrazox. Wear chemical resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, and shoes during planting and handling of treated seeds and any other activities involving calibrating, clean-up, and repair. Wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested when transferring treated seed to the planter/seeder. This product is toxic to aquatic organisms. Do not use for food, feed or oil processing. Store away from food and feed. Dispose of all excess treated seed. Dispose of seed packaging in accordance with local requirements. Toxic to birds and small wild mammals. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface."

7.0 STORAGE

Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated area away from other pesticides, feed and foodstuffs, and out of the reach of children and animals. To prevent contamination, store this product away from seed, food or feed.

Ideal storage temperature is above freezing and below 30°C. If the product should freeze, bring the product back to room temperature and ensure the contents are mixed well prior to application.

8.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

For Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers:

Do not reuse this container for any purpose. For disposal, the empty container may be returned to the point of purchase (distributor/dealer).

For Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

VIBRANCE® is a trademark of a Syngenta Group Company.

GROUPS	1	4	7	12	U17	FUNGICIDES
---------------	----------	----------	----------	-----------	------------	-------------------

VIBRANCE® TOTAL

SEED TREATMENT FUNGICIDE

SUSPENSION

COMMERCIAL

A seed treatment product for control of seed- and soil-borne diseases in labeled crops.

ACTIVE INGREDIENTS:

Thiabendazole	4.30%
Sedaxane.....	1.43%
Metalaxyl-M and S-isomer	1.07%
Fludioxonil.....	0.72%
Picarbutrazox.....	0.72%

Contains 1,2-benzisothiazolin-3-one at 0.0097%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00058%, 2-methyl-4-isothiazolin-3-one at 0.00019%, and 2-bromo-2-nitropropane-1,3-diol at 0.016% as preservatives.

READ THE LABEL AND THIS PAMPHLET BEFORE USING KEEP OUT OF REACH OF CHILDREN

Warning, contains the allergen soy

REGISTRATION NO: **34890**
PEST CONTROL PRODUCTS ACT

On-farm or commercial seed treatment (facilities and mobile treaters) with open or closed transfer treatment equipment is permitted.

ALL SEED TREATED WITH THIS PRODUCT MUST BE CONSPICUOUSLY COLOURED

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, ON N1G 4Z3
Telephone: 1-877-964-3682

Pamphlet

Table of Contents

	<u>Section Number</u>
NOTICE TO USER	1.0
FIRST AID	2.0
TOXICOLOGICAL INFORMATION	3.0
PRECAUTIONS.....	4.0
PERSONAL PROTECTIVE EQUIPMENT	5.0
ENVIRONMENTAL PRECAUTIONS	6.0
USE RESTRICTIONS	7.0
STORAGE	8.0
DISPOSAL	9.0
PRODUCT INFORMATION	10.0
DIRECTIONS FOR USE	11.0
General Information	11.1
Seed Treatment and Inoculants.....	11.2
CROP USE DIRECTIONS	12.0
Dried Shelled Beans.....	12.1
Dried Shelled Peas	12.2
Chickpeas	12.3
Lentils	12.4
Lupins and Faba Beans	12.5
Tank Mixes.....	12.6
RESISTANCE MANAGEMENT RECOMMENDATIONS	13.0

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IF POISONING IS SUSPECTED, IMMEDIATELY contact a physician or a poison control centre. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

3.0 TOXICOLOGICAL INFORMATION

Treat symptomatically.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS. May irritate eyes. Avoid contact with eyes. Wash hands and face after handling and before eating, using tobacco (including smoking), applying cosmetics, or using the toilet. Avoid contamination of feed and foodstuffs.

Do not contaminate food, feed or domestic water supplies.

Apply only in a way that this product will not contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area when seed is being treated or bagged.

Treat seed in a well-ventilated area.

Keep treated seeds out of reach of children and animals.

NOTE: In case of accident, take the product label to the emergency facility or physician.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Seed Type	Tasks	PPE/ Engineering Controls
For Commercial Seed Treatment (including facility workers and mobile treaters)		
<i>All seed types</i>	Mixing, loading, treating, calibrating, bagging, sewing seed bags, stacking and any other activities involving handling treated seeds.	Open or closed transfer. During mixing, loading, treating and calibrating, bagging, sewing seed bags, stacking and any other activities involving handling treated seeds wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested
	Cleaning and repairing	During cleaning and repairing, wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested.
For On-Farm Seed Treatment		
<i>All seed types</i>	Mixing, loading, treating, calibrating, cleaning, repairing and any other activities involving planting and handling of treated seeds	Open or closed transfer. During mixing, loading, treating, calibrating, cleaning, repairing and any other activities involving planting and handling of treated seeds, wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks, and shoes. In addition, wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested when transferring treated seed to the planter/seedler.
For Planting and Handling Treated Seed		
<i>All seed types</i>	Planting	During planting and handling of treated seeds and any other activities involving calibrating, clean-up, and repair wear chemical resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, and shoes. In addition, wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested when transferring treated seed to the planter/seedler. Gloves are not required within a closed cab.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for

washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

6.0 ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms.

Toxic to birds and small wild mammals. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

7.0 USE RESTRICTIONS

1. Do not use treated seed for food, feed or oil purposes.
2. Store away from food and feed.
3. Do not plant any crop other than cereals, soybeans, members of Crop Subgroup 6C (Dried shelled peas and beans, chickpeas, lentils and faba beans) or members of Crop Subgroup 20A (Canola/rapeseed subgroup), within 60 days to fields in which treated seeds were planted.
4. Do not graze or feed livestock on treated area for 60 days after planting dried shelled pea and bean crops, chickpeas, lentils and faba beans.
5. **All containers or packages containing treated seed for sale or use in Canada must be labeled or tagged as follows:** “This seed has been treated with the fungicides thiabendazole, sedaxane, metalaxyl-M and S-isomer, fludioxonil, and picarbutrazox. Wear chemical resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, and shoes during planting and handling of treated seeds and any other activities involving calibrating, clean-up, and repair. Wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested when transferring treated seed to the planter/seeder. This product is toxic to aquatic organisms. Do not use for food, feed or oil processing. Store away from food and feed. Dispose of all excess treated seed. Dispose of seed packaging in accordance with local requirements. Toxic to birds and small wild mammals. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.”

8.0 STORAGE

Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated area away from other pesticides, feed and foodstuffs, and out of the reach of children and animals. To prevent contamination, store this product away from seed, food or feed.

Ideal storage temperature is above freezing and below 30°C. If the product should freeze, bring the product back to room temperature and ensure the contents are mixed well prior to application.

9.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

For Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers:

Do not reuse this container for any purpose. For disposal, the empty container may be returned to the point of purchase (distributor/dealer).

For Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

10.0 PRODUCT INFORMATION

VIBRANCE TOTAL is a ready-to-apply seed treatment product containing the fungicides, thiabendazole, sedaxane, metalaxyl-M and S-isomer, fludioxonil, and picarbutrazox. This product controls a broad spectrum of seed- and soil-borne diseases in listed dried shelled pea and bean crops, chickpeas, lentils, lupins and faba beans.

Note: Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigour and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination, reduced emergence and/or reduction of seed and seedling vigour. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of the seed before committing the total seed lot to a selected seed treatment. Due to seed quality and seed storage conditions beyond the control of Syngenta Canada Inc. no claims are made to guarantee the germination of carry-over treated seed.

11.0 DIRECTIONS FOR USE

11.1 General Information

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

VIBRANCE TOTAL is a seed treatment formulation for use in commercial seed treatment (in facilities or with mobile treaters) and for on-farm treatment using seed treatment equipment that accurately meters, mixes and applies a flowable seed treatment. The equipment must provide uniform coverage of VIBRANCE TOTAL on the seed. Uneven seed coverage may not give the desired level of disease control. Allow the seed to dry before bagging, storing or seeding.

Thoroughly mix the recommended amount of VIBRANCE TOTAL Seed Treatment with the required amount of water for the slurry treater and dilution rate to be used. Slurry volumes will vary depending on seed size. Follow the manufacturer's application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment.

When mixing with products from other manufacturers, the compatibility should be tested prior to use by conducting a jar test; mixing all intended seed treatments with the appropriate amount of water in a clear glass container, mix well, and allow to sit for one hour. Remix and observe for incompatibility.

11.2 Seed Treatment and Inoculants

VIBRANCE TOTAL is compatible with Rhizobium based inoculants. Please check with inoculant manufacturers for specific planting windows and methods of application prior to use. Consult the manufacturer of the application equipment planned to be used for suitability for this application and for instructions on operation and calibration of the equipment.

Seed treated with VIBRANCE TOTAL, or a combination of VIBRANCE TOTAL and seed inoculants, may not flow through a seed drill at the same rate as untreated seed. Recalibrate the seed drill before planting treated seed. **Mixing with inoculants may increase drying time while treating, extending the processing time.**

This product contains a pigment that will colour the treated seed; however, users are responsible for ensuring that the treated seed, when dried and ready for bagging, storage or seeding has an unnatural colour. If the pigment contained in the formulation does not colour the seed adequately, additional colourant must be added to the mixture while treating the seed.

12.0 CROP USE DIRECTIONS

For control of various seed-borne and soil-borne diseases on beans, lentils, lupins, faba beans, and peas, apply 325 mL of VIBRANCE TOTAL Fungicide per 100 kg seed.

12.1 Dried Shelled Beans

CROP	Dried Shelled Beans [<i>Phaseolus</i> spp. (field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); <i>Vigna</i> spp. (includes adzuki bean, blackeyed pea, catjang, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); guar (<i>Cyamopsis tetragonoloba</i>); lablab bean (<i>Lablab purpureus</i>)]
DISEASES CONTROLLED	Seed rot/pre-emergence damping-off and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Pythium</i> spp. and <i>Rhizoctonia</i> spp. Seedling blight caused by <i>Pythium</i> spp. and <i>Rhizoctonia</i> spp. Seed-borne <i>Sclerotinia sclerotiorum</i> Anthracnose caused by seed-borne <i>Colletotrichum</i> spp.
RATE	325 mL/100 kg seed

12.2 Dried Shelled Peas

CROP	Dried Shelled Peas (<i>Pisum</i> spp. and <i>Cajanus cajan</i>)
DISEASES CONTROLLED	Seed-borne ascochyta blight and foot rot caused by <i>Ascochyta pinodes</i> Seed rot/pre-emergence damping-off, post-emergence damping-off and seedling blight caused by <i>Fusarium</i> spp., <i>Pythium</i> spp. and <i>Rhizoctonia</i> spp. Seed-borne <i>Sclerotinia sclerotiorum</i>
RATE	325 mL/100 kg seed

12.3 Chickpeas

CROP	Chickpeas
DISEASES CONTROLLED	Seed-borne ascochyta blight caused by <i>Ascochyta rabiei</i> Seed rot/pre-emergence damping-off and post-emergence damping-off and seedling blight caused by <i>Fusarium</i> spp., <i>Pythium</i> spp. and <i>Rhizoctonia</i> spp. Seed rot and seedling blight caused by seed-borne <i>Botrytis</i> spp. Seed-borne <i>Sclerotinia sclerotiorum</i>
RATE	325 mL/100 kg seed

Lentils

CROP	Lentils
DISEASES CONTROLLED	Seed-borne ascochyta blight caused by <i>Ascochyta lentis</i> Seed rot/pre-emergence damping-off, post-emergence damping-off and seedling blight caused by <i>Fusarium</i> spp., <i>Pythium</i> spp. and <i>Rhizoctonia</i> spp. Seedling root rot caused by <i>Fusarium</i> spp. Seed-borne <i>Sclerotinia sclerotiorum</i> Seed rot and seedling blight caused by seed-borne <i>Botrytis</i> spp.
RATE	325 mL/100 kg seed

12.4 Lupins and Faba Beans

CROP	Lupins and Faba Beans
DISEASES CONTROLLED	Seed rot/pre-emergence damping-off, post-emergence damping-off and seedling blight caused by <i>Fusarium</i> spp., <i>Pythium</i> spp. and <i>Rhizoctonia</i> spp.
RATE	325 mL/100 kg seed

12.5 Tank Mixes

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including

Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Syngenta Canada Inc. at 1-877-SYNGENTA / 1-877-964-3682 for information before applying any tank mix that is not specifically recommended on this label.

Syngenta Canada Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) of all potential tank mixes under all environmental conditions or for all crop varieties. Tank mixes that are not specifically listed on this label should be tested on a small area first, under local conditions and using standard practices, to confirm the tank mix is suitable for widespread application.

To determine the physical compatibility of this product with other products, use a jar test.

Always read and follow label directions including WALES mixing order.

13.0 RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that VIBRANCE TOTAL contains Group 1, 4, 7, 12 and U17 fungicides. Any fungal population may contain individuals naturally resistant to VIBRANCE TOTAL and other Group 1, 4, 7, 12 and U17 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of VIBRANCE TOTAL or other Group 1, 4, 7, 12 and U17 fungicides with different groups that control the same pathogens.

Use tank mixtures with fungicide from a different group that is effective on the target pathogen when such use is permitted.

Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for resistance development. Notify Syngenta Canada Inc. if reduced sensitivity of the pathogen to VIBRANCE TOTAL is suspected.

If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information or to report suspected resistance, contact company representatives at 1-87-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.

VIBRANCE® is a trademark of a Syngenta Group Company.