

2024-1295
2024-09-13

Container Label

GROUP	2	HERBICIDE
-------	---	-----------

FBN ZETHMOX 70 WDG

Water-Dispersible Granular Herbicide
in water-soluble bags

COMMERCIAL (AGRICULTURAL)

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND
PEACE RIVER REGION OF BRITISH COLUMBIA ONLY

ACTIVE INGREDIENT: Imazamox..... 35% a.e.
Imazethapyr...35% a.e.

Warning, contains the allergen sulfites

REGISTRATION NO. 34895 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 20 g – 1.5 kg; 692 g (8 x 86.5g)

CAUTION – SKIN IRRITANT

FOR EMERGENCY MEDICAL ASSISTANCE (HUMAN OR ANIMAL)
CONTACT ROCKY MOUNTAIN POISON CONTROL AT 1-866-767-5041.
FOR CHEMICAL EMERGENCY ASSISTANCE (SPILL, LEAK, FIRE OR
ACCIDENT) CONTACT CHEMTREC AT 1-800-424-9300 (NORTH
AMERICA) OR 1-703-527-3887 (INTERNATIONAL).

DO NOT SELL INDIVIDUAL WATER-SOLUBLE BAGS SEPARATELY

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

Farmer's Business Network Canada, Inc.
PO Box 5607
High River, Alberta
Canada T1V 1M7
1-844-200-FARM (3276)

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

May irritate the skin. Avoid contact with skin

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours after application.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial and aquatic plants. Observe spray buffer zones specified under Directions for Use.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

DISPOSAL

1. When all the water-soluble bags are used, make the empty outer package unsuitable for further use.
2. Dispose of the outer package in accordance with provincial requirements.
3. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

STORAGE

Store above 5°C.

Store this product away from food or feed. Store in a cool, dry place away from children, animals, or fertilizers. Keep unused water-soluble bags in resealed original container. Keep packages dry at all times. Keep packages out of rain. Keep from freezing.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

GROUP	2	HERBICIDE
-------	---	-----------

FBN ZETHMOX 70 WDG

Water-Dispersible Granular Herbicide
in water-soluble bags

COMMERCIAL (AGRICULTURAL)

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND
PEACE RIVER REGION OF BRITISH COLUMBIA ONLY

ACTIVE INGREDIENT: Imazamox..... 35% a.e.
Imazethapyr ... 35% a.e.

Warning, contains the allergen sulfites

REGISTRATION NO. 34895 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 20 g – 1.5 kg; 692 g (8 x 86.5 g)

CAUTION – SKIN IRRITANT

FOR EMERGENCY MEDICAL ASSISTANCE (HUMAN OR ANIMAL)
CONTACT ROCKY MOUNTAIN POISON CONTROL AT 1-866-767-5041.
FOR CHEMICAL EMERGENCY ASSISTANCE (SPILL, LEAK, FIRE OR
ACCIDENT) CONTACT CHEMTREC AT 1-800-424-9300 (NORTH
AMERICA) OR 1-703-527-3887 (INTERNATIONAL).

DO NOT SELL INDIVIDUAL WATER-SOLUBLE BAGS SEPARATELY

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

Farmer's Business Network Canada, Inc.
PO Box 5607
High River, Alberta
Canada
T1V 1M7
1-844-200-FARM (3276)

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

May irritate the skin. Avoid contact with skin.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours after application.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label, or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

DISPOSAL

1. When all the water-soluble bags are used, make the empty outer package unsuitable for further use.
2. Dispose of the outer package in accordance with provincial requirements.
3. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean -up of spills.

STORAGE

Store above 5 °C.

Store this product away from food or feed. Store in a cool, dry place away from children, animals, or fertilizers. Keep unused water-soluble bags in resealed original container. Keep packages dry at all times. Keep packages out of rain. Keep from freezing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial and aquatic plants. Observe spray buffer zones specified under Directions for Use.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

TANK MIXING

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Farmer's Business Network Canada, Inc. at 1-844-200-FARM (3276) for information before applying any tank mix that is not specifically recommended on this label.

For field peas, fenugreek (for seed uses and forage production only), Clearfield lentils¹ (lentil varieties with the Clearfield trait), Clearfield sunflowers¹ (sunflower varieties with the Clearfield trait), soybeans (Western Canada only), fababeans, seedling and established alfalfa, seedling clover (red, alsike, and sweet) and bird's foot trefoil grown for seed production

DIRECTIONS FOR USE (Prairie Provinces and Peace River Region of British Columbia)

FBN Zethmox 70 WDG is a selective herbicide that can be applied as an early post-emergence treatment in field peas, fenugreek (for seed uses only), **Clearfield** lentils¹, **Clearfield** sunflowers¹ (sunflower varieties with the **Clearfield** trait), soybeans, fababeans, seedling and established alfalfa, seedling clover (red, alsike, and sweet) and bird's foot trefoil grown for seed production. When FBN Zethmox 70 WDG is applied early post-emergence, absorption may occur through both the roots and foliage. Susceptible weeds stop growing and eventually die.

Apply only when the potential for drift to areas of human habitation or areas of human activity (houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment, and sprayer settings.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field Sprayer Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply by air.

Spray Buffer Zones

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas, and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands).

Method of Application	Crop	Spray Buffer Zones (metres) Required for the Protection of:	
		Freshwater Habitat	Terrestrial Habitat
Field sprayer	All crops	1	1

When tank mixes are permitted, consult the labels of the tank mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray drift buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

Pre-harvest Interval: After 60 days, field peas, fenugreek (for seed uses and forage production only), sunflowers, lentils, and fababeans can be harvested. After 85 days, soybeans can be harvested. FBN Zethmox 70 WDG must not be applied within 60 days of harvest of bird's foot trefoil.

Rainfastness Interval: FBN Zethmox 70 WDG is rainfast 3 hours after application.

DO NOT graze treated soybeans or cut for hay; sufficient data are not available to support such uses.

Field peas treated with FBN Zethmox 70 WDG may be fed to livestock 30 days after application.

Do not harvest forage or cut hay for forage.

Do not graze treated bird's foot trefoil or cut for hay. Do not use refuse or hay from treated bird's foot trefoil for animal feed or forage.

APPLY FBN Zethmox 70 WDG in a water volume of 50–100 L/ha using ground equipment only.

Do not apply more than once per year.

For Field Peas

Apply FBN Zethmox 70 WDG as a single post-emergent application at 30 g a.e./ha (43 g product/ha). Application to field peas should be from 1–6 true leaf stage and after weeds have emerged. Apply when broadleaf weeds are from the cotyledon to 4-leaf stage and when grassy weeds are at the 1–4 true leaf or early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.

For Fenugreek (for seed uses only)

Apply FBN Zethmox 70 WDG as a single post-emergent application at 30 g a.e./ha (43 g product/ha). Application to fenugreek should be from the 1- to 4-leaf stage and after weeds have emerged. Apply when grassy and broadleaf weeds are at the 1–4 true leaf stage.

NOTE TO USER

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Farmer's Business Network Canada, Inc. under the User Requested Minor Use Label Expansion program. For these uses, Farmer's Business Network Canada, Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

FOR FENUGREEK (Forage production)

Apply FBN Zethmox 70 WDG as a single post-emergent application at 30 g a.e./ha (43 g product/ha). Application to fenugreek should be from the 1- to 4-leaf stage and after weeds have emerged. Apply when grassy and broadleaf weeds are at the 1–4 true leaf stage.

Fenugreek forage is to be used for animal feed purposes only.

FBN Zethmox 70 WDG must not be applied within 60 days of harvest of fenugreek for forage production.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply in 50– 100 L/ha of water with MERGE Adjuvant at a rate of 0.5% v/v.

For Clearfield lentils¹ (lentil varieties with the Clearfield trait)

Apply FBN Zethmox 70 WDG as a single post-emergent application at 30 g a.e./ha (43 g product/ha). Application to **Clearfield** lentils¹ should be from 1–9 node stage and after weeds have emerged. Apply when grassy and broadleaf weeds are at the 1–4 true leaf stage. Grassy weeds can be treated up to early tillering provided they are not beyond the 4 true leaf stage.

For Clearfield sunflowers¹ (sunflower varieties with the Clearfield trait)

Apply FBN Zethmox 70 WDG as a single post-emergent application at 30 g a.e./ha (43 g product/ha).

Application should be made from the 2- to 8-leaf stage of the sunflower crop and after the weeds have emerged. Grassy and broadleaf weeds should be at the 1–4 true leaf stage. Grassy weeds can be treated up to early tillering provided they are not beyond the 4 true leaf stage.

FBN Zethmox 70 WDG should only be used on **Clearfield** sunflowers¹ where production is not destined for export.

Initial transient crop yellowing may be observed after application, but this is outgrown and should not affect yield.

For Soybeans

Apply FBN Zethmox 70 WDG as a single post-emergent application at 30 g a.e./ha (43 g product/ha). Application to soybeans should be from the 1-3 true leaf stage and after weeds have emerged. Apply when broadleaf weeds are from the cotyledon to 4-leaf stage and when grassy weeds are at the 1–4 true leaf or early tillering.

For Seedling and Established Alfalfa Grown for Seed

Apply FBN Zethmox 70 WDG as a single post-emergent application at 30 g a.e./ha (43 g product/ha). Apply by ground up to the 4-leaf stage of actively growing weeds; apply at the 1- to 4-leaf stage of alfalfa.

For Seedling Clover (red, alsike, and sweet) Grown for Seed Production

Apply FBN Zethmox 70 WDG as a single post-emergent application at 30 g a.e./ha (43 g product/ha) to seedling clover (red, alsike, and sweet). Apply by ground up to the 4-leaf stage of actively growing weeds; apply at the 1- to 4-leaf stage of seedling clover.

Injury to clover (red, alsike, and sweet) in the form of discolouration and stunting maybe observed shortly after application. However, this injury is typically outgrown and should not affect seed yield under appropriate field management.

For seed production only. Harvested crop not to be fed to livestock.

For Bird's Foot Trefoil (for seed production)

Apply FBN Zethmox 70 WDG as a single post-emergent application at 30 g a.e./ha (43 g product/ha). Apply by ground up to the 4-leaf stage of actively growing weeds; apply at the 1- to 4-leaf stage of bird's foot trefoil.

Early crop injury to bird's foot trefoil may be observed but this is outgrown and should not affect seed yield under appropriate field management.

NOTE TO USER

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Farmer's Business Network Canada, Inc. under the User Requested Minor Use Label Expansion program. For these uses, Farmer's Business Network Canada, Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

FOR FABABEANS

Apply FBN Zethmox 70 WDG as a single post-emergent application at 30 g a.e./ha (43 g product/ha) for control of labeled weeds. Application to fababeans should be from the 1- to 6-leaf stage and after weeds have emerged. Apply when grassy and broadleaf weeds are at the 1–4 true leaf stage.

APPLY FBN Zethmox 70 WDG in a water volume of 50–100 L/ha with MERGE® Adjuvant at a rate of 0.5% v/v.

Yellow flash or chlorosis may occur after application. The crop should be sprayed when actively growing and not under stress. Application should not be made to crops that are behind normal stage of development or whose maturity puts them at risk for fall frost.

FBN Zethmox 70 WDG must not be applied within 60 days of harvest of fababeans.

Make only one application of FBN Zethmox 70 WDG per year.

FBN Zethmox 70 WDG at 29 g/ha (20 g a.e./ha) will control only:

Redroot pigweed
Volunteer tame mustard
Wild mustard

FBN Zethmox 70 WDG at 43 g/ha (30 g a.e./ha) will control:

Chickweed	Barnyard grass
Cleavers	Green foxtail
Flixweed	Persian darnel
Green smartweed	Volunteer barley
Hemp-nettle*	Volunteer tame oats
Kochia**	Volunteer wheat
Lamb's-quarters***	Wild oats
Redroot pigweed	
Russian thistle****	
Shepherd's purse	
Stinkweed	
Stork's bill	
Volunteer canola (non-Clearfield canola only)	
Volunteer tame mustard	
Wild buckwheat*	
Wild mustard	

- * = suppression in field peas and **Clearfield** lentils¹
- ** = not controlled in **Clearfield** lentils¹; suppression in field peas and **Clearfield** sunflowers¹
- *** = suppression in field peas, **Clearfield** lentils,¹ and **Clearfield** sunflowers¹
- **** = suppression in **Clearfield** lentils¹

Apply in 50–100 L/ha of water with MERGE Adjuvant alone at a rate of 0.5% v/v.

For 1000 L of spray solution

USE:	SURFACTANT (L)	+ FERTILIZER (L)
MERGE® Adjuvant, MPOWER ASSASSIN® Adjuvant or ASSASSIN II Adjuvant	5.0	No fertilizer required

MIXING INSTRUCTIONS

Use 50–100 L/ha of water. Use a 50-mesh (or coarser) filter screen. Fill the spray tank with three-quarters of the required amount of clean water, start agitation, and continue agitation throughout the entire mixing and spraying procedure. Add the required amount of FBN Zethmox 70 WDG soluble bag(s) directly into the sprayer through the tank opening. Agitate for at least 10 minutes to dissolve the herbicide. After the herbicide is dissolved, continue agitation and add the required amount of MERGE Adjuvant, MPOWER ASSASSIN Adjuvant, or ASSASSIN II Adjuvant. If excess foaming occurs, a silicone anti-foaming agent may be added (e.g., HALT). Complete filling the tank to the desired level with water. If agitation is stopped for more than 5 minutes, re-suspend spray solution by full agitation prior to commencing spraying again. Between loads of FBN Zethmox 70 WDG, check in-line and nozzle screens, and rinse and clean if necessary. Upon completion of spraying FBN Zethmox 70 WDG, thoroughly flush tank, boom, hoses, and in-line and nozzle screens with clean water to avoid possible injury to other crops.

ROTATIONAL CROPS

There is the possibility of residual soil activity for FBN Zethmox 70 WDG the year following application.

Research studies have shown that the following crops may be safely planted the year following FBN Zethmox 70 WDG application:

- Field corn
- Field peas
- Clearfield** canola¹ (canola varieties with the **Clearfield** trait)
- Clearfield** lentils¹ (lentil varieties with the **Clearfield** trait)
- Lentils
- Spring wheat
- Durum wheat
- Tame oats
- Canary seed
- Spring barley
- Chickpeas

Research studies have shown that flax, canola, and sunflower may be safely planted **2 years** following FBN Zethmox 70 WDG application in all regions of Western Canada except the Peace River Region of Alberta and British Columbia. In the Peace River Region, observe a minimum 34-month recropping interval for non-**Clearfield** canola, flax, and non-**Clearfield** sunflower, and perform a field bioassay the year prior to planting to confirm normal growth of the crops. If the crops show signs of injury, delay planting by an additional year and perform an additional field bioassay the following year.

WARNING: Certain environmental conditions may delay the breakdown of herbicide residues in soil. These conditions include but are not limited to drought, extremes in soil pH, and excessive cold. Under these conditions, the level of phytotoxic herbicide residues present in the field the season following an application may result in an increased potential for injury to succeeding crops to occur. This potential for increased residues under these environmental conditions is not unique to any specific herbicide or herbicide group but is a property of those herbicides that persist in the soil and are dependent on soil microbial activity and other non-microbial processes (e.g., hydrolysis) to breakdown.

There are insufficient data for other follow crops. Conduct a field bioassay (a test strip grown to maturity) the year before growing any crop other than those listed above.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, FBN Zethmox 70 WDG is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to FBN Zethmox 70 WDG and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of FBN Zethmox 70 WDG or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields

by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact FBN at 1-844-200-FARM (3276).

Clearfield is a registered trademark of BASF Agrochemical Products B.V., all used with permission by BASF Canada Inc.

MERGE is a registered trademark of BASF Canada, Inc.

ASSASSIN is a trademark of AgraCity Crop & Nutrition, Ltd., or its affiliates.

® All other products listed are registered trademarks of their respective companies.

Pouch Label

FBN ZETHMOX 70 WDG

NET CONTENTS: 86.5 g

ACTIVE INGREDIENTS:

: Imazamox 35% a.e.
Imazethapyr 35% a.e.

Warning, contains the allergen sulfites

CAUTION - SKIN IRRITANT

KEEP OUT OF REACH OF

CHILDREN