

2023-6454
2024-06-25

This label has been updated according to the re-evaluation decision of (Z)-9-Tricosene RVD2023-13. While users are encouraged to follow this updated label immediately, the previously approved label is valid until 31/07/2025 in accordance with the phase out period set out in the re-evaluation/special review decision of (Z)-9-Tricosene RVD2023-13. This previously approved label will be provided upon request by emailing hc.pmra.info-arla.sc@canada.ca. In your email please include the product name and Registration number of the label you are requesting.

SOFAST Granular Fly Bait

Insecticide

AGRICULTURAL

FOR INDOOR USE AT LIVESTOCK FACILITIES ONLY

Attracts and Kills House Flies

ACTIVE INGREDIENT:

Imidacloprid.....0.5%
Muscalure.....0.1%

REGISTRATION NUMBER 33562 PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 10 g to 50 kg

SHARDA Cropchem Limited
2nd Floor, Prime Business Park
Dashrathlal Joshi Road
Vile Parle (West)
Mumbai - 400056, India

Canadian Agent:
SHARDA Cropchem Limited
63 Kingsview Blvd
Etobicoke, Ontario, CA
M9R1V1
1-844-810-5720
1-416-840-5639

GENERAL INFORMATION

- Attracts and kills house flies inside livestock facilities
- Ready-to-use bait (containing 0.5% imidacloprid)
- Contains a specially formulated mixture of attractants
- Contains the attractant muscalure
- Contains Bitrex® bittering agent to help prevent ingestion by animals, children, and pets.

DIRECTIONS FOR USE

Use SOFAST Granular Fly Bait as part of an overall integrated pest management program for house flies which may include:

- Proper sanitation and manure management.
- Environmental controls, such as proper light management and exclusion.
- Other registered pest control products.

DO NOT formulate this product into other end-use products.


HOW TO APPLY:

SOFAST Granular Fly Bait can be used:

1. Indoor broadcast treatment
2. Indoors using bait station

Indoor broadcast application is to broad expanses of indoor structural surfaces such as walls, floors, ceilings and indoor foundation walls/crawlspace.

TO ATTRACT AND KILL HOUSE FLIES	
WHERE TO APPLY	<p>FOR INDOOR USE ONLY.</p> <p>For use inside livestock facilities such as broiler houses, feedlots, livestock housing structures, and horse stables.</p> <p>DO NOT place bait in areas accessible to livestock, pets and children.</p> <p>Apply SOFAST Granular Fly Bait in locations where house flies congregate. In caged layer houses, apply only in the walkways. Avoid application in cool locations because house flies prefer warm sites for resting. Avoid application in areas with air movement such as near fans.</p> <p>When used in areas accessible to birds, bait must be placed in a tamper-resistant bait station that prevents birds from consuming the bait.</p>
WHEN TO APPLY	Time applications to begin at the start of the season before fly populations have reached their peak.
USE RATE	200 g SOFAST Granular Fly Bait /100 m ²

<p>METHODS OF APPLICATION</p>	<p><u>BROADCAST APPLICATION:</u> Scatter the bait at a rate of 200 g/100m² directly from the container onto dry level surfaces so that the individual granules lie near each other without forming small piles. DO NOT use this method of application in horse stables.</p> <p><u>BAIT STATIONS:</u> Place in tamper-resistant fly bait station. Add 50 g of bait per station. Use 1 bait station to every 25 m². Secure bait stations at least 1.25 m above the ground. Bait stations should be inaccessible to livestock, children and pets.</p> <p>Treated baits must be placed in tamper-resistant bait stations OR locations NOT accessible to children, pets, livestock or wildlife.</p> <p>Tamper-resistant bait stations must:</p> <ol style="list-style-type: none"> 1. be constructed of high-strength material that is hard for children or animals to destroy; 2. be constructed so that the bait cannot be shaken out; 3. have an entrance which prevents children and non-target animals from reaching bait; 4. have an access panel which fastens securely and locks; 5. be able to be securely fastened to a surface; and 6. be clearly labelled: DANGER POISON.
<p>RESTRICTIONS</p> 	<ul style="list-style-type: none"> • Treated baits must be placed in tamper-resistant bait stations OR locations NOT accessible to children, pets, livestock or wildlife. • DO NOT apply to overhead areas or in confined spaces without appropriate respiratory and eye protection. • Do not allow bait to come into contact with food or water, livestock feed or their water. • Do not apply to milking equipment. • Do not apply inside food processing plants, restaurants or other commercial facilities. • As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests. • Do not contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that SOFAST Granular Fly Bait contains a Group 4 insecticide. Any insect population may contain individuals naturally resistant to SOFAST Granular Fly Bait and other Group 4 insecticides. The resistant individuals may dominate the insect population if this group of insecticides is used repeatedly in the same location. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of SOFAST Granular Fly Bait or other Group 4 insecticides with different groups that control the same pests.

- Insecticide use should be based on an Integrated Pest Management (IPM) program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist for any additional pesticide resistance management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information and to report suspected resistance, contact a Sharda Cropchem Limited representative at 1-844-810-5720.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

DO NOT APPLY IN THE PRESENCE OF CHILDREN

DO NOT APPLY TO AREAS ACCESSIBLE TO CHILDREN

Wear a long-sleeved shirt, long pants and chemical resistant gloves during application and clean-up activities.

Avoid contamination of food and feed.

Avoid contact with eyes, skin or clothing.

Wash thoroughly with soap and water after handling.

ENVIRONMENTAL PRECAUTIONS

Toxic to birds, small wild animals and aquatic organisms.

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites).

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

No specific antidote is available. Treat the patient symptomatically.

STORAGE

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

1. Thoroughly empty the contents of the container before disposal.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

ACCIDENTAL SPILLS

Handle and open container in a manner as to avoid spillage. If the container is leaking or material is spilled for any reason or cause, carefully sweep up spilled bait and wash exposed area. In spill or leak incidents, keep unauthorized people away.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.