

2021-5878

2022-07-21

This label has been updated according to the re-evaluation decision of Imidacloprid RVD2021-05. While users are encouraged to follow this updated label immediately, the previously approved label is valid until 19/05/2023 in accordance with the phase out period set out in the re-evaluation/special review decision of Imidacloprid RVD2021-05. This previously approved label will be provided upon request by emailing hc.pmra.info-arla.sc@canada.ca. In your email please include the product name and Registration number of the label you are requesting.

2021-5878
2022-07-21

Box label

Primidacide Flea & Tick for Dogs Over 25Kg

Flea and Tick Adulticide

Once-A-Month Topical Flea and Tick Treatment

Liquid

FOR USE ONLY ON DOGS OVER 25 KG

DO NOT USE ON CATS OR KITTENS OR ANY OTHER ANIMAL
DOMESTIC

READ THE LABEL AND INSERT BEFORE USING
KEEP OUT OF REACH OF CHILDREN

Cat glyph in red with diagonal line

WARNING POISON AND EYE IRRITANT
TOXIC TO CATS

ACTIVE INGREDIENT: Imidacloprid 8.8%
Permethrin 44%

REG. NO. 33629 P.C.P. ACT

NET CONTENTS: One to six 4 mL Tubes

Fusion Animal Health

71 Main St.

Exeter, ON

NOM 1S3

416-627-9425

2021-5878
2022-07-21

Primidacide Flea & Tick for Dogs Over 25Kg Flea and Tick Adulticide will control fleas and ticks on dogs within one day of treatment and will continue to protect the dog from further flea and tick infestation for at least four weeks. Re-infestation from the emergence of new fleas and ticks in the environment may continue to occur for six weeks or longer after treatment is initiated. Re-treatment at monthly intervals is required if re-infestation is continuous or if flea and tick control is desired. Shampooing of the dog may shorten the duration of protection. For use on pregnant/lactating animals, consult your veterinarian.

PRECAUTIONS:

HUMANS: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed, or absorbed through the skin. Causes eye irritation. **DO NOT** get in eyes. Avoid contact with skin and clothing. Avoid contact with pets until dry. Wear rubber gloves when applying product. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. This product is extremely toxic to fish. Do not contaminate water, feed or food by storage or disposal.

PETS: Toxic to cats. Cats that actively groom or engage in close physical contact with recently treated dogs may be at risk of serious harm, including death. Sensitivities may occur after using any pesticide for pets. Smaller animals are more likely to experience an adverse reaction. Observe carefully. Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment. Do not use on animals under 8 weeks of age.

Side Effects

Monitor your dog after application. Side effects may include: skin irritation such as redness or scratching; changes in behaviour such as agitation or lethargy; or gastrointestinal effects such as vomiting or loss of appetite. If these or other side effects occur consult your veterinarian or Registrant at 416-627-9425 .

FIRST AID:

HUMANS: If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

DISPOSAL: Wrap empty tube and cap in newspaper and dispose in household garbage.

2021-5878
2022-07-21

Label insert

Primidacide Flea & Tick for Dogs Over 25Kg
Flea and Tick Adulticide
Once-A-Month Topical Flea and Tick Treatment
Liquid

FOR USE ONLY ON DOGS OVER 25 KG
DO NOT USE ON CATS OR KITTENS OR ANY OTHER ANIMAL
DOMESTIC

READ THE LABEL AND INSERT BEFORE USING
KEEP OUT OF REACH OF CHILDREN
cat glyph in red
WARNING POISON AND EYE IRRITANT
TOXIC TO CATS

ACTIVE INGREDIENT: Imidacloprid 8.8%
Permethrin 44%
REG. NO. 33629 P.C.P. ACT

NET CONTENTS: One to six 4 mL Tubes

Fusion Animal Health
71 Main St.
Exeter, ON
N0M 1S3
416-627-9425

2021-5878
2022-07-21

Primidacide Flea & Tick for Dogs Over 25Kg Flea and Tick Adulticide will control fleas and ticks on dogs within one day of treatment and will continue to protect the dog from further flea and tick infestation for at least four weeks. Re-infestation from the emergence of new fleas and ticks in the environment may continue to occur for six weeks or longer after treatment is initiated. Re-treatment at monthly intervals is required if re-infestation is continuous or if flea and tick control is desired. For effective flea and tick control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets, and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters, and surrounding areas. If pest problems persist, an insecticidal premise treatment may be required. Shampooing of the dog may shorten the duration of protection.. This product contains imidacloprid and permethrin. Do not apply another pest control product such as a shampoo, collar, or powder that contains imidacloprid or permethrin to the treated animal after applying Primidacide Flea & Tick For Dogs 25kg And Over.

The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with Primidacide Flea & Tick for Dogs Over 25Kg Flea and Tick Adulticide rapidly kills fleas and may reduce incidence of this condition. For use on pregnant/lactating animals, consult your veterinarian.

PRECAUTIONS:

HUMANS: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed, or absorbed through the skin. Causes eye irritation. DO NOT get in eyes. Avoid contact with skin and clothing. Avoid contact with pets until dry. Wear rubber gloves when applying product. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. This product is extremely toxic to fish. Do not contaminate water, feed or food by storage or disposal.

PETS: Toxic to cats. Cats that actively groom or engage in close physical contact with recently treated dogs may be at risk of serious harm, including death. Sensitivities may occur after using any pesticide for pets. Smaller animals are more likely to experience an adverse reaction.

Observe carefully. Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment.

Do not use on animals under 8 weeks of age.

Side Effects

Monitor your dog after application. Side effects may include: skin irritation such as redness or scratching; changes in behaviour such as agitation or lethargy; or gastrointestinal effects such as vomiting or loss of appetite. If these or other side effects occur consult your veterinarian or Registrant at 416-627-9425 .

FIRST AID:

HUMANS: If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

2021-5878
2022-07-21

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

DISPOSAL: Wrap empty tube and cap in newspaper and dispose in household garbage.

DIRECTIONS FOR USE: Do NOT administer by mouth. Do not use on puppies under 8 weeks of age.

A veterinarian should be consulted when pest infestation continues to be a problem on the pet. Do not use on cats or kittens. Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals under drug or other pesticide treatment. Use only on dogs weighing over 25 kg. This product is packaged in single dosage tubes. This package contains one to six monthly 4.0 mL dosages. Treat your dog monthly in the manner described below. Do not apply a dosage greater than one 4.0 mL tube at one time. Re-treatment at monthly intervals is required if re-infestation is continuous or if flea and tick control is desired.

Remove one applicator from the package.

Hold applicator tube in upright position.

Twist tip to break seal.

The dog should be standing for easy application. The entire contents of the Primidacide Flea and Tick for Dogs Over 25 kg tube should be applied evenly to six or eight spots on the top of the back from the shoulder to the base of the tail. At each spot, part the hair until the skin is visible. Place the tip of the tube on the skin and gently squeeze to dispense a portion of the solution on the skin. Do not apply an excessive amount of solution at any one spot that could cause some of the solution to run off the side of the dog. Do not get this product in your pet's eyes or mouth.

Discard the empty tube as described in "DISPOSAL".

Fusion Animal Health
71 Main St.
Exeter, ON
N0M 1S3
416-627-9425

2021-5878

2022-07-21

Tube label

Primidacide Flea & Tick for Dogs Over 25Kg
READ THE LABEL AND INSERT BEFORE USING
KEEP OUT OF REACH OF CHILDREN
WARNING POISON AND EYE IRRITANT
TOXIC TO CATS
REG. NO. 33629 P.C.P. ACT