Doktor Doom For Your Dog Only Residual & Contact Tick & Flea Killer

For use only on dogs

Kills fleas and ticks on dogs. Do not use on puppies less than 8 weeks old. Do not use on cats, kittens or any other animal

DOMESTIC

SOLUTION

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

ACTIVE INGREDIENT: Permethrin 0.25%

Contains 1,2-benzisothiazolin-3-one at 0.019% as a preservative.

REG. No. 34214 P.C.P.A.

NET CONTENT: 150 mL to 1 L

753146 AB Ltd o/a Ultrasol Industries 10755-69 Ave. Edmonton, Alberta T6H 2C9 780-432-6535

DIRECTIONS FOR USE:

FOR USE ONLY ON DOGS OVER 8 WEEKS OLD: DO NOT use on dogs under 8 weeks of age or nursing animals. Consult a veterinarian before using on sick, aged, pregnant animals or animals receiving drug or other pesticide treatment. DO NOT Use on Cats. Do NOT administer by mouth. Use ONLY in well-ventilated areas, preferably outdoors. This product contains permethrin. DO NOT apply another pest control product such as a spot on, collar or powder which contains permethrin to the treated animal when using this product. Avoid contact with treated animals until dried. This product is ready to use. No further dilution is required. Its special formula is non-irritating to dogs and has no conspicuous odor. Water, not petroleum distillates, is used as the diluent.

SHAKE WELL BEFORE USING. Apply this product with the attached trigger sprayer.

HOW TO PRIME AND CARE FOR YOUR TRIGGER SPRAYER

- 1. After shaking product well, remove cap from container and replace with sprayer hose and cap assembly. Tighten securely.
- 2. Turn sprayer nozzle one-half turn counter-clockwise. Hold sprayer below the level of container and prime by pumping trigger 10-15 times until product is dispensed. Adjust nozzle to fine spray or jet stream as required.
- 3. After each use, thoroughly flush sprayer and hose with water. Return hose and cap assembly to container, tighten securely. Turn nozzle to "OFF" position and store above level of liquid in container to prevent leakage.

NOTE: If sprayer does not function properly, remove nozzle cover and check nozzle hole and holes in rubber component for obstructions. Check to make sure nozzle is not in "OFF" position. Replace nozzle cover, return sprayer assembly to container and proceed.

To control fleas and ticks thoroughly cover the entire animal with a light mist. DO NOT wet fur or skin. Spray ticks directly. Only spray until the fur is damp - not wet. Apply 10 mL for each kg of weight of animal. Hold container in upright position and spray the dog. Spray the entire body of the dog, beginning at the tail and working towards the head. Wearing rubber gloves, fluff the fur of the animal to distribute this product evenly over the fur. DO NOT apply/spray to animal's face, eyes, nose, mouth or genitals. Spray ticks directly. Pay special attention to neck, hindquarters, areas under legs, around the tail and between toes. Do not use this spray more often than once every two weeks. Treatment must only be repeated if pest problem reoccurs. A veterinarian should be consulted when pest infestation continues to be a problem on the pet.

For effective flea and tick control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets, and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters, and surrounding areas. Then use this product on sleeping quarters, cleaned bedding and area around where animal sleeps most often. If pest problems persist, a premise treatment with an insecticide labelled for that use may be required.

PRECAUTIONS:

HUMANS: KEEP OUT OF REACH OF CHILDREN. Do not use as a space spray. Causes skin irritation. Avoid contact with skin. Avoid skin contact by wearing rubber gloves. After handling

or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Toxic to fish and other aquatic life. Cover fish aquaria in the vicinity of treatment area. Toxic to birds. Avoid contamination of food, feed, cooking utensils or drinking water. PETS: Discontinue use if adverse effects are observed.

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

HUMANS: If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15- 20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

Call a poison control centre or doctor for further treatment advice.

PETS: Sensitivities may occur after using ANY pesticide for pets. If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

TOXICOLOGICAL INFORMATION:

Treat symptomatically

DISPOSAL:

Do not re-use empty container. Dispose of empty container in household garbage.

OPTIONAL MARKETING STATEMENTS Ready-to-Use Water-based Insecticide Non-Staining on water safe surfaces No sticky residue Contact & residual action Contains cedar scent