

2022-12-19
2022-2947

[Container Label]

GROUP

2

HERBICIDE

ADVANTAGE FLORASULAM 50 SC

Suspension

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INLCUDING THE PEACE RIVER REGION) ONLY

ADVANTAGE FLORASULAM 50 SC is a selective herbicide for postemergent control or suppression of annual broadleaf weeds including cleavers, wild buckwheat and chickweed in spring wheat (including durum), spring barley, and oats (tank mixes only).

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: florasulam 50 g/L
Contains 1,2-benzisothiazolin-3-one at 0.02% as a preservative

REGISTRATION NO. **34336**
PEST CONTROL PRODUCTS ACT

READ THE LABEL AND THIS ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

**IN CASE OF A MEDICAL EMERGENCY,
CALL TOLL FREE, DAY OR NIGHT:
1-888-CAN-UTEC (226-8832) or 613-996-6666**

Advantage Crop Protection Inc.
620 4th Ave E
Regina, Saskatchewan
S4N 4Z5 Canada
1-888-931-2530

NET CONTENTS: 0.1-1000 L

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

DO NOT APPLY USING AERIAL APPLICATION EQUIPMENT

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.

At completion of spraying or end of the day: Take a shower immediately. Wash thoroughly with soap and water before eating, drinking or smoking. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient. Treat symptomatically.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

- TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.
- This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE

Store this product away from food or feed. Store in original containers in a secure, dry heated storage. ADVANTAGE FLORASULAM 50 SC will freeze at -10°C. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. Do not contaminate food, feedstuffs or domestic water supplies. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Disposal of unused, unwanted product:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in any way that is inconsistent with the directions on the label.

[Booklet Label]

GROUP

2

HERBICIDE

ADVANTAGE FLORASULAM 50 SC

Suspension

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING PEACE RIVER REGION) ONLY

ADVANTAGE FLORASULAM 50 SC is a selective herbicide for postemergent control or suppression of annual broadleaf weeds including cleavers, wild buckwheat and chickweed in spring wheat (including durum), spring barley, and oats (tank mixes only).

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: florasulam 50 g/L
Contains 1,2-benzisothiazolin-3-one at 0.02% as a preservative

REGISTRATION NO. **34336**
PEST CONTROL PRODUCTS ACT

READ THE LABEL AND THIS BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

**IN CASE OF A MEDICAL EMERGENCY,
CALL TOLL FREE, DAY OR NIGHT:
1-888-CAN-UTEC (226-8832) or 613-996-6666**

Advantage Crop Protection Inc.
620 4th Ave E
Regina, Saskatchewan
S4N 4Z5 Canada
1-888-931-2530

NET CONTENTS: 0.1-1000 L

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

DO NOT APPLY USING AERIAL APPLICATION EQUIPMENT

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.

At completion of spraying or end of the day: Take a shower immediately. Wash thoroughly with soap and water before eating, drinking or smoking. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient. Treat symptomatically.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

- TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.
- This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE

Store this product away from food or feed. Store in original containers in a secure, dry heated storage. ADVANTAGE FLORASULAM 50 SC will freeze at -10°C. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. Do not contaminate food, feedstuffs or domestic water supplies. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Disposal of unused, unwanted product:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

GENERAL INFORMATION

ADVANTAGE FLORASULAM 50 SC is a selective postemergence herbicide for the control of hard-to-kill annual broadleaf weeds such as chickweed, wild buckwheat and cleavers in spring wheat (including durum), spring barley, and oats (tank mixes only) not underseeded with legumes. ADVANTAGE FLORASULAM 50 SC is mixed with water and applied as a uniform broadcast spray. It is non-corrosive, nonflammable, and nonvolatile.

ADVANTAGE FLORASULAM 50 SC must be applied early postemergence, to the main flush of actively growing broadleaf weeds. Warm, moist growing conditions promote active weed growth and enhance the activity of ADVANTAGE FLORASULAM 50 SC by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds. See "DIRECTIONS FOR USE" section of this label for complete use details.

ADVANTAGE FLORASULAM 50 SC stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 2 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

MODE OF ACTION

ADVANTAGE FLORASULAM 50 SC inhibits the production of the ALS enzyme in plants. This enzyme is essential for the production of certain amino acids which are essential for plant growth. ADVANTAGE FLORASULAM 50 SC is a Group 2 mode of action herbicide.

GENERAL USE PRECAUTIONS

Sensitive Plants

Do not apply ADVANTAGE FLORASULAM 50 SC directly to, or otherwise permit it to come in direct contact with susceptible crops or desirable plants including alfalfa, edible beans, canola, flowers and ornamentals, lentils, lettuce, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes or tobacco.

Non-Target Sites

Do not apply where proximity of susceptible crops (e.g. canola and legumes) or other desirable plants is likely to result in exposure to spray or spray drift. See Environmental Precautions section of the label.

Crop Rotation

Fields previously treated with ADVANTAGE FLORASULAM 50 SC can be seeded the following year to barley, canola, peas, wheat or fields can be summerfallowed.

Do not use in successive years at the same site.

This product has potential to leach. Do not apply excessive irrigation.

Tank Mixtures

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Advantage Crop Protection Inc. representatives at 1-888-931-2530 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

Spray Equipment Precaution

Do not apply through any type of irrigation system.

DO NOT APPLY USING AERIAL APPLICATION EQUIPMENT.

To Reduce Spray Drift:

1. Use nozzles delivering higher volumes and coarser droplets.
2. Use low pressures (200 to 275 kPa).
3. Use 100 L/ha of spray solution.
4. Spray when the wind velocity is 15 km/hr or less.
5. Spot treatments should only be applied with a calibrated boom to prevent over-application.

Sprayer Clean-Out Instructions

To avoid injury to desirable plants, thoroughly clean equipment used to apply this product before re-use or using it to apply other chemicals.

1. Immediately after spraying, completely drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.

2. First rinse:

- Spray the inside of tank with clean water and fill the sprayer with at least one tenth of the spray tank volume.
- Agitate and circulate for 15 minutes, and flush through booms and hoses.
- Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
- Drain tank completely.

3. Second rinse:

- Fill the tank with clean water.
- Add All Clear Spray Tank Decontaminator, or Clean-Out Spray Tank Cleaner, or 1 L of household ammonia (containing a minimum of 3 % ammonia) per 100 L of water, or similar tank cleaning agent as per manufacturer's recommendations while filling the tank with clean water.
- Agitate and then flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full. Allow to stand for 15 minutes with agitation. Flush the solution out of the spray tank through the spray booms. Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
- If possible, let the solution stand in the sprayer tank and booms for an extended period of time, overnight if possible.
- After flushing the boom and hoses, drain tank completely.
- Remove nozzles and screens and clean separately with a cleaning agent or an ammonia solution (100 mL in 10 L water).

4. Third rinse:

- Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume.
- Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
- Drain tank completely.

Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty odour which may cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application: **DO NOT** apply using aerial application equipment.

SPRAY BUFFER ZONES

Spot treatments using hand-held equipment do not require a spray buffer zone.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Crop	Method of application	Spray Buffer Zones (metres) Required for the Protection of:		
		Freshwater Habitat of Depths:		Terrestrial Habitat:
		Less than 1 m	Greater than 1 m	
All labelled crops	Field sprayer (medium)	1	0	3

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product that are larger than 1m can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE.

DO NOT APPLY TO CROPS UNDERSEEDED WITH LEGUMES.

ADVANTAGE FLORASULAM 50 SC ALONE

Crops Registered

Spring wheat (including durum) and spring barley

Application Directions

Apply 100 mL of ADVANTAGE FLORASULAM 50 SC per hectare in a minimum of 100 L per hectare of water. Add Agral 90 at 0.2% v/v. See weeds species controlled under “Weeds Controlled or Suppressed by ADVANTAGE FLORASULAM 50 SC Alone at 100 mL/ha.”

Apply to actively growing wheat and barley from the 2 to 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds.

Application Timing

Apply to actively growing weeds in the 2-4 leaf stage. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Only weeds which are emerged at the time of application will be affected. If foliage is wet at the time of application control may be decreased. Under conditions of low crop and high weed density, control may be reduced.

Weeds Controlled or Suppressed by ADVANTAGE FLORASULAM 50 SC Alone at 100mL/ha

Weeds Controlled:

buckwheat, wild	shepherd's-purse
canola, volunteer♦	smartweed
chickweed, common	stinkweed
cleavers	mustard, wild

♦Including herbicide-tolerant canola varieties except Clearfield canola.

Weeds Suppressed:

hempnettle	sowthistle, annual
pigweed, redroot	sowthistle, perennial♦

♦Applications made at advanced leaf stages will reduce product effectiveness.

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of ADVANTAGE FLORASULAM 50 SC
4. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
5. Add Agral 90 as the last ingredient at the rate of 0.2% v/v (2 L per 1000 L of spray volume).
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
7. Follow sprayer clean-up directions.

Preharvest/Grazing Intervals

Livestock may be grazed on treated crops 7 days following application. Do not harvest the treated crop within 60 days after application.

TANK-MIX COMBINATIONS:

ADVANTAGE FLORASULAM 50 SC + Foax HERBICIDE

For control of annual grasses (see table below) tank mix ADVANTAGE FLORASULAM 50 SC with Foax Herbicide. Refer to the above table for broadleaf weeds controlled or suppressed with ADVANTAGE FLORASULAM 50 SC. Read all tank mix partner labels thoroughly for more information on precautions, preharvest intervals, buffer zone requirements and additional directions for use.

Tank-Mix Combinations with ADVANTAGE FLORASULAM 50 SC for Annual Grass Control

Tank-Mix Partner	Crops Registered	Rate/ha	Adjuvant and Rate	Additional Weeds Controlled
FOAX Herbicide	spring wheat durum wheat	230 mL	0.8% v/v*	wild oats, green foxtail, yellow foxtail, volunteer (tame) oats, volunteer canary seed, barnyard grass
		290 mL	1.0% v/v*	wild oats, green foxtail, yellow foxtail, volunteer (tame) oats, volunteer canary seed, barnyard grass, plus: Persian dandel

*Always use an adjuvant of either SCORE, ASSIST, or CROPOIL 83/17 ADJUVANT with FOAX HERBICIDE.

To control additional broadleaf weeds, MCPA Ester may be added to either tank mix up to a maximum of 420 g ae/ha (equivalent to 840 mL/ha of MCPA LV500). Refer to the MCPA Ester tank mix partner label for rates less than 420 g ae/ha and weed controlled.

Mixing Instructions

1. Begin to fill sprayer tank with clean water, and engage agitator. **Agitation must be continued throughout the entire mixing and spraying procedure.**
2. When the sprayer is three quarters full of water, add ADVANTAGE FLORASULAM 50 SC and agitate for 2-3 minutes.
3. If 2,4-D Ester or MCPA Ester are also being added, add these next.
4. Add Foax Herbicide. Agitate for 2-3 minutes.
5. Add required Adjuvant.
6. Agitate for 1-2 minutes before adding remainder of water and then maintain constant agitation.
7. After any break in spraying operations, agitate thoroughly before spraying again.
8. **Use the spray suspension as soon as it is prepared.**
9. If an oil film starts to build up in the tank, drain tank and then clean as directed on this label.

TANK MIX COMBINATIONS:

ADVANTAGE FLORASULAM 50 SC + MCPA LV 500 HERBICIDE

Crops Registered

Spring wheat (including durum), spring barley, and oats

Application Directions

For control of a wide spectrum of broadleaf weeds apply ADVANTAGE FLORASULAM 50 SC at 100 mL/ha tank mixed with MCPA LV 500 Herbicide at 0.70 L/ha (350 g ae/ha) in 100 L per hectare of water. Apply to actively growing wheat, barley, and oats from the 2 to 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds. Follow all precautions, directions for use, and limitations on the tank mix partner labels.

Weeds Controlled or Suppressed by the Tank Mixture of ADVANTAGE FLORASULAM 50 SC + MCPA LV 500 Herbicide

Weeds Controlled:

buckwheat, wild		ragweed, common
burdock◆◆	lamb'squarters	shepherd's purse
canola, volunteer◆	mustard, ball	smartweed
chickweed, common	mustard, wild	stinkweed
cleavers	pigweed, redroot◆◆◆	sunflower, annual◆◆
flixweed (spring rosettes only)	pigweed, Russian◆◆	
hempenettle◆◆◆	prickly lettuce◆◆	

Weeds Suppressed:

dandelion (seedlings and overwintered rosettes less than 15 cm) ◆◆◆		sowthistle, perennial
plantain (top growth)		stork's bill◆◆◆
sowthistle, annual		thistle, Canada◆◆◆

◆Including herbicide-tolerant canola varieties

◆◆Up to 4 leaf stage

◆◆◆Optimum control can be achieved with the addition of 140 mL (70 g ae/ha) of MCPA LV 500 Herbicide

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of ADVANTAGE FLORASULAM 50 SC
4. Add the correct amount of MCPA LV 500 Herbicide and continue to agitate.
5. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
7. Follow sprayer clean-up directions.

Note: Do not add a surfactant to this tank mixture.

Pre-Harvest/Grazing Intervals (ADVANTAGE FLORASULAM 50 SC + MCPA LV 500 Herbicide)

1. Do not cut the treated crop for hay or graze treated crop within 7 days after application.
2. Do not harvest the treated crop within 60 days after application.

TANK-MIX COMBINATION:

ADVANTAGE FLORASULAM 50 SC + MCPA LV 500 HERBICIDE + OTHER HERBICIDES

Crops Registered

Spring wheat (including durum) and spring barley

Do not apply these tank mixes to oats.

Application Directions

Tank mixtures of ADVANTAGE FLORASULAM 50 SC + MCPA LV 500 Herbicide with other herbicides will provide control of broadleaf weeds and specified annual grasses. Apply to actively growing wheat and barley from the 2 to 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds. Follow all precautions, minimum interval to harvest and directions for use on the ADVANTAGE FLORASULAM 50 SC and tank mix partner labels.

Tank-Mix Combinations with ADVANTAGE FLORASULAM 50 SC + MCPA LV 500 Herbicide + Other Herbicides

Herbicide Tank-Mix Partner	Crops Registered	Rate (Product/ha)	Adjuvant and Rate	Additional Weeds Controlled
Assert² 300 SC Herbicide	Wheat (spring & durum), Spring barley	1.6 L	Refer to Assert 300 SC Herbicide label	wild oat
Foax Herbicide	Wheat (spring & durum)	230 mL	0.8% v/v*	wild oats, green foxtail, yellow foxtail, volunteer (tame) oats, volunteer canary seed, barnyard grass
		290 mL	1.0% v/v*	wild oats, green foxtail, yellow foxtail, volunteer (tame) oats, volunteer canary seed, barnyard grass, plus: Persian darnel
Everest³ Solupak 70DF ♦ Herbicide	Wheat (spring & durum)	43 g	Ag-Surf or Agral 90 0.25% v/v	Wild oat and green foxtail

*Always use an adjuvant of either SCORE, ASSIST, or CROPOIL 83/17 ADJUVANT with FOAX HERBICIDE.

♦Wheat (including durum) exposed to water-logged or saturated soils, or temperature extremes such as heat or freezing weather, or drought, low fertility or plant disease at application time could show unacceptable injury symptoms. Weed control may also be reduced by these same conditions.

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add ADVANTAGE FLORASULAM 50 SC (see Notes 2 and 3 below).
4. Add MCPA LV 500 Herbicide and continue to agitate.
5. Add the required amount of tank mix partner and continue to agitate.
6. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
7. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
8. Follow sprayer clean-up directions.

Note 1: Add only the adjuvant recommended above. Follow tank-mix partner label for order of mixing and any specific instructions for mixing.

Note 2: When using Assert 300 SC Herbicide, add tank mix partners in the following order: ADVANTAGE FLORASULAM 50 SC, Spray Water pH Adjuster, Assert 300 SC Herbicide, MCPA LV 500 Herbicide.

Note 3: When using Everest Solupak 70DF Herbicide add tank mix partners in the following order: Everest Solupak 70DF Herbicide, tank mix partner, adjuvant.

**TANK-MIX COMBINATIONS:
ADVANTAGE FLORASULAM 50 SC + 2,4-D LV 600 EC HERBICIDE**

Crops Registered

Spring wheat (including durum)

Application Directions

For control of a wide spectrum of broadleaf weeds apply ADVANTAGE FLORASULAM 50 SC at 100 mL/ha tank-mixed with 2,4-D LV 600 Herbicide at 1.0 L/ha. Follow all precautions, directions for use and limitations on the tank -mix partner labels. Apply when wheat is in the 3 leaf expanded stage to the 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be affected. Best results are obtained from application made to seedling weeds.

Weeds Controlled or Suppressed by the Tank Mixture of ADVANTAGE FLORASULAM 50 SC + 2,4-D LV 600 Herbicide

Weeds Controlled:

bluebur	flixweed	ragweed, common
burdock	kochia♦♦	shepherd'spurse
buckwheat, tartary	lady's-thumb	smartweed
buckwheat, wild	lamb'squarters	sowthistle, annual
canola, volunteer♦	mustard, ball	stinkweed
chickweed, common	mustard, wild	sunflower, annual
cleavers	pigweed, redroot	thistle, Russian
cocklebur	plantain	vetch
dandelion (seedlings and overwintered rosettes)	prickly lettuce	wild radish

♦Including all herbicide-tolerant canola varieties.

♦♦Up to 5 cm in height

Weeds Suppressed: hempnettle

sowthistle, perennial (top growth control) thistle, Canada (top growth control)

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add ADVANTAGE FLORASULAM 50 SC
4. Add 2,4-D LV 600 Herbicide and continue to agitate.
5. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
7. Follow sprayer clean-up directions.

Pre-Harvest/Grazing Intervals (ADVANTAGE FLORASULAM 50 SC + 2,4-D LV 600 Herbicide)

1. Do not permit lactating dairy animals to graze fields within 7 days after application.
2. Do not harvest forage or cut hay within 30 days after application.
3. Withdraw meat animals from treated fields at least 3 days before slaughter.
4. Do not harvest the treated mature crop within 60 days after application.

TANK-MIX COMBINATIONS:

ADVANTAGE FLORASULAM 50 SC + 2,4-D LV 600 HERBICIDE + OTHER HERBICIDES

Crops Registered

Spring wheat (including durum)

Tank mixtures of ADVANTAGE FLORASULAM 50 SC + 2,4-D LV600 Herbicide with other herbicides will provide control of broadleaf weeds and specified annual grasses. Apply when crops and weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds. Follow all precautions, minimum interval to harvest and directions for use on the ADVANTAGE FLORASULAM 50 SC and tank-mix partner labels.

Tank-Mix Combinations with ADVANTAGE FLORASULAM 50 SC + 2,4-D LV 600 Herbicide + Other Herbicides

Herbicide Tank-Mix Partner	Rate (Product/ha)	Adjuvant ¹ and Rate	Additional Weeds Controlled
Everest ³ Solupak 70DF ♦ Herbicide	43 g	Ag-Surf or Agral 90 0.25% v/v	wild oat and green foxtail

♦Wheat (including durum) exposed to water-logged or saturated soils, or temperature extremes such as heat or freezing weather, or drought, low fertility or plant disease at application time could show unacceptable injury symptoms. Weed control may also be reduced by these same conditions.

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add ADVANTAGE FLORASULAM 50 SC (see Note 2 below).
4. Add 2,4-D LV 600 Herbicide and continue to agitate.
5. Add tank mix partner and continue to agitate.
6. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
7. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
8. Follow sprayer clean-up directions.

Note 1: Add only the adjuvant recommended below. Follow tank-mix partner label for order of mixing and any specific instructions for mixing.

Note 2: When using Everest Solupak 70DF Herbicide add tank mix partners in the following order: Everest Solupak 70DF Herbicide, tank mix partner, adjuvant. Vigorous agitation is required to dissolve Everest Solupak 70DF Herbicide before adding other tank mix partners.

TANK MIX COMBINATIONS:

ADVANTAGE FLORASULAM 50 SC + GLYPHOSATE PRESENT AS ISOPROPYLAMINE SALT OR DIMETHYLAMINE SALT

Application Directions

ADVANTAGE FLORASULAM 50 SC may be tank mixed with glyphosate present as isopropylamine salt (e.g. Glyphos Soluble Concentrate Herbicide) or dimethylamine salt to control volunteer glyphosate-tolerant canola, wild buckwheat (up to 5 leaf) and top growth control of dandelion (up to 6 leaf stage), in addition to the labelled weeds controlled by ADVANTAGE FLORASULAM 50 SC or glyphosate.

Apply ADVANTAGE FLORASULAM 50 SC at 100 mL/ha with glyphosate herbicide (present as isopropylamine salt or dimethylamine salt) at 450 g a.e./ha in 100 L water per hectare.

Application Timing

Apply to actively growing weeds in the 2-4 leaf stage, except where noted above. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may reduce weed control. Only weeds which are emerged at the time of application will be affected. If foliage is wet at the time of application, control may be decreased. Under conditions of high weed density, control may be reduced.

Apply in the spring prior to planting barley, oats and wheat, or as an initial summerfallow treatment, or in the fall prior to planting barley, oats and wheat the following spring. For annual weed control in the fall, only the weeds at labelled growth stage will be controlled. Heavy frost prior to application in the fall may decrease control.

Do not use in successive years at the same site.

Apply only one of the following treatments to the same site in the same year: a tank-mix of ADVANTAGE FLORASULAM 50 SC and glyphosate (present as isopropylamine salt or dimethylamine salt, prior to planting or as the initial treatment in summerfallow).

RESISTANCE MANAGEMENT RECOMMENDATION

For resistance management, ADVANTAGE FLORASULAM 50 SC is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to ADVANTAGE FLORASULAM 50 SC and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of ADVANTAGE FLORASULAM 50 SC or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific

crops and weed biotypes.

- For further information or to report suspected resistance, contact Advantage Crop Protection Inc. representatives at 1-888-931-2530 or at www.advantagecropprotection.ag.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in any way that is inconsistent with the directions on the label.

ADVANTAGE FLORASULAM 50 SC is a trademark of Advantage Crop Protection Inc.

1 Assert is a trademark of Nufarm Agriculture Inc.

2 FOAX Herbicide is a trademark of Farmer's Business Network Canada Inc.

3 Everest is a trademark of Bayer Corp. used under licence by Bayer Inc.

4 Solupak is a registered trademark of Bayer Inc.