

## A22070 Fungicide

COMMERCIAL

SUSPENSION

Broad-spectrum fungicide for control of turf diseases.

### ACTIVE INGREDIENTS:

Pydiflumetofen .....	10.2 g/L
Azoxystrobin .....	62.5 g/L
Propiconazole .....	104 g/L

Contains 1,2-benzisothiazolin-3-one at 0.029% , 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00122% and 2-methyl-4-isothiazolin-3-one at 0.000407% as preservatives.

**READ THE LABEL AND PAMPHLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**



**POISON**

**WARNING: EYE IRRITANT AND SKIN IRRITANT**

REGISTRATION NUMBER: **33798**  
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **0.5 L to Bulk**

**Syngenta Canada Inc.**  
140 Research Lane, Research Park  
Guelph, Ontario N1G 4Z3  
Telephone: 1-877-964-3682

Label

## 1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

## 2.0 FIRST AID

**IN CASE OF POISONING**, contact a doctor or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**If swallowed**, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing**, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes**, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

## 3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote. Treat symptomatically.

## 4.0 PRECAUTIONS

### **KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS.**

Harmful if swallowed or inhaled. Causes eye and skin irritation. DO NOT get in eyes or on skin. Avoid breathing spray mist. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Prolonged or frequent, repeated skin contact may cause allergic reactions in some individuals.

For sod farms, **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. For golf courses, **DO NOT** enter or allow entry into treated areas until sprays have dried.

Avoid inhalation of vapours/dust and spray mist, and working in spray mist.

Apply only when the potential for drift to areas of human habitation and human activity (other than

golf courses) such as parks, school grounds, and playing fields, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Read and follow all tank mix partner labels and observe the most stringent label restrictions, precautions and instructions for each tank mixture.

Not for use in residential areas. Residential areas are defined as any use site where bystanders including children could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public including children could be exposed.

**DO NOT apply using aerial application equipment.**

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

## **5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing and loading. Gloves are not required during application within a closed cab.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash splashes from skin IMMEDIATELY with plenty of water. Remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. As soon as possible, wash thoroughly and change into clean clothing. After spraying, wash hands and shower thoroughly with soap and water.

Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Wear freshly laundered clothes daily.

Do not wear contaminated shoes.

Users should wash hands and exposed skin before eating, drinking, chewing gum, using tobacco, applying cosmetics or using the toilet.

While using product, do not eat, drink or use tobacco (including smoking).

## **6.0 ENVIRONMENTAL PRECAUTIONS**

This product is TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones and precautionary measures specified under DIRECTIONS FOR USE.

This product demonstrates the properties and characteristics associated with chemicals detected

in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to, heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted, fine textured or low in organic matter such as clay). Avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body. For the protection of aquatic plants, overspray or drift to important wildlife habitats such as wetlands, sloughs or dry slough borders and water bodies should be avoided. Drift to emerging vegetation or newly emerged vegetation in shelterbelts or woodlots should be avoided.

The active ingredient propiconazole is toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland.

Pydiflumetofen and azoxystrobin are persistent and may carryover. It is recommended that this product not be used in areas treated with any products containing either pydiflumetofen or azoxystrobin or both during the previous season.

**DO NOT** apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands), estuarine/marine habitats.

**DO NOT** discharge effluent containing this product into sewer systems, lakes, streams, ponds, estuaries, oceans and other waters.

## 7.0 STORAGE

Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated secure area away from feed, foodstuffs, beverages and tobacco, and out of the reach of children, unauthorized persons and animals. To prevent contamination store this product away from seed, food, feed or fertilizer.

Repeated freeze-thawing of A22070 Fungicide will not affect the physical integrity of the product. If the product should freeze, allow to thaw and agitate thoroughly prior to use.

Not for use or storage in or around the home.

## 8.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

### **For Recyclable Containers:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsate to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

**For Returnable Containers:**

Do not reuse this container for any purpose. For disposal, the empty container may be returned to the point of purchase (distributor/dealer).

**For Refillable Containers:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,  
CALL 1-800-327-8633 (FASTMED)***

GROUP 7 3 11 FUNGICIDES

## A22070 Fungicide

COMMERCIAL

SUSPENSION

Broad-spectrum fungicide for control of turf diseases.

### ACTIVE INGREDIENTS:

Pydiflumetofen ..... 10.2 g/L

Azoxystrobin ..... 62.5 g/L

Propiconazole ..... 104 g/L

Contains 1,2-benzisothiazolin-3-one at 0.029% , 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00122% and 2-methyl-4-isothiazolin-3-one at 0.000407% as preservatives.

**READ THE LABEL AND PAMPHLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**



POISON

**WARNING: EYE IRRITANT AND SKIN IRRITANT**

REGISTRATION NUMBER: **33798**  
PEST CONTROL PRODUCTS ACT

**Syngenta Canada Inc.**  
140 Research Lane, Research Park  
Guelph, Ontario N1G 4Z3  
Telephone: 1-877-964-3682

Pamphlet

---

## Table of Contents

	<u>Section Number</u>
NOTICE TO USER .....	1.0
FIRST AID .....	2.0
TOXICOLOGICAL INFORMATION .....	3.0
PRECAUTIONS.....	4.0
PERSONAL PROTECTIVE EQUIPMENT (PPE) .....	5.0
ENVIRONMENTAL PRECAUTIONS .....	6.0
STORAGE .....	7.0
DISPOSAL .....	8.0
PRODUCT INFORMATION .....	9.0
DIRECTIONS FOR USE .....	10.0
General Information .....	10.1
Ground Application .....	10.2
General Precautions .....	10.2.1
Mixing Instructions .....	10.2.2
Spraying Instructions .....	10.2.3
Equipment Clean-Up.....	10.2.4
Spray Buffer Zones .....	10.3
Spray Drift Management and Application Requirements .....	10.4
CROP USE DIRECTIONS .....	11.0
MINOR USES .....	12.0
RESISTANCE MANAGEMENT .....	13.0

## 1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

## 2.0 FIRST AID

**IN CASE OF POISONING**, contact a doctor or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**If swallowed**, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing**, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes**, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

## 3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote. Treat symptomatically.

## 4.0 PRECAUTIONS

### **KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS.**

Harmful if swallowed or inhaled. Causes eye and skin irritation. DO NOT get in eyes or on skin. Avoid breathing spray mist. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Prolonged or frequent, repeated skin contact may cause allergic reactions in some individuals.

For sod farms, **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. For golf courses, **DO NOT** enter or allow entry into treated areas until sprays have dried.

Avoid inhalation of vapours/dust and spray mist, and working in spray mist.

Apply only when the potential for drift to areas of human habitation and human activity (other than



golf courses) such as parks, school grounds, and playing fields, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Read and follow all tank mix partner labels and observe the most stringent label restrictions, precautions and instructions for each tank mixture.

Not for use in residential areas. Residential areas are defined as any use site where bystanders including children could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public including children could be exposed.

### **DO NOT apply using aerial application equipment**

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

## **5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing and loading. Gloves are not required during application within a closed cab.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash splashes from skin IMMEDIATELY with plenty of water. Remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. As soon as possible, wash thoroughly and change into clean clothing. After spraying, wash hands and shower thoroughly with soap and water.

Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Wear freshly laundered clothes daily.

Do not wear contaminated shoes.

Users should wash hands and exposed skin before eating, drinking, chewing gum, using tobacco, applying cosmetics or using the toilet.

While using product, do not eat, drink or use tobacco (including smoking).

## **6.0 ENVIRONMENTAL PRECAUTIONS**

This product is TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones and precautionary measures specified under DIRECTIONS FOR USE.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to, heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted, fine textured or low in organic matter such as clay). Avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body. For the protection of aquatic plants, overspray or drift to important wildlife habitats such as wetlands, sloughs or dry slough borders and water bodies should be avoided. Drift to emerging vegetation or newly emerged vegetation in shelterbelts or woodlots should be avoided.

The active ingredient propiconazole is toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland.

Pydiflumetofen and azoxystrobin are persistent and may carryover. It is recommended that this product not be used in areas treated with any products containing either pydiflumetofen or azoxystrobin or both during the previous season.

**DO NOT** apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands), estuarine/marine habitats.

**DO NOT** discharge effluent containing this product into sewer systems, lakes, streams, ponds, estuaries, oceans and other waters.

## 7.0 STORAGE

Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated secure area away from feed, foodstuffs, beverages and tobacco, and out of the reach of children, unauthorized persons and animals. To prevent contamination store this product away from seed, food, feed or fertilizer.

Repeated freeze-thawing of A22070 Fungicide will not affect the physical integrity of the product. If the product should freeze, allow to thaw and agitate thoroughly prior to use.

Not for use or storage in or around the home.

## 8.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

---

**For Recyclable Containers:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsate to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

**For Returnable Containers:**

Do not reuse this container for any purpose. For disposal, the empty container may be returned to the point of purchase (distributor/dealer).

**For Refillable Containers:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,  
CALL 1-800-327-8633 (FASTMED)***

**9.0 PRODUCT INFORMATION**

A22070 Fungicide is a fungicide for control of diseases in turf. A22070 Fungicide is best used preventatively for the control of plant diseases. Read all label directions before use. All applications must be made in accordance with the DIRECTIONS FOR USE.

**10.0 DIRECTIONS FOR USE**

**10.1 General Information**

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

**DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or containers, or disposal of wastes.

**10.2 Ground Application**

**10.2.1 General Precautions**

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural and Biological Engineers (ASABE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply using aerial application equipment.

DO NOT apply through irrigation equipment.

DO NOT apply through any type of ultra-low volume (ULV) spray system (less than 30 litres per hectare)

### 10.2.2 Mixing Instructions

Protect sprayer operators from drift or mist. Additional information on spray drift management for ground application is provided in the section "Spray Drift Management and Application Requirements". When low volumes of spray are applied, complete coverage and thorough application are essential for most effective results. Schedule applications in accordance with local conditions. Consult your local agricultural authorities for specific use information. Where possible, use a nurse tank to fill your spray tank.

#### Mixing Steps:

1. Spray equipment should be thoroughly flushed with clean water before mixing A22070 Fungicide.
2. Fill spray tank 1/2 full with carrier. Engage gentle agitation.
3. Pour in the required amount of add the A22070 Fungicide.
4. Finish filling tank. Maintain gentle agitation during mixing and spraying operations.
5. Do not let stand without agitation.

If tank mixes are desired, observe all directions, precautions, and limitations on tank mix partner labels.

Do not use silicone based products with A22070 Fungicide due to possible phytotoxicity.

To determine the physical compatibility of A22070 Fungicide with other products, follow the WALES mixing order and use a jar test.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Syngenta Canada Inc. at 1-87-SYNGENTA (1-877-964-3682) for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

### 10.2.3 Spraying Instructions

1. Water Volume: Apply in a minimum spray volume of 200 L/ha OR the volume given in the crop and pest specific instructions tabulated below, whichever is LARGER.
2. Spray Nozzles: Use only ASABE medium or coarse nozzles. The use of higher water volumes and larger spray tip sizes will generally increase spray deposition and reduce drift.
3. Pressure: As recommended by the nozzle manufacturer to achieve ASABE medium or coarse sized droplets.

4. Ensure tank agitation is engaged whenever possible.
5. Keep bypass line at bottom of tank to minimize foaming.
6. Use 16 mesh suction screens, 50 mesh line and tip screens, and spray pressure of 200-30 kPa.

### 10.2.4 Equipment Clean-Up

#### Before Spraying:

Prior to using A22070 Fungicide, ensure that the spray tank, chemical handling equipment, lines and filter are thoroughly cleaned.

#### After Spraying:

Thoroughly clean application equipment immediately after spraying. **DO NOT ALLOW** A22070 Fungicide residue to dry within application equipment.

When using tank mixes, consult the tank-mix partner label for additional clean-up instructions. The following recommendations are provided:

1. Drain and flush tank walls, boom and all hoses for ten minutes with a clean water/detergent mixture. Rinse with clean water. **DO NOT** clean application equipment near desirable vegetation, wells or other water sources.
2. Remove all nozzles and screens and wash separately.
3. Dispose of all rinsate in accordance with provincial regulations.

### 10.3 Spray Buffer Zones

Spot treatments using hand-held equipment do not require a spray buffer zone.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop	Spray Buffer Zones (metres) Required for the Protection of:				
		Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths:		Terrestrial Habitat:
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Turf	2	1	2	1	2

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the

---

coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

#### **10.4 Spray Drift Management and Application Requirements**

Use good pesticide practices and apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and parks is minimal. Take into consideration wind speed, wind direction, temperature, application equipment and sprayer settings used for application.

For the protection of non-target habitats, over-spray or drift to any body of water or other environmentally sensitive habitat must be avoided. The interaction of equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

## 11.0 CROP USE DIRECTIONS

<b>TURF (including golf courses and sod farms)</b>	
<b>Diseases Controlled</b>	<b>Dollar spot (<i>Sclerotinia homeocarpa</i>)</b>
<b>Use rate (product/ha)</b>	48 to 76 mL/100 m <sup>2</sup> or 4.8 to 7.6 L/ha
<b>Application Interval</b>	14 to 28 days
<b>Minimum recommended application of water volume</b>	200 to 800 L/ha for ground application
<b>Maximum number of applications per crop</b>	4
<b>Maximum rate of active ingredients applied</b>	0.78 g ai/100 m <sup>2</sup> or 78 g ai/ha of pydiflumetofen 4.75 g ai/100 m <sup>2</sup> or 475 g ai/ha of azoxystrobin 7.90 g ai/100 m <sup>2</sup> or 790 g ai/ha propiconazole
<b>Application Directions</b>	<p>For best control, use as a component of an integrated disease management program.</p> <p>Begin application when conditions are favourable for disease infection and prior to disease symptom expression.</p> <p>Use the higher rate under severe disease pressure.</p> <p>During prolonged periods of favourable conditions apply on the shorter application interval.</p> <p>Apply in sufficient water to ensure thorough coverage. 2-8 litres of water per 100 m<sup>2</sup> (200-800 L/ha) is recommended.</p> <p>Use AI, XR, or Syngenta XC nozzles.</p> <p>Apply after mowing or allow spray to completely dry before mowing.</p> <p>Allow sprayed area to completely dry before irrigation.</p>
<b>Diseases Controlled</b>	<b>Waitea patch (<i>Waitea circinata</i> var. <i>circinata</i>)</b>
<b>Use rate (product/ha)</b>	48 to 95 mL/100 m <sup>2</sup> or 4.8 to 9.5 L/ha
<b>Application Interval</b>	14 to 28 days
<b>Minimum recommended application of water volume</b>	200 to 800 L/ha for ground application
<b>Maximum number of applications per crop</b>	4
<b>Maximum rate of active ingredients applied</b>	1 g ai/100 m <sup>2</sup> or 100 g ai/ha of pydiflumetofen 5.94 g ai/100 m <sup>2</sup> or 594 g ai/ha of azoxystrobin 9.88 g ai/100 m <sup>2</sup> or 988 g ai/ha propiconazole

<b>Application Directions</b>	Apply when conditions are favourable for disease development, i.e., warmer temperatures between 20 to 30°C and during periods of high humidity and extended leaf wetness. Use the higher rate and shorter application interval under moderate to severe disease pressure. During prolonged periods of favourable conditions apply to the shorter application interval.
<b>Diseases Controlled</b>	<b>Leaf spot, melting out</b> ( <i>Bipolaris sorokiniana</i> , <i>Dreschlera poae</i> )
<b>Use rate (product/ha)</b>	48 to 95 mL/100 m <sup>2</sup> or 4.8 to 9.5 L/ha
<b>Application Interval</b>	14 to 28 days
<b>Minimum recommended application of water volume</b>	200 to 800 L/ha
<b>Maximum number of applications per crop</b>	4
<b>Maximum rate of active ingredients applied</b>	1 g ai/100 m <sup>2</sup> or 100 g ai/ha of pydiflumetofen 5.94 g ai/100 m <sup>2</sup> or 594 g ai/ha of azoxystrobin 9.88 g ai/100 m <sup>2</sup> or 988 g ai/ha propiconazole
<b>Application Directions</b>	Apply when conditions are favourable for disease development.  Use the higher rate under severe disease pressure.  During prolonged periods of favourable conditions apply on the shorter application interval
<b>Diseases Controlled</b>	<b>Summer patch</b> ( <i>Magnaporthe poae</i> )
<b>Use rate (product/ha)</b>	48 to 95 mL/100 m <sup>2</sup> or 4.8 to 9.5 L/ha
<b>Application Interval</b>	14 to 28 days
<b>Minimum recommended application of water volume</b>	200 to 800 L/ha for ground application
<b>Maximum number of applications per crop</b>	4
<b>Maximum rate of active ingredients applied</b>	1 g ai/100 m <sup>2</sup> or 100 g ai/ha of pydiflumetofen 5.94 g ai/100 m <sup>2</sup> or 594 g ai/ha of azoxystrobin 9.88 g ai/100 m <sup>2</sup> or 988 g ai/ha propiconazole
<b>Application Directions</b>	Apply on a preventative basis. The first application should be when the soil temperature is greater than 16°C at a 5 cm depth for 4 consecutive days with a second application 30 days later. Additional applications may be required if hot, humid weather occurs.
<b>Diseases Controlled</b>	<b>Microdochium or Fusarium patch</b> ( <i>Microdochium nivale</i> )
<b>Use rate (product/ha)</b>	95 mL/100 m <sup>2</sup> or 9.5 L/ha
<b>Application Interval</b>	21 to 28 days
<b>Minimum recommended application of water volume</b>	200 to 800 L/ha
<b>Maximum number of applications per crop</b>	4
<b>Maximum rate of active ingredients applied</b>	1 g ai/100 m <sup>2</sup> or 100 g ai/ha of pydiflumetofen 5.94 g ai/100 m <sup>2</sup> or 594 g ai/ha of azoxystrobin 9.88 g ai/100 m <sup>2</sup> or 988 g ai/ha propiconazole



<b>Application Directions</b>	Use preventatively. Begin applications when conditions are favourable for disease development, i.e., extended periods of cool (<15.5°C) and wet weather in spring and fall.  During prolonged periods of favourable conditions apply on the shorter application interval.
<b>Diseases Controlled</b>	<b>Pythium blight</b> ( <i>Pythium aphanidermatum</i> )
<b>Use rate (product/ha)</b>	95 mL/100 m <sup>2</sup> or 9.5 L/ha
<b>Application Interval</b>	10 to 14 days
<b>Minimum recommended application of water volume</b>	200 to 800 L/ha
<b>Maximum number of applications per crop</b>	4
<b>Maximum rate of active ingredients applied</b>	1 g ai/100 m <sup>2</sup> or 100 g ai/ha of pydiflumetofen 5.94 g ai/100 m <sup>2</sup> or 594 g ai/ha of azoxystrobin 9.88 g ai/100 m <sup>2</sup> or 988 g ai/ha propiconazole
<b>Application Directions</b>	Use preventatively. Begin applications when conditions are favourable for disease infection, i.e., relative humidity greater than 90% for at least 9 hours, maximum daily temperature of 28°C and low temperature of 20°C.  During prolonged periods of favourable conditions apply on the shorter application interval.
<b>Diseases Controlled</b>	<b>Fairy ring</b> (Basidiomycete, including <i>Lycoperdon spp.</i> )
<b>Use rate (product/ha)</b>	95 mL/100 m <sup>2</sup> or 9.5 L/ha
<b>Application Interval</b>	28 day
<b>Minimum recommended application of water volume</b>	200 to 800 L/ha
<b>Maximum number of applications per crop</b>	4
<b>Maximum rate of active ingredients applied</b>	1 g ai/100 m <sup>2</sup> or 100 g ai/ha of pydiflumetofen 5.94 g ai/100 m <sup>2</sup> or 594 g ai/ha of azoxystrobin 9.88 g ai/100 m <sup>2</sup> or 988 g ai/ha propiconazole
<b>Application Directions</b>	Apply as soon as possible after fairy ring symptoms develop. Severely damaged or thin turf may require reseeding. Fairy ring symptoms may take 2 to 3 weeks to disappear following application. Reapplication after 28 days may be required.

<b>Diseases Controlled</b>	<b>Anthracnose (<i>Colletotrichum graminicola</i>)</b>
<b>Use rate (product/ha)</b>	48 to 76 mL/100 m <sup>2</sup> or 4.8 to 7.6 L/ha
<b>Application Interval</b>	14 to 21 days
<b>Minimum recommended application of water volume</b>	200 to 800 L/ha
<b>Maximum number of applications per crop</b>	4
<b>Maximum rate of active ingredients applied</b>	0.78 g ai/100 m <sup>2</sup> or 78 g ai/ha of pydiflumetofen 4.75 g ai/100 m <sup>2</sup> or 475 g ai/ha of azoxystrobin 7.90 g ai/100 m <sup>2</sup> or 790 g ai/ha propiconazole
<b>Application Directions</b>	Use preventatively. Begin applications when conditions are favourable for disease infection, prior to disease symptom development.  Use the higher rate under severe disease pressure.  During prolonged periods of favourable conditions apply on the shorter application interval.
<b>Diseases Controlled</b>	<b>Red thread (<i>Laetisaria fuciformis</i>)</b>
<b>Use rate (product/ha)</b>	76 mL/100 m <sup>2</sup> or 7.6 L/ha
<b>Application Interval</b>	14 day
<b>Minimum recommended application of water volume</b>	200 to 800 L/ha
<b>Maximum number of applications per crop</b>	4
<b>Maximum rate of active ingredients applied</b>	0.78 g ai/100 m <sup>2</sup> or 78 g ai/ha of pydiflumetofen 4.75 g ai/100 m <sup>2</sup> or 475 g ai/ha of azoxystrobin 7.90 g ai/100 m <sup>2</sup> or 790 g ai/ha propiconazole
<b>Application Directions</b>	Apply when conditions are favourable for disease development. Prolonged periods of high humidity air favour the development of the disease.
<b>Diseases Controlled</b>	<b>Brown patch (<i>Rhizoctonia solani</i>)</b>
<b>Use rate (product/ha)</b>	48 mL/100 m <sup>2</sup> or 4.8 L/ha
<b>Application Interval</b>	14 day
<b>Minimum recommended application of water volume</b>	200 to 800 L/ha
<b>Maximum number of applications per crop</b>	4
<b>Maximum rate of active ingredients applied</b>	0.49 g ai/100 m <sup>2</sup> or 49 g ai/ha of pydiflumetofen 3.0 g ai/100 m <sup>2</sup> or 300 g ai/ha of azoxystrobin 4.99 g ai/100 m <sup>2</sup> or 499 g ai/ha propiconazole
<b>Application Directions</b>	Begin applications before disease is present when conditions are favorable for disease development (high surface moisture or humidity and warm night temperatures).

---

## 12.0 RESISTANCE MANAGEMENT

For resistance management, A22070 Fungicide contains Group 7, 11 and Group 3 fungicides. Any fungal population may contain individuals naturally resistant to A22070 Fungicide and other Group 7, 11 and 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

### **To delay fungicide resistance:**

Where possible, rotate the use of A22070 Fungicide or other Group 7, 11 or 3 fungicides with different groups that control the same pathogens.

Apply no more than 2 sequential applications before alternating to a non-Group 3, 7 or 11 fungicide.

Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.

Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for sign of resistance development. Notify Syngenta Canada Inc. if reduced sensitivity of the pathogen to A22070 Fungicide is suspected.

If disease continues to progress after treatment with this product, DO NOT increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, to which pathogen resistance has not developed.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information or to report suspected resistance, Contact Company representatives at 1-87-SYNGENTA (1-877-964-3682) or at [www.syngenta.ca](http://www.syngenta.ca).