

2021-05-20
2020-3922

(Box label)

Primidacide II for Cats Over 4 Kg

Flea Adulticide

For cats 7 months and older weighing over 4 kg

[Once-A-Month] [Once Per Month] Topical Flea Treatment

LIQUID

[Rapidly kills fleas] [Kills fleas rapidly] [and] [may reduce flea allergy dermatitis (FAD)]

[Controls fleas within one (1) day of treatment]

[Flea infestation protection for up to four (4) weeks]

[Three] [Four] [Six] [3] [4] [6] [months' supply]

READ THE LABEL AND INSERT BEFORE USING
FOR USE ONLY ON CATS

Do not use on other animals

KEEP OUT OF REACH OF CHILDREN



POISON

WARNING EYE IRRITANT

DOMESTIC

ACTIVE INGREDIENT: Imidacloprid 9.1%

Pyriproxyfen 0.46%

REGISTRATION. NO. 34137 P.C.P. ACT

[Three] [Four] [Six] [x] 0.8 mL Tubes

TWIST N' USE TUBE™

[Simple] [and] [easy to open]

[Tamper-evident] [cap] [tip] [with internal twist-off pin release]

[Hygienic] [single-dose application] [Pharma compliant tube]

[Fusion Animal Health Inc.]

71 Main Street

Exeter, Ontario

NOM 1S3

416-627-9425]

Primidacide II for Cats Over 4 Kg will control fleas on cats within one day of treatment and will continue to protect the cat from further flea infestation for up to four weeks. Reinfestation from the emergence of new fleas in the environment may continue to occur for six weeks or longer after treatment is initiated. Reapply every 4 weeks for continual pest control in situations where there is an on going risk of pest infestation. Treatment must only be repeated if pest problem reoccurs.

Do not shampoo or immerse the animal in water immediately after treatment. The animal can be shampooed, immersed in water or exposed to rain 3 days after treatment. Primidacide II for Cats Over 4 Kg retains its ovicidal and larvicidal efficacy up to 4 weeks following a shampoo treatment, swimming or exposure to rain. The adulticidal effect may be shortened by repeated shampooing, water immersion or exposure to rain. Reapplication may be required after such activities.

[For effective flea control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, a premise treatment with an insecticide labelled for that use may be required.

The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with Primidacide II for Cats Over 4 Kg rapidly kills fleas and may reduce the incidence of this condition.

DIRECTIONS:

This product contains imidacloprid and pyriproxyfen. Do not apply another pest control product such as a shampoo, collar, or powder that contains imidacloprid or pyriproxyfen to the treated animal after applying Primidacide II for Cats Over 4 Kg. Do NOT administer by mouth. Do NOT use on other animals. Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment. A veterinarian should be consulted when pest infestation continues to be a problem on the pet.

Treatment must only be repeated if the pest problem reoccurs.

Do not use on cats under 7 months of age.

Use only on cats weighing over 4 kg.

This product is packaged in single dosage tubes. This package contains [Three] [Four] [Six] [3] [4] [6] monthly 0.8 ml dosages. Treat your cat monthly in the manner described below. Do not apply a dosage higher than one 0.8 ml tube at a time.

Remove one applicator from the package.

Hold applicator tube in upright position.

Twist tip to break seal.

Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to completely empty the contents directly on the skin. Do not get this product in your pet's eyes or mouth. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product. Do not allow recently treated animals to groom each other. Until Primidacide II for Cats Over 4 Kg dries, keep pet off furniture and away from hardwood floors.

Discard the empty tube as described in "DISPOSAL".]

PRECAUTIONS:

HUMANS: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed, or absorbed through the skin. Causes eye irritation. DO NOT get in eyes. Avoid contact with skin and clothing. Wear rubber gloves when applying product. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Avoid contact with pets until dry. Do not contaminate water, feed or food by storage or disposal.

PETS: Do not use on animals under 7 months of age. Care should be taken to prevent exposure of other pets who may be in close physical contact with the treated animal. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. Smaller animals are more likely to experience an adverse reaction. Observe carefully. If signs persist, or become more severe, consult a veterinarian immediately.

Side Effects: Monitor your cat after application. Side effects may include: skin irritation such as scratching or hair loss at the application site, or changes in behaviour such as agitation or lethargy.

Gastrointestinal effects such as drooling, vomiting, or loss of appetite may also occur. If these or other side effects occur, consult your veterinarian or Fusion Animal Health Inc. at 1-800-265-9754.

FIRST AID:

HUMANS: IF IN EYES, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

STORAGE: Store container at room temperature.

DISPOSAL: Do not reuse empty container. Wrap empty tube and cap in newspaper and dispose with household garbage.

[RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE.]

[Fusion Animal Health Inc.

71 Main Street

Exeter, Ontario

N0M 1S3

416-627-9425]

[(Insert label)]

Primidacide II for Cats Over 4 Kg

Flea Adulticide

For cats 7 months and older weighing over 4 kg

[Once-A-Month] [Once Per Month] Topical Flea Treatment

LIQUID

[Rapidly kills fleas] [Kills fleas rapidly] [and] [may reduce flea allergy dermatitis (FAD)]

[Controls fleas within one (1) day of treatment]

[Flea infestation protection for up to four (4) weeks]

[Three] [Four] [Six] [3] [4] [6] [months' supply]

READ THE LABEL AND INSERT BEFORE USING

FOR USE ONLY ON CATS

Do not use on other animals

KEEP OUT OF REACH OF CHILDREN



POISON

WARNING EYE IRRITANT

DOMESTIC

ACTIVE INGREDIENT: Imidacloprid 9.1%

Pyriproxyfen 0.46%

REG. NO. 34137 P.C. P. ACT

[Three] [Four] [Six] 0.8 mL Tubes

TWIST N' USE TUBE™

[Simple] [and] [easy to open]

[Tamper-evident] [cap] [tip] [with internal twist-off pin release]

[Hygienic] [single-dose application] [Pharma compliant tube]

Primidacide II for Cats Over 4 Kg will control fleas on cats within one day of treatment and will continue to protect the cat from further flea infestation for up to four weeks. Reinfestation from the emergence of new fleas in the environment may continue to occur for six weeks or longer after treatment is initiated. Reapply every 4 weeks for continual pest control in situations where there is an ongoing risk of pest infestation. Treatment must only be repeated if pest problem reoccurs.

Do not shampoo or immerse the animal in water immediately after treatment. The animal can be shampooed, immersed in water or exposed to rain 3 days after treatment. Primidacide II for Cats Over 4 Kg retains its ovicidal and larvicidal efficacy up to 4 weeks following a shampoo treatment, swimming or exposure to rain. The adulticidal effect may be shortened by repeated shampooing, water immersion or exposure to rain. Reapplication may be required after such activities.

For effective flea control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, a premise treatment with an insecticide labelled for that use may be required.

The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with Primidacide II for Cats Over 4 Kg rapidly kills fleas and may reduce the incidence of this condition.

PRECAUTIONS:

HUMANS: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed, or absorbed through the skin. Causes eye irritation. DO NOT get in eyes. Avoid contact with skin and clothing. Wear rubber gloves when applying product. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Avoid contact with pets until dry. Do not contaminate water, feed or food by storage or disposal.

PETS: Do not use on animals under 7 months of age. Care should be taken to prevent exposure of other pets who may be in close physical contact with the treated animal. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. Smaller animals are more likely to experience an adverse reaction. Observe carefully. If signs persist, or become more severe, consult a veterinarian immediately.

Side Effects: Monitor your cat after application. Side effects may include: skin irritation such as scratching or hair loss at the application site, or changes in behaviour such as agitation or lethargy. Gastrointestinal effects such as drooling, vomiting, or loss of appetite may also occur. If these or other side effects occur, consult your veterinarian or Fusion Animal Health Inc. at 1-800-265-9754.

FIRST AID:

HUMANS: IF IN EYES, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Take container, label or product name and Pest Control Product Registration Number with you, when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

STORAGE: Store container at room temperature.

DISPOSAL: Do not reuse empty container. Wrap empty tube and cap in newspaper and dispose with household garbage.

DIRECTIONS:

This product contains imidacloprid and pyriproxyfen. Do not apply another pest control product such as a shampoo, collar, or powder that contains imidacloprid or pyriproxyfen to the treated animal after applying Primidacide II for Cats Over 4 Kg. Do NOT administer by mouth. Do NOT use on other animals. Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment. A veterinarian should be consulted when pest infestation continues to be a problem on the pet.

Treatment must only be repeated if the pest problem reoccurs.

Do not use on cats under 7 months of age.

Use only on cats weighing over 4 kg.

This product is packaged in single dosage tubes. This package contains [Three] [Four] [Six] [3] [4] [6] monthly 0.8 ml dosages. Treat your cat monthly in the manner described below. Do not apply a dosage higher than one 0.8 ml tube at a time.

Remove one applicator from the package.

Hold applicator tube in upright position.

Twist tip to break seal.

Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to completely empty the contents directly on the skin. Do not get this product in your pet's eyes or mouth. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product. Do not allow recently treated animals to groom each other. Until Primidacide II for Cats Over 4 Kg dries, keep pet off furniture and away from hardwood floors.

Discard the empty tube as described in "DISPOSAL".

RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE.

Fusion Animal Health Inc.

71 Main Street

Exeter, Ontario

NOM 1S3

416-627-9425

(Tube label)

Primidacide II for Cats Over 4 Kg

READ [THE] LABEL [&] [AND] INSERT BEFORE USING



POISON

WARNING EYE IRRITANT

DOMESTIC

EXTERNAL USE ONLY

ACTIVE INGREDIENT: IMIDACLOPRID 9.1%

Pyriproxyfen 0.46%

REG NO 34137 PCP ACT

Fusion Animal Health Inc.

[71 Main Street

Exeter, Ontario

NOM 1S3]

416-627-9425

0.8 mL