

2023-5988
2023-11-02

Base container label

GROUP	10	HERBICIDE
-------	----	-----------

Tide Glufosinate ammonium 150SL Herbicide and Desiccant

SOLUTION

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

REGISTRATION NUMBER: 34996 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENT: Glufosinate ammonium ... 150 g/L



WARNING: SKIN AND EYE IRRITANT

NET CONTENTS: 1L – Bulk

Tide International Canada Inc.
245 Hanlon Creek Boulevard, Unit 102,
Guelph, Ontario,
N1C 0A1
+1 866 699 7717

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Harmful or fatal if absorbed through the skin. May irritate the skin. Harmful if swallowed. Causes eye irritation. DO NOT get in eyes, on skin or on clothing. Avoid breathing spray mist. Wash thoroughly after using and before eating, drinking or smoking.

Protective Clothing and Equipment: Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes, and protective eyewear (goggles or face shield) during mixing, loading, application, clean-up and repair. Gloves and protective eyewear (goggles or face shield) are not required during application within a closed cab and/or cockpit.

If used in a tank mix, follow the most protective PPE directions of the products used.

Re-entry Restriction: Workers should not enter treated fields within 24 hours of treatment. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice.

Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. DO NOT give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

Note to Physician: If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Avoid spray drift to susceptible plants and use only for recommended purposes and at recommended rates.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

STORAGE

DO NOT store product below freezing.

If stored for 1 year or longer, shake well before using.

Do not use or store in or around the home.

Keep in original container during storage.

Store this product away from food or feed.

DISPOSAL

RECYCLABLE CONTAINER DISPOSAL (less than 23 litres):

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

RETURNABLE CONTAINER DISPOSAL

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

REFILLABLE CONTAINER DISPOSAL

For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

DISPOSAL OF UNUSED, UNWANTED PRODUCT

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

DISPOSAL OF UNUSED SPRAY SOLUTION

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the area just sprayed, away from water supplies, ditches, and irrigation canals.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Tide Glufosinate ammonium 150SL Herbicide and Desiccant

SOLUTION

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

REGISTRATION NUMBER: 34996 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENT: Glufosinate ammonium.....150 g/L



WARNING: SKIN AND EYE IRRITANT

NET CONTENTS: 1L – Bulk

Tide International Canada Inc.
245 Hanlon Creek Boulevard, Unit 102,
Guelph, Ontario,
N1C 0A1
+1 866 699 7717

GENERAL INFORMATION

Tide Glufosinate ammonium 150SL Herbicide and Desiccant provides control of a broad spectrum of grassy and broadleaf weeds in Canola and Soybean varieties or hybrids that are specially developed to be tolerant to glufosinate ammonium.

Tide Glufosinate ammonium 150SL Herbicide and Desiccant is registered for use on glufosinate ammonium tolerant Canola and Soybean varieties or hybrids [example: varieties or hybrids labelled with LibertyLink®].

Tide Glufosinate ammonium 150SL Herbicide and Desiccant may also be applied to glufosinate ammonium tolerant lines or varieties grown for seed production.

For hybrid seed production in canola, Tide Glufosinate ammonium 150SL Herbicide and Desiccant may be used to remove segregating wild type plants within the female population.

Two applications are required to do this. A third application may be necessary if it is determined that the segregating wild type plants are not fully removed from the field.

Glufosinate ammonium tolerant Corn can be used as separator rows in hybrid Canola seed production, but must be destroyed and not used for food or feed.

Tide Glufosinate ammonium 150SL Herbicide and Desiccant may be used as a harvest aid (desiccant) on Alfalfa grown for seed.

Tide Glufosinate ammonium 150SL Herbicide and Desiccant will desiccate weeds that are present in the field at time of application when used as a harvest aid.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Harmful or fatal if absorbed through the skin. May irritate the skin. Harmful if swallowed. Causes eye irritation. **DO NOT** get in eyes, on skin or on clothing. Avoid breathing spray mist. Wash thoroughly after using and before eating, drinking or smoking.

Protective Clothing and Equipment: Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes, and protective eyewear (goggles or face shield) during mixing, loading, application, clean-up and repair. Gloves and protective eyewear (goggles or face shield) are not required during application within a closed cab and/or cockpit.

If used in a tank mixture, follow the most protective PPE directions of the products used.

Re-entry Restriction: Workers should not enter treated fields within 24 hours of treatment. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice.

Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so

by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

Note to Physician: If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Avoid spray drift to susceptible plants and use only for recommended purposes and at recommended rates.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

STORAGE

DO NOT store product below freezing.

If stored for 1 year or longer, shake well before using.

Do not use or store in or around the home.

Keep in original container during storage.

Store this product away from food or feed.

DISPOSAL

RECYCLABLE CONTAINER DISPOSAL (less than 23 litres):

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

3. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
4. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

RETURNABLE CONTAINER DISPOSAL

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

REFILLABLE CONTAINER DISPOSAL

For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

DISPOSAL OF UNUSED, UNWANTED PRODUCT

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

DISPOSAL OF UNUSED SPRAY SOLUTION

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the area just sprayed, away from water supplies, ditches, and irrigation canals.

DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

POST EMERGENT USES

CANOLA – EASTERN CANADA AND COASTAL BRITISH COLUMBIA ONLY

Tide Glufosinate ammonium 150SL Herbicide and Desiccant is registered for aerial and ground application to glufosinate ammonium tolerant Canola varieties or hybrids [example: varieties or hybrids labelled with LibertyLink®].

To assure crop safety and optimal herbicide performance, only use Tide Glufosinate ammonium 150SL Herbicide and Desiccant on glufosinate ammonium tolerant (*i.e.*, LibertyLink) Canola grown from certified seed.

Application of Tide Glufosinate ammonium 150SL Herbicide and Desiccant to non-tolerant Canola varieties or hybrids or other non-tolerant crops will result in severe crop injury or death of the crop.

Apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant from the cotyledon stage up until, but prior to, the early bolting stage of Canola.

Refer to weed/rate chart in this label for application rates and weeds controlled.

For control/suppression of late emerging weeds or if new weed germination or growth is present re-apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant.

Up to three applications of Tide Glufosinate ammonium 150SL Herbicide and Desiccant, each at a maximum of 4 L/ha may be made. The third application must occur prior to the early bolting stage of Canola.

DO NOT apply more than a total of 12 L/ha per year.

Apply when the new weed growth is in the correct leaf stage and up until, but prior to, the bolting stage of Canola.

For best results, apply to emerged, young, actively growing weeds. Weeds that emerge after application will not be controlled.

Tide Glufosinate ammonium 150SL Herbicide and Desiccant will have an effect on weeds that are larger than the recommended leaf stage; however speed of activity and control may be reduced.

Slight discoloration of the Canola may be visible after application. This effect is temporary and will not influence crop growth, maturity or yield.

For application to glufosinate ammonium and glyphosate tolerant canola varieties or hybrids (example: canola varieties or hybrids labelled with LibertyLink[®] and Roundup Ready[®] or TruFlex[™] or Optimum[®] Gly): Tide Glufosinate ammonium 150SL Herbicide and Desiccant should be applied in a sequential program with glyphosate for effective weed control. Tank-mixing Tide Glufosinate ammonium 150SL Herbicide and Desiccant with glyphosate may result in reduced control on certain weed species.

Tide Glufosinate ammonium 150SL Herbicide and Desiccant may also be applied to glufosinate ammonium tolerant Canola lines or varieties grown for seed production.

FOR HYBRID SEED PRODUCTION

Two applications are required to remove the segregating wild type plants. The first application should occur when the Canola is in the 2-4 leaf stage, the second application when the Canola plants are in the 4-6 leaf stage and the third application, if necessary, may be applied to the Canola up until, but prior to, the bolting stage. All applications of Tide Glufosinate ammonium 150SL Herbicide and Desiccant for hybrid seed production should be made at 3.33 – 4 L/ha.

USE OF CORN SEPARATOR ROWS IN HYBRID CANOLA SEED PRODUCTION

Tide Glufosinate ammonium 150SL Herbicide and Desiccant is registered for use on glufosinate ammonium tolerant Corn varieties or hybrids when used as separator rows [example: varieties or hybrids labelled with Liberty Link[®] in hybrid Canola seed production.

Corn should be seeded at 20,000 – 30,000 plants/ha.

Row spacing for glufosinate ammonium tolerant Corn should be 18-23 cm.

NOTE: Corn must be destroyed and not used for food or feed.

Follow use instructions for canola and general instructions in this label.

CANOLA - PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA ONLY

Tide Glufosinate ammonium 150SL Herbicide and Desiccant is registered for aerial and ground application to glufosinate ammonium tolerant Canola varieties or hybrids [example: varieties or hybrids labelled with LibertyLink[®]] grown in the Prairie Provinces and Interior of British Columbia only.

To assure crop safety and optimal herbicide performance, only use Tide Glufosinate ammonium 150SL Herbicide and Desiccant on glufosinate ammonium tolerant (*i.e.*, LibertyLink[®]) Canola grown from certified seed.

Application of Tide Glufosinate ammonium 150SL Herbicide and Desiccant to non-tolerant Canola varieties or hybrids or other non-tolerant crops will result in severe crop injury or death of the crop.

Apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant from the cotyledon stage up until, but prior to, the early bolting stage of Canola.

Refer to weed/rate chart in this label for application rates and weeds controlled.

For control/suppression of late emerging weeds or if new weed germination or growth is present re-apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant.

Up to three applications of Tide Glufosinate ammonium 150SL Herbicide and Desiccant, each at a maximum of 4 L/ha may be made. The third application must occur prior to the early bolting stage of Canola.

DO NOT apply more than a total of 12 L/ha per year.

Apply when the new weed growth is in the correct leaf stage and up until, but prior to, the bolting stage of Canola.

For best results, apply to emerged, young, actively growing weeds. Weeds that emerge after application will not be controlled.

Tide Glufosinate ammonium 150SL Herbicide and Desiccant will have an effect on weeds that are larger than the recommended leaf stage, however speed of activity and control may be reduced.

Slight discoloration of the Canola may be visible after application. This effect is temporary and will not influence crop growth, maturity or yield.

For application to glufosinate ammonium and glyphosate tolerant canola varieties or hybrids (example: canola varieties or hybrids labelled with LibertyLink[®] and Roundup Ready[®] or TruFlex[™] or Optimum[®] Gly): Tide Glufosinate ammonium 150SL Herbicide and Desiccant should be applied in a sequential program with glyphosate for effective weed control. Tank-mixing Tide Glufosinate ammonium 150SL Herbicide and Desiccant with glyphosate may result in reduced control on certain weed species.

Tide Glufosinate ammonium 150SL Herbicide and Desiccant may also be applied to glufosinate ammonium tolerant Canola lines or varieties grown for seed production.

FOR HYBRID SEED PRODUCTION - Two applications are required to remove the segregating wild type plants. The first application should occur when the Canola is in the 2-4 leaf stage, the second application when the Canola plants are in the 4-6 leaf stage and the third application, if necessary, may be applied to the Canola up until, but prior to, the bolting stage. All applications of Tide Glufosinate ammonium 150SL Herbicide and Desiccant for hybrid seed production should be made at 3.33 - 4 L/ha.

USE OF CORN SEPARATOR ROWS IN HYBRID CANOLA SEED PRODUCTION

Tide Glufosinate ammonium 150SL Herbicide and Desiccant is registered for use on glufosinate ammonium tolerant Corn varieties or hybrids when used as separator rows [example: varieties or hybrids labelled with Liberty Link[®]] in hybrid Canola seed production.

Corn should be seeded at 20,000 – 30,000 plants/ha.

Row spacing for glufosinate ammonium tolerant Corn should be 18-23 cm.

NOTE: Corn must be destroyed and not used for food or feed.

Follow use instructions for canola and general instructions in this label.

SOYBEAN

Tide Glufosinate ammonium 150SL Herbicide and Desiccant is registered for aerial and ground application to glufosinate ammonium tolerant Soybean varieties [example: varieties labeled with LibertyLink[®]].

Application of Tide Glufosinate ammonium 150SL Herbicide and Desiccant to non-tolerant Soybean varieties or other non-tolerant crops will result in severe crop injury or death of the crop.

Apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant at 2 L to 3.33 L/ha from the cotyledon to the flowering stage of the crop.

A second application of Tide Glufosinate ammonium 150SL Herbicide and Desiccant can be made to Soybeans treated initially with up to 3.33 L/ha if new weed germination or growth is

present. Apply when the new weed growth is in the correct leaf stage and up to the maximum leaf stage of the crop.

Do not apply beyond the flowering stage of Soybeans.

Do not apply more than a total of 6.67 L/ha in one season.

SUSCEPTIBLE WEEDS:

Tide Glufosinate ammonium 150SL Herbicide and Desiccant has an effect on all weeds and crops except for those crops which are developed to be tolerant to applications of glufosinate ammonium. The following weeds are susceptible to application of Tide Glufosinate ammonium 150SL Herbicide and Desiccant.

Best control will be obtained when Tide Glufosinate ammonium 150SL Herbicide and Desiccant is applied in the recommended leaf stages.

WEED	RECOMMENDED WEED LEAF STAGE								RATE 1.33 L/ha
	1	2	3	4	5	6	7	8	
Cow Cockle	-----								
Green Foxtail	----- Maximum 3 tillers								

WEED	RECOMMENDED WEED LEAF STAGE								RATE 2 L/ha
	1	2	3	4	5	6	7	8	
Barnyard Grass	-----								
Lady's-thumb	-----								
Lamb's-quarters	-----								
Russian Thistle	Up to 8 cm height								
Smartweed	-----								
Stinkweed	-----								
Volunteer Flax	Up to 6 cm height								
Wild Mustard	-----								

WEED	RECOMMENDED WEED LEAF STAGE								RATE 2.67 L/ha
	1	2	3	4	5	6	7	8	
Canada Thistle ¹	Up to 10 cm height								
Common Chickweed	----- Leaf Pairs								
Hemp-nettle	----- Leaf Pairs								
Kochia	Up to 8 cm height								
Perennial Sow Thistle	-----								
Quackgrass ¹	-----								
Redroot Pigweed	-----								
Round-leaved Mallow	-----								
Scentsless Chamomile	Up to 10 cm height								
Shepherd's-purse	-----								
Volunteer Barley ²	----- Maximum 2 tillers								

Volunteer Wheat -----| Maximum 2 tillers
 Wild Buckwheat -----|

WEED	RECOMMENDED WEED LEAF STAGE								RATE
	1	2	3	4	5	6	7	8	
									3.33 L/ha
Cleavers	----- Whorls								
Dandelion	1-15 cm rosette								
Flixweed	Up to 10 cm height								
Hemp-nettle	----- Leaf Pairs								
Jimsonweed	-----								
Quackgrass ³	-----								
Stork's-bill	-----								
Wild Oats	----- Maximum 2 tillers								

WEED	RECOMMENDED WEED LEAF STAGE								RATE
	1	2	3	4	5	6	7	8	
									3.33 L/ha
Heavy Populations									
Canada Thistle ¹	Up to 10 cm height								
Quackgrass ¹	-----								
Volunteer Barley ²	----- Maximum 2 tillers								
Volunteer Wheat	----- Maximum 2 tillers								
Wild Buckwheat	-----								

WEED	RECOMMENDED WEED LEAF STAGE								RATE
	1	2	3	4	5	6	7	8	
									4 L/ha
Canada Thistle ⁴	Up to 10 cm height								
Jimsonweed ⁵	-----								
Quackgrass ⁶	-----								
Japanese Brome ⁷	----- up to emergence of 1st tiller								
Downy Brome ⁷	----- up to emergence of 1st tiller								

NOTES: 1 Top Growth Suppression Only
 2 Suppression Only
 3 Improved Top Growth Control
 4 Better Top Growth Suppression
 5 Improved Control
 6 Season Long Control for Heavy Populations
 7 Spring-germinated Brome Only; best results are obtained after a pre-seed or burn-down application with a glyphosate herbicide.

TANK MIXES

Do not tank mix Tide Glufosinate ammonium 150SL Herbicide and Desiccant with herbicides, fertilizers or chemical additives unless recommended on this label.

When applied as a tank-mix combination, read and observe all label directions, including rates, personal protective equipment, restrictions and precautions for each product used in the tank-mix. Always use in accordance with the most restrictive label restrictions and precautions. Tank mix partners may have additional restrictions for leaf staging. For enhanced activity Tide Glufosinate ammonium 150SL Herbicide and Desiccant may be tank mixed with the following products:

CANOLA

Tank Mix Product	Rate	Directions for Use
Centurion [®] or Select [®]	63 mL/ha	For control of Volunteer Barley and Wild Oats, apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant at a rate of 2.67 – 4.0 L/ha plus Centurion [®] or Select [®] at 63 mL/ha with adjuvant as recommended on the Centurion [®] or Select [®] label. Apply when the weeds are in the 1-4 leaf stage with a maximum of 2 tillers.
Centurion [®] or Select [®]	190 mL/ha	For control of weed species listed on the Tide Glufosinate ammonium 150SL Herbicide and Desiccant and Centurion or Select labels plus foxtail barley, spring-germinated Japanese and downy brome, apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant at a rate of 3.33 to 4.0 L/ha in tank mixture with Centurion [®] or Select [®] at 190 mL/ha and adjuvant as recommended on the Centurion or Select label. Apply when the weeds are at growth stages listed on the Tide Glufosinate ammonium 150SL Herbicide and Desiccant, Centurion [®] and Select [®] labels, foxtail barley is at the 1-4 leaf stage with a maximum of 2 tillers and spring-germinated Japanese and downy brome are at the 1 to 6-leaf up to the emergence of the 1st tiller stage.
Facet [®] L + Merge [®]	0.28 L/ha + 0.5 – 1.0 L/ha	For enhanced and more consistent control of cleavers, apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant at a rate of 3.33 L/ha plus Facet [®] L at 0.28 L/ha. Apply in 100 L/ha of water with Merge [®] Adjuvant at a rate of 0.5 to 1 L/ha. Application should be made from the 2 to 6 true leaf stage of the canola crop.
Facet [®] L + Centurion [®] + Amigo [®] or Merge [®]	0.28 L/ha + 63 mL/ha + 0.5 L/ha	For enhanced and more consistent control of cleavers and annual grasses, apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant at a rate of 3.33 L/ha plus Facet [®] L at 0.28 L/ha plus Centurion [®] at 63 mL/ha. Apply in 100 L/ha of water with Amigo [®] or Merge [®] Adjuvant at 0.5 L/ha. Application should be made from the 2 to 6 true leaf stage of the canola crop.

DESICCATION USE

Tide Glufosinate ammonium 150SL Herbicide and Desiccant may be applied for desiccation, by ground or air, to Alfalfa (grown for seed).

Tide Glufosinate ammonium 150SL Herbicide and Desiccant will also desiccate weeds, which are present in the field at time of application (Wild Buckwheat may not be completely desiccated).

Desiccation of crops and weeds will be best when environmental conditions are favourable (warm temperatures, good moisture conditions, high humidity).

Crop	Rate	Directions for Use
Alfalfa (grown for seed)	2.7 L/ha	Apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant at 50-75% pod turn (brown) stage. Do NOT apply more than once per year.
Note: Do NOT tank mix Tide Glufosinate ammonium 150SL Herbicide and Desiccant with pesticides, fertilizers or chemical additives unless recommended on this label.		

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, Tide Glufosinate ammonium 150SL Herbicide and Desiccant is a Group 10 herbicide. Any weed population may contain or develop plants naturally resistant to Tide Glufosinate ammonium 150SL Herbicide and Desiccant and other Group 10 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Tide Glufosinate ammonium 150SL Herbicide and Desiccant or other Group 10 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates, precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.

- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Tide International Canada Inc at +1 866 699 7717.

GENERAL APPLICATION INSTRUCTIONS

The speed of action of Tide Glufosinate ammonium 150SL Herbicide and Desiccant is influenced by environmental factors. At cool temperatures (below 10° C), poor moisture and low humidity, the speed of action may be reduced. Generally, visual symptoms appear 2-4 days after application.

For best results, apply to emerged, young, actively growing weeds. Weeds that emerge after application will not be controlled.

Tide Glufosinate ammonium 150SL Herbicide and Desiccant will have an effect on weeds that are larger than the recommended leaf stage; however speed of activity and control may be reduced.

Tide Glufosinate ammonium 150SL Herbicide and Desiccant works primarily as a contact herbicide. Thorough coverage of the weeds to be controlled, or the crop to be desiccated, is essential.

When a rate range is given the higher rate should be utilized when the:

- weed or crop growth is dense;
- weeds are large and/or mature - *i.e.*, advanced leaf stages and plant height;
- environmental conditions are cool and dry.

Refer to the section on weeds controlled for details on rates and timing of application.

GROUND APPLICATION INSTRUCTIONS

Ensure that all circuits (pipes, booms etc.) have the correct Tide Glufosinate ammonium 150SL Herbicide and Desiccant/water concentration before application is started.

Apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant in a minimum of 110 L/ha of water, at a pressure of 275 kPa and at a ground speed of 6-8 kph. If check valves are used, apply at 310 kpa. The use of 80° or 110° flat fan nozzles is highly recommended for optimum spray coverage and canopy penetration.

Application of the spray at a 45° angle forward will result in better spray coverage. Refer to sections above for the correct rate and timing of application.

For desiccation uses only, where the crop canopy is dense or weed growth is heavy, achieve better spray coverage with higher spray volumes. Under these conditions, apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant in a minimum of 170-220 L/ha of water.

AERIAL APPLICATION INSTRUCTIONS

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Exercise extreme caution during the aerial application of any insecticide, herbicide or fungicide. Drift of pesticides is not always visible with the human eye. Small droplets may drift into sensitive areas without obvious signs of danger. Follow these directions precisely.

When applying Tide Glufosinate ammonium 150SL Herbicide and Desiccant by aircraft, uniform spray coverage is essential. Applicators are required to use the correct combination of spray nozzle tips, nozzle placement and spray pressures which will provide a coarse droplet size distribution with a volume mean diameter greater than 350 microns. Do not use raindrop nozzles. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Post-emergent Use in Glufosinate Ammonium Tolerant Canola and Soybean Water Volume:

Apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant in a minimum of 55 L/ha of water only.

Desiccation Use Only Water Volume Range: If the crop canopy is dense or weed growth is heavy, achieve better spray coverage with higher spray volumes. Under these conditions, if using aerial application, apply Tide Glufosinate ammonium 150SL Herbicide and Desiccant in a minimum of 33-55 L/ha of water.

APPLICATION PRECAUTIONS

Weed control may be reduced when heavy dew, fog or mist is present at the time of application.

Uniform, thorough spray coverage is important to achieve consistent weed control.

Crop injury may result if Tide Glufosinate ammonium 150SL Herbicide and Desiccant is applied to a crop stressed by severe weather conditions, frost, drought, water-saturated soil, low fertility, disease or insect damage.

Tide Glufosinate ammonium 150SL Herbicide and Desiccant is a non-selective herbicide and has the potential to desiccate, defoliate or kill all green plants. Avoid contact with other desirable plants or crops by direct application or from spray drifts as severe damage may occur.

Application of Tide Glufosinate ammonium 150SL Herbicide and Desiccant to non-tolerant Canola or Soybean varieties, hybrids or other non-tolerant crops will result in severe crop injury or death of the crops.

To assure crop safety and optimal herbicide performance, only use Tide Glufosinate ammonium 150SL Herbicide and Desiccant on glufosinate ammonium tolerant (i.e., LibertyLink) Canola or Soybean grown from certified seed.

PLANT-BACK INTERVALS

No plant-back interval is required for field corn, canola and soybeans, dry common beans (not grown for seed), alfalfa, carrot, lettuce, onion and potato.

70 days for buckwheat, barley, millet, oats, rye, sorghum, triticale, wheat, other root and leafy vegetables.

120 days for all other crops.

GROUND APPLICATION PRECAUTIONS

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Do NOT apply if winds exceed 16 kph when using open boom sprayers for ground application.

Do NOT apply if winds exceed 25 kph when using hooded sprayer for ground application.

Do not use flood jet nozzles, controlled droplet application equipment or air-assisted spray equipment.

AERIAL APPLICATION PRECAUTIONS

Aerial application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

Read and understand the entire label before opening this product. If you have questions, call Tide International Canada Inc. or obtain technical advice from the distributor or your provincial agricultural representative.

AERIAL APPLICATION OPERATOR PRECAUTIONS

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

RAINFALL PERIOD

If rainfall occurs within 4 hours of application, effectiveness may be reduced.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- Uses with hand-held application equipment permitted on this label
- Low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit, or foliage.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Application method	Crop	Spray buffer zones (metres) required for the protection of:		
		Freshwater habitat of depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	
Field sprayer	Alfalfa, Canola, Corn,	1	0	1

	Soybean				
Aerial	Alfalfa, soybean (fixed wing and rotary)	1	0	20	
	Canola	Fixed wing	1	0	35
		Rotary wing	1	0	30

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

MIXING DIRECTIONS

Tide Glufosinate ammonium 150SL Herbicide and Desiccant must be applied with properly calibrated, clean equipment.

Tide Glufosinate ammonium 150SL Herbicide and Desiccant is specially formulated to mix readily in water.

Prior to adding Tide Glufosinate ammonium 150SL Herbicide and Desiccant to the spray tank, ensure that the spray tank is thoroughly clean (see section on Sprayer Cleanup).

1. Fill the tank three-quarters full with clean water.
2. Add the correct amount of Tide Glufosinate ammonium 150SL Herbicide and Desiccant.
3. Add the remaining amount of water, begin agitation, and spray out immediately.

TANK-MIXES

When applied as a tank-mix combination, read and observe all label directions, including rates, personal protective equipment, restrictions and precautions for each product used in the tank mix. Always use in accordance with the most restrictive label restrictions and precautions.

Tide Glufosinate ammonium 150SL Herbicide and Desiccant and Centurion[®] or Select[®]

For use on Glufosinate Ammonium Tolerant Canola only:

When tank mixing Tide Glufosinate ammonium 150SL Herbicide and Desiccant and Centurion[®] or Select[®], always add Amigo[®] adjuvant to the tank first, then add Tide Glufosinate ammonium 150SL Herbicide and Desiccant to the tank followed by the Centurion[®] or Select[®].

The addition of an anti-foaming agent may reduce foaming, especially when using soft water.

1. Thoroughly clean the sprayer by flushing the system with water containing detergent (see section on Sprayer Cleanup).
2. Fill clean spray tank half full with clean water. Start agitation system.
3. Add the correct amount of Amigo[®]. Continue to agitate until Amigo[®] is thoroughly mixed.
4. STOP agitation. Add the required amount of Tide Glufosinate ammonium 150SL Herbicide and Desiccant to the spray tank. Start agitation system.
5. Add the correct amount of Centurion[®] or Select[®] along with the remaining amount of

- water necessary to fill the spray tank.
6. Continue to agitate or run the by-pass system and spray out immediately.
 7. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to remix the spray materials. Do not allow the mixture to sit overnight.
 8. If an oil film starts to build up in the tank, drain it and clean tank with strong detergent solution.
 9. Immediately after use, thoroughly clean the sprayer by flushing the system with water containing detergent (see section on Sprayer Cleanup).

SPRAYER CLEANUP

Before and after using Tide Glufosinate ammonium 150SL Herbicide and Desiccant always complete a thorough cleaning of the spray tank, lines, and filter. Spray equipment should be thoroughly rinsed using a strong detergent solution.

LIVESTOCK FEEDING

CANOLA: Grain and meal from treated crop can be fed to livestock. DO NOT graze the treated crop or cut for hay; sufficient data are not available to support such use.

CORN (used as separator rows in Hybrid Canola Seed Production): DO NOT feed treated crop to livestock; sufficient data are not available to support such use.

SOYBEAN: DO NOT graze the treated fields within 20 days of application.

PRE-HARVEST INTERVALS (PHI)

CANOLA: When Tide Glufosinate ammonium 150SL Herbicide and Desiccant is tank mixed with Centurion or Select, observe a PHI of 60 days from the date of treatment (or last treatment when a second application has been made). When Tide Glufosinate ammonium 150SL Herbicide and Desiccant is tank mixed with Facet L, observe a PHI of 60 days from the date of treatment.

SOYBEAN: Do not apply within 70 days of harvesting soybean seed.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.