

2022-1140
2024-04-26

FIESTA Granule Selective Turf Weed Killer

READY TO USE
GRANULE
HERBICIDE
COMMERCIAL

ACTIVE INGREDIENT: Iron (present as ferric sodium EDTA trihydrate) 1.82%

Warning, contains the allergen soy

REG. NO. xxxxx P.C.P. ACT

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS 1 - 1000 kg

Registrant:

W. Neudorff GmbH KG

An der Muhle 3,

31860 Emmerthal, Germany

www.neudorffpro.com

Canadian Agent:

Neudorff North America

PO Box 178

Brentwood Bay, BC V8M1R3

Telephone: (250) 652-5888

For use on residential, commercial, industrial and municipal lawns and turf on rights of way, landscaping, non-crop areas, golf courses, parks, cemeteries and athletic fields consisting of one or more of Kentucky bluegrass, perennial ryegrass, and fescue (tall fescue, chewings fescue, and/or creeping fescue) for post-emergent control of Black medic (*Medicago lupulina*), Broadleaf plantain (*Plantago major*), Canada thistle (*Cirsium arvense*), Creeping buttercup (*Ranunculus repens*), Dandelion (*Taraxacum officinale*), and White clover (*Trifolium repens*). For pre-emergent control or suppression of crabgrass.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes, skin, or clothing. Avoid inhaling the product. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during loading, application, clean-up and repair. Wash hands thoroughly with soap and water after handling and before eating, drinking, and chewing gum or chewing tobacco. DO NOT enter or allow people or pets to enter treated areas until dusts have settled.

FIRST AID:

If swallowed:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes:

Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Post-Emergent Control: Apply at a rate of 15-40 g/m². The standard rate for this product is 25 g/m². Reapply after 3-4 weeks for best results. See table below for application rate to control specific weed species.

Pre-Emergent Control (Crabgrass): Apply when soil temperatures reach 10–15 °C at 40 g/m² to control emergence of smooth crabgrass and suppress emergence of large crabgrass. Reapply after 3 – 6 weeks for best results.

Rate of Application	Weeds Controlled	(Optional column) Area Treated
15 g/m ² or 1.4 kg /93 m ² (1000 ft ²)	Weed seedlings of Dandelion (<i>Taraxacum officinale</i>) and White clover (<i>Trifolium repens</i>) (suppression only)	67 m ² (721 ft ²)/kg
25 g/m ² or 2.3 kg /93 m ² (1000 ft ²)	Black medic (<i>Medicago lupulina</i>) and Dandelion (<i>Taraxacum officinale</i>)	40 m ² (431 ft ²)/kg
40 g/m ² or 3.7 kg/93 m ² (1000 ft ²)	Broadleaf plantain (<i>Plantago major</i>), Creeping buttercup (<i>Ranunculus repens</i>), Canada thistle (<i>Cirsium arvense</i>), and White clover (<i>Trifolium repens</i>). For Canada thistle, best results are obtained when weeds are less than 15 cm tall and two applications may be required for control. Pre-emergent control of smooth crabgrass and pre-emergent suppression of large crabgrass	25 m ² (269 ft ²)/kg

How to Apply:

Contents may have settled during shipment; invert bag several times to ensure product is mixed.

One kg of product will treat between 25 and 67 m² (269 - 721 ft²). For best results, apply a few days after mowing when temperature is between 10°C and 30°C. If temperature is below 27°C, ensure turf is wet from dew, or water turf lightly before application. If temperature is above 27°C, do not pre-wet grass and apply product in the late

afternoon or evening. Do not exceed the 15 g/m² rate when temperature is above 30°C. Do not apply when daytime temperature exceeds 32°C.

Apply using a broadcast/rotary, drop, or handheld spreader calibrated according to the manufacturer's directions. Uniform coverage is important to prevent turf damage. Turf areas can be re-sown one day after treatment. Apply product only on turf. Application on hard surfaces can cause staining. If product gets on surface, sweep off if the surface is dry, rinse off immediately if the surface is wet. If product gets on clothing or footwear, wash immediately.

Visible signs of control may be seen within 24 hours after application. Susceptible weeds will turn brown or black. Occasionally a darkening of the grass leaf blades can occur after treatment however, the grass will recover within a few days to several weeks. May stain thatch (dead grass under turf); this does not affect the health of the grass. If turf is damaged, irrigation 24 hours after application, followed by mowing to remove any turf tip burn, can promote faster recovery.

Use Precautions:

- Do not apply to drought stressed grass; ensure that the turf is well watered prior to application.
- Do not apply when daytime temperature is above 32°C.
- Do not apply in windy conditions.
- Do not apply to bentgrass. If unsure of grass sensitivity, test apply to a small area of turf.
- Apply this product only to turf, and sweep any product that lands on the driveway, sidewalk or street back onto the treated area.
- Avoid contact with desirable plants. Any desirable plants or structures accidentally contacted should be rinsed immediately with water.
- Do not apply if rainfall is expected within 24 hours of application.
- Do not irrigate within 24 hours of application.

STORAGE:

Store this product away from food or feed.

DISPOSAL

1. Thoroughly empty the contents of the container into the application device.
2. Follow provincial/territorial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial/territorial requirements.
5. **For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for cleanup of spills.**

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Canadian Patent No. 2,477,493

Manufactured for:

W. Neudorff GmbH KG
An der Muhle 3, 31860 Emmerthal, Germany

Canadian Office: Neudorff North America
PO Box 178 • Brentwood Bay, BC • V8M 1R3 • 250-652-5888

FIESTA® is a registered trademark of W. Neudorff GmbH KG



Pull here to open and press to reseal.

Optional marketing statements:

SELECTIVE BROADLEAF WEED KILLER

No unpleasant odor

Made with iron

Ready-to-use. No mixing necessary

(1 kg) Will treat up to 66.7m² (718 sq. ft.)

Prevents crabgrass and broadleaf weeds from germinating

Granular herbicide for weed control in turf
Selective granular broadleaf weed killer
Can apply 1 day before planting turf