

2022-1139
2024-04-26

Fiesta Granule Selective Weed Killer RTU

**GRANULE
HERBICIDE
DOMESTIC
READY TO USE**

ACTIVE INGREDIENT: Iron (present as ferric sodium EDTA trihydrate) 1.82%

Warning, contains the allergen soy

REG. NO. 35169 P.C.P. ACT

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS 1 - 6 kg

Registrant:

W. Neudorff GmbH KG

An der Muhle 3,

31860 Emmerthal, Germany

Canadian Agent:

Neudorff North America

PO Box 178

Brentwood Bay, BC V8M1R3

Telephone: (250) 652-5888

2022-1139

2024-04-26

PESTS/SITES:

For the post-emergent control of the following lawn weeds: Black medic, Broadleaf plantain, Canada thistle, Creeping buttercup, Dandelion, and White clover. For pre-emergent control or suppression of crabgrass.

For use on lawn consisting of one or more of Kentucky bluegrass, perennial ryegrass, and fescue (tall fescue, chewing fescue, and/or creeping fescue).

PRECAUTIONS:

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin, eyes or clothing. Avoid inhaling the product. Wear waterproof gloves during loading, application, clean-up and repair. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. DO NOT enter or allow people or pets to enter treated areas until dusts have settled.

FIRST AID:

If swallowed:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes:

Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

DIRECTIONS FOR USE:

DO NOT apply to any body of water

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Post-Emergent Control: Apply at a rate of 15-40 g/m². The standard rate for this product is 25 g/m². Use the higher rate (40 g/m²) on the following tough perennial weeds: broadleaf plantain, creeping buttercup, white clover and Canada thistle. For Canada thistle, best results are obtained when weeds are less than 15 cm tall. Use the lower rate (15 g/m²) to control seedlings of dandelion and to suppress seedlings of white clover. Reapply after 3-4 weeks for best results.

Pre-Emergent Control (Crabgrass): Apply when soil temperatures reach 10–15 °C at 40 g/m² to control emergence of smooth crabgrass and suppress emergence of large crabgrass. Reapply after 3 – 6 weeks for best results.

How to Apply:

2022-1139

2024-04-26

Contents may have settled during shipment; invert bag several times to ensure product is mixed.

For best results, apply a few days after mowing when temperature is between 10°C and 30°C. If temperature is below 27°C, ensure lawn is wet from dew, or water lawn lightly before application. If temperature is above 27°C, do not pre-wet grass and apply product in the late afternoon or evening. Do not exceed the 15 g/m² rate when temperature is above 30°C. Do not apply when daytime temperature exceeds 32°C.

Uniform coverage is important. Uneven application may cause lawn damage. For more uniform coverage, apply at half the labelled rate and go over lawn two times. Use a rotary or handheld spreader set to the correct application rate according to the manufactures directions. See table below. Lawn can be re-sown one day after treatment.

{Optional table:}

Weed Species	Rate	{Optional column} Scotts® Rotary Spreader Setting (All Models)	{Optional column} Scotts® Drop Spreader Setting (All Models)	{Optional column} Scotts® Hand-Held Spreader Setting (All models)
Weed seedlings of Dandelion, White clover	15 g/m ²	Set dial to 2.5 setting. To apply product begin walking, then squeeze and hold trigger.	Set dial to 5 setting. To apply product begin walking, then squeeze and hold trigger.	Adjust spreader selector setting to 3. Pull trigger and turn crank clockwise while walking at normal speed across area to be covered with product
Black medic, Dandelion	25 g/m ²	Set dial to 3 setting. To apply product begin walking, then squeeze and hold trigger.	Set dial to 6 setting. To apply product begin walking, then squeeze and hold trigger.	Adjust spreader selector setting to 3.5. Pull trigger and turn crank clockwise while walking at normal speed across area to be covered with product
Broadleaf plantain, Creeping buttercup, Canada thistle, White clover, Smooth crabgrass, Large crabgrass	40 g/m ²	Set dial to 4.5 setting. To apply product begin walking, then squeeze and hold trigger.	Set dial to 7.5 setting. To apply product begin walking, then squeeze and hold trigger.	Adjust spreader selector setting to 5. Pull trigger and turn crank clockwise while walking at normal speed across area to be covered with product
Notes		Spreader Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application.		

What to expect:

Visible signs of control may be seen within 24 hours after application. Susceptible weeds, moss or algae will turn brown or black. Occasionally a darkening of the grass leaf blades can occur after treatment; however the grass will recover within a few days to several weeks. If lawn is damaged, irrigation 24 hours after application, followed by mowing to remove the lawn tip burn, can help lawn recover faster. May stain thatch (dead grass under lawn); this does not affect the health of the grass.

Use Precautions:

-Do not apply to drought stressed grass; ensure that the lawn is well watered prior to application.

2022-1139

2024-04-26

- Do not apply when the daytime temperature will exceed 32°C.
- Do not apply in windy conditions.
- Do not apply to bentgrass. If unsure of grass sensitivity, test apply to a small area of your lawn.
- Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk or street back onto the treated area.
- Avoid contact with desirable plants. Any desirable plants or structures accidentally contacted should be rinsed immediately with water.
- Do not apply if rainfall is expected within 24 hours of application.
- Do not irrigate within 24 hours of application.

STORAGE:

Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals. Store unused material, tightly closed, in original container only, away from open flame. **Store this product away from food or feed.**

DISPOSAL:

DO NOT reuse the empty containers. Dispose in household garbage. Unused or partially used products should be disposed at provincially/territorially or municipally designated hazardous waste disposal sites.

[NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.]



[optional marketing statements:

SELECTIVE BROADLEAF HERBICIDE

No unpleasant odor

Made with iron

Ready-to-use

(1 kg) Will treat up to 66.7 m² (718 sq. ft)

Second application may be required

A selective granular herbicide for weed control in lawns

Can apply 1 day before planting lawn