

## ALB AZOXYPROP

Fungicide

COMMERCIAL

SUSPENSION

For Use in Controlling Diseases on Cereals

### ACTIVE INGREDIENT:

Azoxystrobin ..... 143 g/L  
Propiconazole ..... 124 g/L

Contains 1,2-benzisothiazolin-3-one at 0.0251% as a preservative, OR contains Bronopol at 0.0114%, 5-Chloro-2-methyl-4-isothiazolin-3-one at 0.0011% and 2-Methyl-4-isothiazolin-3-one at 0.00036% as preservatives.

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**



**POISON**

**DANGER: EYE & SKIN IRRITANT**

REGISTRATION NO: 35083      PEST CONTROL PRODUCTS ACT

**NET CONTENTS:** 1-1000L

**Albaugh, LLC**  
1525 NE 36<sup>th</sup> Street  
Ankeny, IA 50021  
1-800-247-8013

For medical or treatment information from exposure to this product, call 1-888-347-6732  
(7 days/week, 24-hr).

For 24-hour chemical spill, leak, fire, exposure, or accident response information, call CHEMTREC toll  
free at 1-800-424-9300.

## **NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

## **FIRST AID**

**IN CASE OF POISONING**, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**IF SWALLOWED**, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING**, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF IN EYES**, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**IF INHALED**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

## **TOXICOLOGICAL INFORMATION**

There is no specific antidote if this product is ingested. Treat symptomatically.

## **PRECAUTIONS**

**KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED.** Severely irritating to the eye. **DO NOT** get in eyes. May irritate the skin. Avoid contact with skin and clothing. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing and loading. Gloves are not required during application within a closed cab.

Wash with soap and water after handling, and before eating, drinking or smoking. Remove clothing immediately if pesticide gets inside. Wash contaminated clothing, separately from household laundry, before re-use. Do not wear contaminated shoes. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**DO NOT** allow entry into treated areas for 12 hours following application.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools, and recreation areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

## **ENVIRONMENTAL PRECAUTIONS**

**TOXIC** to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under **DIRECTIONS FOR USE**.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff maybe reduced by including a vegetative filter strip between the treated area and the edge of the water body.

Azoxystrobin is persistent and will carryover. It is recommended that this product not be used in areas treated with azoxystrobin during the previous season.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland.

## **STORAGE**

Store this product away from food or feed. Keep in original container, tightly closed, during storage. Store in a cool, dry, well ventilated area and out of the reach of children and animals.

Do not store below 0°C.

## **SPILL CLEANUP**

Wear appropriate protective equipment (gloves, glasses, apron) when attempting to clean up the spill. If the container is leaking, secure leak and place the container into a drum or heavy gauge plastic bag. See Emergency Number for further information.

For spills and leaks - contain the liquid with dikes of inert material (soil, clay, kitty litter, etc.). Absorb the spill onto inert material and shovel into a sealable waste container.

On hard surfaces - sprinkle spill area with detergent and scrub in a small quantity of water with a coarse broom. Let stand 10 minutes then absorb onto an inert material and shovel into the waste container.

On soil - remove the top 15 cm of soil in the spill area and replace with fresh soil. Dispose of all waste including scrub brush in accordance with provincial requirements.

## **DISPOSAL**

### **DISPOSAL OF CONTAINER:**

#### **For returnable containers:**

**DO NOT** reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

#### **For refillable containers:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. **DO NOT** reuse this container for any other purpose.

#### **For recyclable containers:**

**DO NOT** reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

**Disposal of unused, unwanted product**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

## ALB AZOXYPROP

Fungicide

COMMERCIAL

SUSPENSION

For Use in Controlling Diseases on Cereals

### ACTIVE INGREDIENT:

Azoxystrobin ..... 143 g/L  
Propiconazole ..... 124 g/L

Contains 1,2-benzisothiazolin-3-one at 0.0251% as a preservative, OR contains Bronopol at 0.0114%, 5-Chloro-2-methyl-4-isothiazolin-3-one at 0.0011% and 2-Methyl-4-isothiazolin-3-one at 0.00036% as preservatives.

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**



**POISON**

**DANGER: EYE & SKIN IRRITANT**

REGISTRATION NO: 35083 PEST CONTROL PRODUCTS ACT

**NET CONTENTS:** 1-1000L

**Albaugh, LLC**  
1525 NE 36<sup>th</sup> Street  
Ankeny, IA 50021  
1-800-247-8013

For medical or treatment information from exposure to this product, call 1-888-347-6732  
(7 days/week, 24-hr).

For 24-hour chemical spill, leak, fire, exposure, or accident response information, call CHEMTREC toll  
free at 1-800-424-9300.

## NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

## FIRST AID

**IN CASE OF POISONING**, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**IF SWALLOWED**, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING**, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF IN EYES**, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**IF INHALED**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

## TOXICOLOGICAL INFORMATION

There is no specific antidote if this product is ingested. Treat symptomatically.

## PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED. Severely irritating to the eye. DO NOT get in eyes. May irritate the skin. Avoid contact with skin and clothing. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing and loading. Gloves are not required during application within a closed cab.

Wash with soap and water after handling, and before eating, drinking or smoking. Remove clothing immediately if pesticide gets inside. Wash contaminated clothing, separately from household laundry, before re-use. Do not wear contaminated shoes. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**DO NOT** allow entry into treated areas for 12 hours following application.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools, and recreation areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

## ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff maybe reduced by including a vegetative filter strip between the treated area and the edge of the water body.

Azoxystrobin is persistent and will carryover. It is recommended that this product not be used in areas treated with azoxystrobin during the previous season.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland.”

## **STORAGE**

Store this product away from food or feed. Keep in original container, tightly closed, during storage. Store in a cool, dry, well ventilated area and out of the reach of children and animals.

Do not store below 0°C.

## **SPILL CLEANUP**

Wear appropriate protective equipment (gloves, glasses, apron) when attempting to clean up the spill. If the container is leaking, secure leak and place the container into a drum or heavy gauge plastic bag. See Emergency Number for further information.

For spills and leaks - contain the liquid with dikes of inert material (soil, clay, kitty litter, etc.). Absorb the spill onto inert material and shovel into a sealable waste container.

On hard surfaces - sprinkle spill area with detergent and scrub in a small quantity of water with a coarse broom. Let stand 10 minutes then absorb onto an inert material and shovel into the waste container.

On soil - remove the top 15 cm of soil in the spill area and replace with fresh soil. Dispose of all waste including scrub brush in accordance with provincial requirements.

## **DISPOSAL**

### **DISPOSAL OF CONTAINER:**

#### **For returnable containers:**

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

#### **For refillable containers:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

#### **For recyclable containers:**

**DO NOT** reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

**Disposal of unused, unwanted product**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

**PRODUCT INFORMATION**

ALB AZOXYPROP is a broad spectrum, preventative fungicide with systemic and curative properties recommended for the control of important plant diseases in Crop Group 15 - Cereals. ALB AZOXYPROP may be applied as a foliar spray in alternating spray programs. All applications should be made according to the use directions that follow.

**GENERAL USE PRECAUTIONS**

DO NOT apply ALB AZOXYPROP through irrigation equipment.

DO NOT use in nurseries, greenhouses or landscape plantings.

Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications.

**Rotational Crop Precautions**

To avoid possible residues above the acceptable limit, DO NOT plant any other crop intended for food, grazing, or any component of animal feed or bedding within 105 days of AXOXYPROP fungicide application to the preceding crop unless the second crop appears on this label.

Oats and rye may be planted 45 days after ALB AZOXYPROP application.

**Phytotoxicity**

ALB AZOXYPROP demonstrates some phytotoxic effects when mixed with products that are formulated as Emulsifiable Concentrate. These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone may contribute to phytotoxicity.

**INTEGRATED PEST MANAGEMENT**

ALB AZOXYPROP should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. The DIRECTIONS FOR USE section in this label identifies specific IPM recommendations for each crop. Consult your local agricultural authorities for additional IPM strategies established for your area.

**DIRECTIONS FOR USE**

As this product is not registered for the control of pests in aquatic systems.

**DO NOT** use to control aquatic pests.

**DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of waste.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each



product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Albaugh LLC at 1-800-247-8013 for information before applying any tank mix that is not specifically recommended on this label.

**Field sprayer application:** **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

**Aerial application:** **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

**DO NOT** apply through irrigation equipment.

**DO NOT** apply through any type of ultra-low volume (ULV) spray system (less than 30 liters per hectare).

**Spray Buffer Zones:**

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop	Spray Buffer Zones (meters) Required for the Protection of:					
		Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial habitat	
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m		
Field Sprayer*	Cereals	1	0	1	0	1	
Aerial	Cereals	Fixed Wing	1	0	3	1	20
		Rotary Wing	1	0	1	1	20

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

## **GROUND APPLICATION**

### **Spray Equipment**

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate sprayer before use.

Use a pump with capacity to: (1) maintain at least 35-40 psi at nozzles, and (2) provide sufficient agitation in tank to keep mixture in suspension - this requires recirculation of 10% of tank volume per minute. Use a jet agitator or liquid sparge tube for agitation. **DO NOT** air sparge.

ALB AZOXYPROP is a suspoemulsion concentrate. It is suggested that screens be used to protect the pump and to prevent nozzles from clogging. Screens placed on suction side of pump should be 16-mesh or coarser. **DO NOT** place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations. For more information on spray equipment and calibration, consult sprayer manufacturers and provincial recommendations.

### **Spraying/Mixing**

ALB AZOXYPROP may be applied with all types of spray equipment commonly used for making ground application. Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, highly susceptible varieties, or when environmental conditions conducive to disease exist. Apply ALB AZOXYPROP in sufficient water volume for adequate coverage and canopy penetration.

Prepare no more spray mixture than is required for the immediate operation. Thoroughly clean spray equipment before using this product. Agitate the spray solution before and during application. Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

Add 1/2-2/3 of the required amount of water to the spray or mixing tank. With the agitator running, add ALB AZOXYPROP to the tank and continue agitation while adding the remainder of the water. Begin application of the spray solution after the ALB AZOXYPROP fungicide has completely dispersed into the mix water and maintain agitation until all of the mixture has been sprayed.

## **AERIAL APPLICATION**

### **Instruction for Aerial Application**

Apply only by fixed-wing or rotary aircraft equipment, which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

## **Use Precautions**

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

## **Operator Precautions**

**DO NOT** allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear long pants, a long-sleeved shirt, shoes and socks and chemical-resistant gloves during mixing/loading, application, clean-up and repair activities. In addition, wear protective eyewear (goggles or a face shield) during mixing and loading. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

## **Product-Specific Precautions**

Read and understand the entire label before opening this product. If you have questions, call the manufacturer, Albaugh, LLC at 1-800-247-8013 or obtain technical advice from the distributor or your provincial agricultural representative.

Application of this specific product must meet and/or conform to the following:

- Volume: Apply the recommended rate in a minimum spray volume of 45 litres of water per hectare.
- Avoid spraying when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

## **SPRAY DRIFT MANAGEMENT**

### **ATTENTION**

ALB AZOXYPROP is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

DO NOT spray ALB AZOXYPROP where spray drift may reach apple trees.

AVOID spraying when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc.

DO NOT use spray equipment which has been previously used to apply ALB AZOXYPROP to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

## USE DIRECTIONS

<b>CEREALS</b>	
<b>CROPS</b>	Barley
<b>DISEASE</b>	Barley net blotch ( <i>Pyrenophora teres</i> ), Barley scald ( <i>Rhynchosporium secalis</i> ), Barley leaf rust ( <i>Puccinia hordei</i> ), Septoria leaf spot ( <i>Septoria</i> sp.), Tan spot ( <i>Pyrenophora tritici-repentis</i> ), Stripe rust ( <i>Puccinia striiformis</i> )
<b>PRODUCT RATE (L/ha)</b>	0.53
<b>APPLICATION TIMING</b>	Apply once between stem elongation and half-head emergence (G.S. 29-55). Good spray coverage and canopy penetration are important for best results.
<b>CROPS</b>	Rye
<b>DISEASE</b>	Barley scald ( <i>Rhynchosporium secalis</i> ), Septoria leaf spot ( <i>Septoria</i> sp.), Tan spot ( <i>Pyrenophora tritici-repentis</i> )
<b>PRODUCT RATE (L/ha)</b>	0.53
<b>APPLICATION TIMING</b>	Apply once between stem elongation and half-head emergence (G.S. 29-55). Good spray coverage and canopy penetration are important for best results.
<b>CROPS</b>	Triticale
<b>DISEASE</b>	Septoria leaf spot ( <i>Septoria</i> sp.), Tan spot ( <i>Pyrenophora tritici-repentis</i> )
<b>PRODUCT RATE (L/ha)</b>	0.53
<b>APPLICATION TIMING</b>	Apply once between stem elongation and half-head emergence (G.S. 29-55). Good spray coverage and canopy penetration are important for best results.
<b>CROPS</b>	Wheat
<b>DISEASE</b>	Septoria leaf spot ( <i>Septoria</i> sp.), Tan spot ( <i>Pyrenophora tritici-repentis</i> ), Stripe rust ( <i>Puccinia striiformis</i> ), Wheat leaf rust ( <i>Puccinia triticina</i> )
<b>PRODUCT RATE (L/ha)</b>	0.53
<b>APPLICATION TIMING</b>	Apply once between stem elongation and half-head emergence (G.S. 29-55). Good spray coverage and canopy penetration are important for best results.
<b>CROPS</b>	Oats
<b>DISEASE</b>	Barley net blotch ( <i>Pyrenophora teres</i> ), Crown Rust ( <i>Puccinia coronata</i> var. <i>avenae</i> ), Septoria leaf spot ( <i>Septoria</i> sp.)
<b>PRODUCT RATE (L/ha)</b>	0.53
<b>APPLICATION TIMING</b>	Apply once between stem elongation and half-head emergence. Good spray coverage and canopy penetration are important for best results.
<p><b>Specific Use Restrictions:</b> DO NOT make more than one application per season. DO NOT apply within 45 days of harvest for grain and straw (45 day PHI). DO NOT harvest within 30 days for forage and hay (30 day PHI). Not all of these cereal crops have been tested for efficacy and phytotoxicity at the recommended label rates and ALB AZOXYPROP should be used at the discretion of the user. ALB AZOXYPROP may be applied by air or ground equipment.</p> <p><b>GROUND APPLICATION:</b> Apply specified rates in a minimum of 100 L of water per hectare.</p> <p><b>AERIAL APPLICATION:</b> Apply specified rates in a minimum of 45 L of water per hectare.</p>	

## RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that ALB AZOXYPROP fungicide contains both a Group 3 (propiconazole) and Group 11 (azoxystrobin) fungicides. ALB AZOXYPROP has two modes of action: (1) DMI (Demethylation Inhibitor of sterol biosynthesis) [Group 3], and (2) inhibitor of the Qo (quinone outside) site within the electron transport system (QoI) as well as disrupting membrane synthesis by blocking demethylation [Group 11]. Any fungal population may contain individuals naturally resistant to ALB AZOXYPROP and other Group 3 or Group 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

### To delay fungicide resistance:

- DO NOT exceed the total number of applications of ALB AZOXYPROP per season per crop as stated in label.
- Follow the specific crop recommendations that limit the total number of sprays on a crop and the required alternations with fungicides from other resistance management groups.
- In situations requiring multiple sprays, develop season long spray programs for Group 11 (QoI) fungicides. The program should meet the goal of no more than 1/3 of the total sprays per season, when a Group 11 fungicide is used as a solo product, or 1/2 the total sprays when a Group 11 fungicide is used in a mixture. Programs that include both solo Group 11 products and/or mixes containing Group 11 products should be no more than 1/2 the total sprays.
- Where possible, rotate the use of ALB AZOXYPROP or other Group 3 and 11 fungicides with different fungicide groups that control the same pathogens.
- DO NOT apply sequential treatments of ALB AZOXYPROP, or other fungicides in the same Fungicide Group, in a season. DO NOT apply at rates lower than recommended on the label.
- Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.
- Monitor treated fungal populations for sign of resistance development.
- If disease continues to progress after treatment with this product, DO NOT increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, to which pathogen resistance has not developed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance, contact a Albaugh, LLC company representative at 1-800-247-8013

### Application Limitation and Preharvest Interval (PHI) Summary

Crop	PHI Interval	Maximum number of applications per year
Grain & Straw: wheat, barley, oats, triticale, rye	45 days	1
Forage and Hay: wheat, barley, oats, triticale, rye	30 days	1