

2023-5833
2023-11-25

Insert (English)

Parapet Praventia Small Cat
Flea Adulticide/Larvicide/Ovicide

For cats 8 weeks and older weighing between 2.3 kg and 4 kg

Once-a-month topical flea treatment

Solution

READ THE LABEL AND INSERT BEFORE USING

FOR USE ONLY ON CATS

Do not use on other animals

KEEP OUT OF REACH OF CHILDREN



POISON

WARNING EYE IRRITANT

DOMESTIC

ACTIVE INGREDIENTS: Imidacloprid 9.1% and Pyriproxyfen 0.46%

REG. NO. 34932 P.C.P. ACT

Contents: One to Six 0.4 mL Tubes

Tevra Brands LLC
9100 F Street, Suite 200
Omaha, NE, USA
68127

Rolf C. Hagen Inc.
20500 Trans Canada Hwy
Baie d'Urfé, QC H9X 0A2
Telephone: 1-833-408-0768

[Contains Imidacloprid, Pyriproxyfen, the active ingredients found in Advantage II.] [Contains the same active ingredients as Advantage II.]

Parapet Praventia Small Cat kills 98-100% of the existing fleas on cats within 1 day after treatment. Protection against further flea infestation can last up to 4 weeks on cats. Retreatment at monthly intervals is required if re-infestation is continuous or if flea control is desired. Fleas, eggs, and larvae are killed following contact with skin debris from an Parapet Praventia Small Cat treated pet. Parapet Praventia Small Cat provides multi-stage flea control effectively breaking all flea life-cycle stages for quick and lasting control of flea populations.

Do not shampoo or immerse the animal in water immediately after treatment. The animal can be shampooed, immersed in water or exposed to rain 3 days after treatment. Parapet Praventia Small Cat retains its ovicidal and larvicidal efficacy for up to 4 weeks on cats following a shampoo treatment, swimming or exposure to rain. The adulticidal effect may be shortened by repeated shampooing, water immersion or exposure to rain. Reapplication on cats may be required after such activities.

The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with Parapet Praventia Small Cat rapidly kills fleas and may reduce the incidence of this condition. For effective flea control, treatment of the pet should be combined with sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, an insecticidal premise treatment may be required. A veterinarian should be consulted when pest infestation continues to be a problem on the pet.

PRECAUTIONS:

HUMANS: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed. Causes eye irritation. DO NOT get in eyes. Avoid contact with skin or clothing. Wear rubber gloves during application of this product. After handling or applying, wash hands (or any other skin that came in contact with the product), with soap and water. Avoid contact with pets until dry.

PETS: Do not use on cats under 8 weeks of age. Care should be taken to prevent exposure of other pets that may be in close physical contact with the treated animal. Sensitivities may occur after using ANY pesticide product for pets. Smaller animals are more likely to experience an adverse reaction. Observe carefully. If signs persist, or become more severe, consult a veterinarian immediately.

Side Effects: Monitor your pet after application. Side effects may include: skin irritation such as scratching or hair loss at the application site, or changes in behaviour such as agitation or lethargy. Gastrointestinal effects such as drooling, vomiting, or loss of appetite may also occur. If these or other side effects occur consult your veterinarian or Rolf C. Hagen Inc. at 1-833-408-0768.

FIRST AID:

HUMANS:

IF IN EYES, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

STORAGE: Store container at room temperature.

DISPOSAL: Wrap empty tube and cap in newspaper and dispose in household garbage.

DIRECTIONS FOR USE:

This product contains Imidacloprid and Pyriproxyfen. Do not apply another pest control product such as a shampoo, collar, or powder which contains Imidacloprid, or Pyriproxyfen to the treated animal after applying Parapet Praventia Small Cat.

Do NOT administer by mouth.

This product is packaged in single dosage tubes. This package contains one to six monthly 0.4 mL dosages. Do not apply a dosage higher than one 0.4 mL tube at one time. Treat your cat monthly in the manner described below. Do not use on other animals.

Cats

2023-5833
2023-11-25

Use only on cats 8 weeks of age and older weighing between 2.3 and 4 kg. Do not use on cats under 8 weeks of age. Do not retreat earlier than 4 weeks. Repeat every 4 weeks for continual pest control in situations where there is an ongoing risk of pest infestation.

How to Apply

1. Remove one applicator from the package.
2. Hold applicator tube in upright position. Cut cap off tube.
3. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to completely empty the contents directly on the skin as a single spot. Do not get this product in your pet's eyes or mouth. The product is bitter tasting and salivation may occur for a short time if the animal licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the animal to lick the product. Do not allow recently treated animals to groom each other.
4. Discard the empty tube as described in "DISPOSAL".

Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment. A veterinarian should be consulted when pest infestation continues to be a problem on the pet.

Until Parapet Praventia Small Cat dries, keep pet off furniture and away from hardwood floors.

Retain packaging information for future reference.

2023-5833
2023-11-25

Carton (English)

Parapet Praventia Small Cat

Flea Adulticide/Larvicide/Ovicide

For cats 8 weeks and older weighing between 2.3 kg and 4 kg

Once-A-Month Topical Flea Treatment

Solution

READ THE LABEL AND INSERT BEFORE USING

FOR USE ONLY ON CATS

Do not use on other animals

KEEP OUT OF REACH OF CHILDREN



POISON

WARNING EYE IRRITANT

DOMESTIC

ACTIVE INGREDIENTS: Imidacloprid 9.1% and Pyriproxyfen 0.46%

REG. NO. 34932 P.C.P. ACT

Contents: One to Six 0.4 mL Tubes

Rolf C. Hagen Inc.
20500 Trans Canada Hwy
Baie d'Urfé, QC H9X 0A2
Telephone: 1-833-408-0768

[Contains Imidacloprid, Pyriproxyfen, the active ingredients found in Advantage II.] [Contains the same active ingredients as Advantage II.]

Parapet Praventia Small Cat kills 98-100% of the existing fleas on cats within 1 day after treatment. Protection against further flea infestation can last up to 4 weeks on cats. Retreatment at monthly intervals is required if re-infestation is continuous or if flea control is desired. Fleas, eggs, and larvae are killed following contact with skin debris from an Parapet Praventia Small Cat treated pet. Parapet Praventia Small Cat provides multi-stage flea control effectively breaking all flea life-cycle stages for quick and lasting control of flea populations.

Do not shampoo or immerse the animal in water immediately after treatment. The animal can be shampooed, immersed in water or exposed to rain 3 days after treatment. Parapet Praventia Small Cat retains its ovicidal and larvicidal efficacy up to 4 weeks on cats following a shampoo treatment, swimming or exposure to rain. The adulticidal effect may be shortened by repeated shampooing, water immersion or exposure to rain. Reapplication on cats may be required after such activities.

PRECAUTIONS:

HUMANS: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed. Causes eye irritation. DO NOT get in eyes. Avoid contact with skin or clothing. Wear rubber gloves during application of this product. After handling or applying, wash hands (or any other skin that came in contact with the product), with soap and water. Avoid contact with pets until dry.

PETS: Do not use on cats under 8 weeks of age. Care should be taken to prevent exposure of other pets who may be in close physical contact with the treated animal. Sensitivities, may occur after using ANY pesticide product for pets. Smaller animals are more likely to experience an adverse reaction. Observe carefully. If signs persist, or become more severe, consult a veterinarian immediately.

| |
|--|
| <p>Side Effects: Monitor your pet after application. Side effects may include: skin irritation such as scratching or hair loss at the application site, or changes in behaviour such as agitation or lethargy. Gastrointestinal effects such as drooling, vomiting, or loss of appetite may also occur. If these or other side effects occur consult your veterinarian or Rolf C. Hagen Inc. at 1-833-408-0768.</p> |
|--|

FIRST AID:

HUMANS:

IF IN EYES, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting

2023-5833
2023-11-25

unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

STORAGE: Store container at room temperature.

DISPOSAL: Wrap empty tube and cap in newspaper and dispose in household garbage.

DIRECTIONS FOR USE

This product contains Imidacloprid and Pyriproxyfen. Do not apply another pest control product such as a shampoo, collar, or powder which contains Imidacloprid, or Pyriproxyfen to the treated animal after applying Parapet Praventia Small Cat.

2023-5833
2023-11-25

Tube (English)

Parapet Praventia Small Cat

READ THE LABEL AND INSERT BEFORE USING



POISON

WARNING EYE IRRITANT

DOMESTIC

EXTERNAL USE ONLY

ACTIVE INGREDIENTS: Imidacloprid 9.1% and Pyriproxyfen 0.46%

REG. NO. 34932 P.C.P. ACT

Rolf C. Hagen Inc.
20500 Trans Canada Hwy
Baie d'Urfé, QC H9X 0A2
Telephone: 1-833-408-0768

0.4 mL