

14-OCT-1998

AVAILABLE THROUGH VETERINARIANS

VET-KEM

OVITROL (TM) PLUS

Dual Action Flea & Tick Collar for Dogs

Kills adult fleas and ticks  
Kills flea eggs for 7 months

With PRECOR®  
Insect Growth Regulator (IGR)

DOMESTIC

GUARANTEE:

Tetrachlorvinphos ..... 14.55%  
(S)-Methoprene ..... 1.02%

REG. NO. 25667 P.C.P. ACT

READ THE LABEL BEFORE USING

CONTENTS: 1 COLLAR, NET WT. 33g

Wellmark International  
P.O. Box 20040  
Woodlawn Postal Outlet  
Guelph, Ontario N1H 8H6

distributed by  
Hoechst Roussel Vet  
240 Henderson Drive  
Regina, SK S4N 5P7

Vet-Kem® Ovitrol™ Plus Dual Action Flea and Tick Collar for Dogs will begin to kill fleas, ticks and flea eggs immediately. Its full protection against harder to kill ticks will be built up within a few days after being placed on the pet. The collar will continue to control fleas for 7 months by killing eggs laid on your pet. Under conditions where pets are

exposed to severe flea or tick infestations, it may be necessary to replace the collar more frequently. Wetting will not impair the collar's effectiveness or the pet's protection. If the pet goes swimming or is out in the rain, it is not necessary to remove the collar. The Vet-Kem® Ovitrol™ Plus Dual Action Flea and Tick Collar for Dogs may be worn with a regular collar. For use when fleas and ticks are a problem. If only ticks are problematic, a collar without the insect growth regulator should be used.

**DIRECTIONS:** Remove collar from package, unroll and stretch to activate insecticide generator. As the collar begins to work, a fine white powder will appear on the surface. Place the Vet-Kem® Ovitrol™ Plus Dual Action Flea and Tick Collar for Dogs around the dog's or puppy's neck, adjust for proper fit, and buckle in place. The collar must be worn loosely to allow for growth of the animal and to permit the collar to move around the neck. A properly fitted collar is one that, when fastened, will snugly slide over the dog's head. Leave 5 to 8 cm on the collar for extra adjustment. Cut off and dispose of extra length in empty pouch. Wash hands with soap and water after handling collar.

For effective flea and tick control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpet, furniture, (discard vacuum bag after use) and wash the pet's bedding, living quarters, and surrounding areas. If pest problems persist, an insecticidal premise treatment (e.g. Vet-Kem Siphotrol (TM) Premise Spray) may be required.

**PRECAUTIONS:** DO NOT ALLOW CHILDREN TO PLAY WITH COLLAR. Do not unroll collar until ready to use. This collar is intended for use only as an insecticide/IGR generator dog collar. Consult a veterinarian before treating pregnant or sick animals or animals under drug or other pesticide treatment. Do not use on nursing animals or animals under 8 weeks of age. Any irritation usually indicates the collar has been affixed too tightly. If irritation occurs, loosen the collar. If irritation continues, remove the collar from the animal and consult your veterinarian.

**FIRST AID:** If contact with collar causes irritation, or if ingested, consult physician or Poison Control Centre immediately.

**TOXICOLOGICAL INFORMATION:** Tetrachlorvinphos is a cholinesterase inhibitor. Atropine is antidotal.

DISPOSAL: Do not reuse empty pouch. Wrap in newspaper and discard expired collar, any excess collar and pouch with household garbage.

Hotline: 1-800-263-3532

Vet-Kem, Siphotrol and Precor are registered trademarks of Wellmark International

E.P.A. Est. No. 11715-TN-1

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))