

15-DEC-1998

COMMERCIAL

Net Contents 22.7 kg

Folpet 50-WP

Wettable Powder Fungicide for Use
in Water for Plant Disease Control

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

GUARANTEE: Folpet 50%

REGISTRATION NO. 07480 PEST CONTROL PRODUCTS ACT

ZENECA Agro, a business of ZENECA Corp.
250-3115 12th Street N.E.
Calgary, AB T2E 7J2

A93/A-2

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

* IRRITATING TO SKIN, EYES, NOSE AND THROAT. * Avoid Breathing Dust or Spray Mist. * Avoid Contact with Skin. * Wear Gloves. * Wash Hands Thoroughly After Use. * Do not contaminate food or feed. * This product is toxic to fish, do not contaminate any body of water by cleaning of equipment, disposal of wastes or application where run-off is likely to occur.

FIRST AID

If a known exposure occurs or is suspected, immediately initiate the recommended procedures below. If further treatment is required, contact a Poison Control Centre, a physician, or the nearest hospital.

If Swallowed: Immediately give several glasses of water and induce vomiting by gagging the victim with a finger or blunt object placed on the back of the victim's tongue. Give fluids until vomitus is clear. If victim is unconscious, do not induce vomiting or give anything by mouth.

If Inhaled: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

If in Eyes: Immediately flush eyes with large amounts of running water for at least 15 minutes. Hold eyelids apart during flushing to ensure rinsing of the entire surface of the eye and lids with water. Get medical attention.

If on Skin: Flush all affected areas with plenty of water for 15 minutes. Remove contaminated clothing and shoes. Seek medical attention if skin irritation occurs. Wash clothing before reuse.

EMERGENCY TELEPHONE NUMBER

All hours, 1-800-327-8633 (FASTMED) ONLY for health and environmental information.

STORAGE

Store in a cool dry well ventilated area away from foodstuffs and out of reach of children and animals. Protect from excessive heat. This product is not affected by freezing.

SPILL CLEANUP

Wear Appropriate protective equipment (gloves, glasses, apron) when attempting to clean up the spill. If the container is leaking, secure leak and place the container into a drum or heavy gauge plastic bag. Contact ZENECA Agro (See EMERGENCY NUMBER) for further information.

For spills and leaks; clean up as much of the product as possible with a broom and shovel. If available, use an absorbent such as DUSTBANE to clean the spill area. Shovel the spilled product and spend absorbent in sealable waste container.

On hard surfaces - sprinkle spill area with detergent and scrub in a small quantity of water with a coarse broom. Let stand for 10 minutes then absorb onto an inert material and shovel waste into a waste container.

On soil - remove the top 15 cm of soil in the spill area and replace with fresh soil. Dispose of all waste including scrub brush in accordance with provincial requirements. For more information on the disposal of waste and the clean up of spills, contact the provincial regulatory agency or the manufacturer.

DISPOSAL

Rinse the empty container thoroughly and add the rinsings to the spray mixture in the tank. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the provincial regulatory agency or the manufacturer.

DIRECTIONS FOR USE

GENERAL USE PRECAUTIONS

Use only for claims listed and only as specified on this label. Folpet 50-WP may be combined with most commonly used insecticides, adjuvants and fungicides. Fungicidal activity is reduced if combined with strongly alkaline materials. This material should not be used in combination with or closely following oil sprays.

Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for damages caused by drift from his property to that of others.

Consult local agricultural authorities for additional information, as the timing of applications needed will vary with the local conditions.

APPLICATION RATES

Ornamentals

Unless otherwise specified, dosage rates are in terms of Folpet 50-WP per 1,000 litres of water. Thorough coverage is required.

Asters, Phlox: Powdery Mildew - Use 2 kg. Make first application before disease appears and repeat at 7 to 10 day

intervals. Increase frequency during periods of rapid new growth and severe infection caused by rains or high humidity.

Azaleas (Cuttings): Damping Off - Use 1.5 to 2 kg. Soak cuttings for 15 to 30 minutes before planting in rooting media.

Carnations: Alternaria Leaf Spot - Use 2 kg. Make first application 2 weeks after planting and treat 3 to 4 times at bi-weekly intervals.

Chrysanthemums: Powdery Mildew, Septoria Leaf Spot - Use 2 kg. Make first application immediately after planting and repeat at 7 to 10 day intervals. Increase frequency during periods of rapid new growth and severe infection caused by rains or high humidity.

Iris: Didymellina Leaf Spot - Use 2 kg. Make first application when plants emerge and repeat at 7 to 10 day intervals.

Marigolds, Zinnias: Alternaria Leaf Spot - Use 2 kg. Make first application when plants emerge and repeat at 3 to 7 day intervals. Use the higher frequency during periods of rapid new growth and severe infection caused by rains or high humidity.

Poinsettias (Greenhouse): Pythium, Root Rot - Use 2 kg plus a spreader and use 600 ml of suspension per 20 centimetre pan. Make application at time of panning and repeat 10 days later.

Roses: Black Spot, Powdery Mildew - Use 2 kg. Make first application when first leaves unfold and repeat at 7 to 10 day intervals throughout the season. Do not use wetting agents on roses as injury may result.

Snapdragons: Anthracnose, Powdery Mildew - Use 2 kg. Make first application when plants emerge and repeat at 3 to 7 day intervals. Use the higher frequency during periods of rapid new growth and severe infection caused by rains or high humidity.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))