

97.06.19

RHONE-POULENC

A liquid flowable for control of  
DAMPING OFF and ROOT ROT  
caused by Rhizoctonia Solani,  
BLACKLEG and FLEABEETLE in CANOLA

FOR USE BY COMMERCIAL SEED COATERS ONLY

ROVRAL (R) CST  
CANOLA SEED TREATMENT

Contains Iprodione and Lindane

AGRICULTURAL

DANGER            POISON

READ THE LABEL BEFORE USING

ROLL DRUM OR STIR WELL BEFORE USING

GUARANTEE: Iprodione 167 G/L Lindane 500 g/L

REGISTRATION NO. 24972 PEST CONTROL PRODUCTS ACT

RHONE-POULENC CANADA INC.  
2000 ARGENTIA ROAD, PLAZA 3  
MISSISSAUGA, ON  
L5N 1V9

905-821-4450

In case of spills, poisoning or fire  
telephone emergency response number  
1-800-334-7577 (24 hours a day).

NET CONTENTS 100 LITRES

ROVRAL (R) is a registered trademark of Rhone-Poulenc Agrochimie,

Lyon - Authorized user Rhone-Poulenc Canada Inc.

DIRECTIONS FOR USE:

ROVRAL CST is a liquid dual purpose seed treatment for use on canola to control fleabeetles, damping off and root rot caused by *Rhizoctonia solani* and "seed borne" Blackleg on emerging seedlings.

Rate: 30 mL per kg of seed or 750 mL per 25 kg bag of seed.  
(The 100 L container treats 3333 kg of seed).

A colourant MUST be added to this ROVRAL CST to colour the seed, in accordance with the Pest Control Products Act and the Seeds Act regulations.

ROVRAL CST treated seed can be mixed with the granular insecticides \*Furadan CR-10 and \*Counter 5G. Refer to labels of both products for use recommendations and all safety precautions.

METHOD OF APPLICATION:

ROVRAL CST treated seed should not require drying after treatment and can be stored or bagged immediately. Add ROVRAL CST to the seed slowly and mix well for at least 3-5 minutes in treating equipment. Thorough seed coverage is required.

USE LIMITATIONS:

Do not use treated seed for feed, food or oil processing.

Do not store ROVRAL CST near feed or foodstuffs.

Do not contaminate feed or foodstuffs.

Do not contaminate domestic or irrigation waters.

A colourant MUST be added to this product to colour the seed in accordance with the Pest Control Products Act and the Seeds Act regulations.

Treated seed must be labelled with the date of treatment as follows:

This seed has been treated with ROVRAL CST containing Iprodione and lindane. Do not use for feed, food or oil processing.

Do not store treated seed for more than 6 months. Treated seed stored for periods in excess of 6 months may decrease in germination at a faster rate than untreated seed. Treated seed stored, for more than 6 months should be tested for germination before planting.

Thoroughly clean auger after handling treated seed before using same auger for handling commercial or feedgrains.

Leftover treated seed should not be stored near feed, or foodstuffs, or where there is a chance of it becoming mixed with untreated seed. Do not contaminate domestic or irrigation waters.

Store treated seed in cool, dry conditions.

\*Furadan CR-10 is a registered trademark of FMC Corporation. Counter 5G is a registered trademark of Cyanamid Canada Inc.

#### PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN, AVOID CONTACT WITH SKIN AND EYES. Wear protective gloves and coveralls when handling ROVRAL CST or treated seed. When handling ROVRAL CST and when treating seed or while augering or handling treated seed, work in a well ventilated area and wear a suitable respirator. Wash hands and exposed skin thoroughly after handling the concentrate and after application.

#### FIRST AID:

Remove person from contaminated area, remove contaminated clothing. Keep patient warm, comfortable, at rest.

Eye Contact: Irrigate with water for 15 minutes. Get medical attention.

Skin Contact: Wash exposed skin with soap and cold water.

Inhalation: If breathing stops, start artificial respiration.

Ingestion: If patient is conscious, wash out mouth. Induce vomiting by touching back of the throat with a

finger or blunt object. Do not give milk or oils.  
Get medical attention.

TOXICOLOGICAL INFORMATION:

Contains lindane, a chlorinated hydrocarbon which may be rapidly absorbed following oral ingestion or skin contact. Gastric lavage with saline purgative should be employed; oil purgatives are contraindicated. Symptomatic treatment is recommended, including an appropriate anti-convulsant therapeutic (eg. diazepam) if required.

STORAGE CONDITIONS:

Protect from frost.

DISPOSAL:

1. After use, return the container to the point of purchase or designated locations.
2. This container must only be refilled with ROVRAL CST. Do not reuse container for any other purpose.
3. Prior to refilling, inspect thoroughly for damage such as cracks, punctures, abrasions, and damaged or worn out threads on closure devices.
4. Check for leaks after refilling and before transportation. Do not refill or transport damaged or leaking containers.
5. If the container is not being refilled, return to point of purchase.
6. For information on the disposal of unused, unwanted product and the clean-up of spills contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto the buyer assumes the risk to persons or property arising from the use or handling

of this product and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))