

17-JAN-2007

29-JAN-2008 Notification - Change in Registrant

THROW BAG

**LATER'S
WARFARIN BAIT**

KILLS NORWAY (SEWER) RATS, ROOF RATS AND MICE.

DOMESTIC

REG. NO. 23554 P.C.P. ACT

GUARANTEE: Warfarin 0.025%

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

250 g

Notification Change

~~Spectrum Brands IP Inc.,~~
P.O. Box 21001,
Brantford, ON N3R 7W9

Sure-Gro IP Inc.

DIRECTIONS FOR USE: Place the bait in inaccessible locations or in protected bait stations in areas such as along walls, in corners, near rat holes and in concealed areas where rats and mice seek food, water or travel. Check daily to make sure that an uninterrupted supply of bait is available for at least 15 days, continue baiting until all signs of feeding have stopped. Replace bait which becomes damp or moldy. 1 m = 1 long pace (approx.)

HOW TO USE: For Rats: Place 250 g bait bags 6 to 9 m apart. **For Mice:** Place 250 g bait bags 2 to 3 m apart.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Do not open pre-measured place packs. Rubber gloves must be worn when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.**

USE LIMITATIONS: Bait **MUST** either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. **DO NOT** place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

ENVIRONMENTAL HAZARDS: Do not contaminate any body of water.

FIRST AID: If swallowed, give patient one or two glasses of water and **INDUCE VOMITING** by inserting a finger down the throat. Repeat until vomit fluid is clear. For all cases of human ingestion, immediately notify a physician or poison control centre. Never give anything by mouth to an unconscious person. **If in eyes,** flush with water for 15 minutes and get medical attention. **If on skin,** wash with soap and water. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

TOXICOLOGICAL INFORMATION: Vitamin K₁, in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE: Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

DISPOSAL: Do not reuse empty container. Dispose of empty container by placing in household garbage. **RODENT DISPOSAL:** Dispose of dead rodents in garbage or by burying.

17-JAN-2007

CARTON

**LATER'S
WARFARIN BAIT**

EFFECTIVE CONTROL FOR RATS AND MICE
IN AND AROUND THE HOUSE

KILLS NORWAY (SEWER) RATS, ROOF RATS AND MICE.

DOMESTIC

REG. NO. 23554 P.C.P. ACT

GUARANTEE: Warfarin 0.025%

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

500 g
2 x 250 g SEALED BAIT BAGS

Notification Change

~~Spectrum Brands IP Inc.,~~
P.O. Box 21001,
Brantford, ON N3R 7W9

Sure-Gro IP Inc.

LATER'S WARFARIN BAIT IS AN ANTICOAGULANT RODENTICIDE THAT REDUCES THE CLOTTING ABILITY OF BLOOD AND AFTER REPEATED FEEDINGS WILL CAUSE FATAL HEMORRHAGING IN RATS AND MICE. KILLS NORWAY RATS, ROOF RATS AND HOUSE MICE.

DIRECTIONS FOR USE: Place the bait in inaccessible locations or in protected bait stations in areas such as along walls, in corners, near rat holes and in concealed areas where rats and mice seek food, water or travel. Check daily to make sure that an uninterrupted supply of bait is available for at least 15 days, continue baiting until all signs of feeding have stopped. Replace bait which becomes damp or moldy.

1 m = 1 long pace (approx.)

HOW TO USE:

For Rats: Place 250 g bait bags 6 to 9 m apart.

For Mice: Place 250 g bait bags 2 to 3 m apart.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Do not open pre-measured place packs. Rubber gloves must be worn when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. KEEP AWAY FROM FEED AND FOODSTUFFS.

USE LIMITATIONS: Bait MUST either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

ENVIRONMENTAL HAZARDS: Do not contaminate any body of water.

FIRST AID: **If swallowed**, give patient one or two glasses of water and INDUCE VOMITING by inserting a finger down the throat. Repeat until vomit fluid is clear. For all cases of human ingestion, immediately notify a physician or poison control centre. Never give anything by mouth to an unconscious person. **If in eyes**, flush with water for 15 minutes and get medical attention. **If on skin**, wash with soap and water. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

TOXICOLOGICAL INFORMATION: Vitamin K₁, in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE: Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

DISPOSAL: Do not reuse empty container. Dispose of empty container by placing in household garbage. **RODENT DISPOSAL:** Dispose of dead rodents in garbage or by burying.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.