

95/2000

CO-OP

Captan 50% W.P.

WETTABLE POWDER FUNGICIDE

READ THE LABEL BEFORE USING

COMMERCIAL

GUARANTEE: CAPTAN 50%

REGISTRATION NO. 14171 PEST CONTROL PRODUCTS ACT

CONTROLS MANY COMMON DISEASES ON FRUITS, VEGETABLES, ORNAMENTALS AND TURF

PRECAUTIONS: KEEP OUT OF THE REACH OF CHILDREN.

Exposure to this product may cause skin and eye irritation. If safety precautions listed below are followed, exposure will be significantly reduced or eliminated. Keep away from heat, fire or sparks. Store away from feeds and foods. Never handle material with bare hands. Use rubber gloves; do not use leather or cloth gloves. Avoid spilling on skin or clothing. Do not breathe spray mist (dust). Wear a respirator and goggles while mixing, loading and spraying. When handling or applying, wear goggles and clothing that completely covers arms and legs to minimize exposure. Change contaminated clothing daily and wash thoroughly before re-use. Wash hands, face and arms thoroughly after handling captan and before drinking, eating or smoking. Shower after use or at the end of the work day. If an accident occurs, remove contaminated clothing and wash skin thoroughly with soap and water. Wash clothing and gloves before re-use. Always work out of the wind during mixing and loading operations. Keep unprotected persons out of operating area or vicinity where there may be drift.

2 kg

WARNING:

Exposure to Captan may produce long term health effects.
To minimize exposure, follow directions outlined on this label.

INTERPROVINCIAL COOPERATIVE LIMITED,
P.O. BOX 1050,
SASKATOON, SASKATCHEWAN
S7K 3M9

WINNIPEG MISSISSAUGA MONTREAL MONCTON

DIRECTIONS: Add Captan to water and mix thoroughly. Agitate solution during spraying operation to prevent Captan from settling out.

Crop	Disease	kg. product per 1000 litres of water	Comments
Apple	Scab Sooty Blotch Fly Speck	2 1-2	Apply in a regular protective schedule from green tip up to and including second cover spray. After second cover, cut rate in half if scab infection is at a low level. Consult provincial spray calendars for dosage and timing.
Peach	Scab Brown Rot	2-2.5	Apply in late pink, full bloom, petal fall, shuck fall, cover sprays and the day before picking.
Pear	Scab Sooty Blotch	2	For scab apply in the pre-bloom, petal fall and first cover sprays. For sooty blotch, apply in the second cover spray and two weeks later. Do not use on D'Anjou pears.

Plum	Black Knot Brown Rot	2-2.5	For Black knot, spray in green tip, pre-bloom and blossom sprays. For brown rot, apply in the pre-bloom, bloom, shuck and cover sprays.
Grape	Dead Arm	2	Apply when new shoots are 1-5 cm long and again when 10-15 cm long.
	Downy Mildew	2-2.5	Apply just before bloom, just after bloom in first cover and 2-3 weeks later.
Raspberry	Fruit Rot Spur Blight	2	Apply when blossoms first appear and repeat every 7-10 days until harvest.
Strawberry	Gray Mold Rot, Leaf Spot	4-6	Apply when the new growth starts in spring. Repeat at weekly intervals.
Cucumber	Anthracoese Scab	3-6	Apply at the first sign of disease. Repeat at intervals of 5-7 days. Use lower rates on young plants.
Potato (Seed Pieces)	Brown Eye Disease, Common Scab Rhizoctonia Seed Piece Decay	2.25-3	Dip in suspension then drain. Treat within 6 hours of cutting. Dry thoroughly unless planted within 1 day.
Tomato	Early and late Blight	4.5-7	Apply when plants are 5-10 cm high. Repeat at intervals of 5-10 days as long as disease threatens. Use high rates for older plants under disease pressure.
Turf (Ornamentals, Sport)	Brown Patch Damping-off, Leaf Spot Melting Out	150-220 g 100 m ²	Apply before infection occurs in early spring. Repeat 7-14 days throughout summer.
Rose	Black Spot	2-2.5	Apply at intervals of 7-10 days as necessary.

NOTE: Do not apply to apple, pear or grape within seven days of harvest. All other edible crops may be treated up to two days before harvest unless otherwise specified.

SYMPTOMS OF POISONING:

FIRST AID: In case of poisoning, call a physician immediately. IF ON SKIN, remove contaminated clothing and wash skin thoroughly with soap and water. IF IN EYES, flush with water for 5-10 minutes and obtain medical attention. IF SWALLOWED, give the patient one to two glasses of water, and cause vomiting by giving one dose of syrup of ipecac*. If the patient does not vomit within 20 minutes, give a second dose. If syrup of ipecac is not available, give the patient one to two glasses of water and cause vomiting by inserting a finger down the throat. Repeat with water until vomit fluid is clear. The patient should be lying down with the head below the level of the feet. DO NOT TRY TO CAUSE VOMITING IF THE PATIENT IS UNCONSCIOUS OR IN A CONVULSIVE STATE. *(15 mL)

TOXICOLOGICAL INFORMATION:

Decontamination and Disposal.

1. Empty bag thoroughly into spray tank.
2. Make the empty bag unsuitable for further use.
3. Dispose of the bag in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted product, and cleanup of spills, contact the regional office of Conservation and Protection, Environment Canada.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the

information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))