

Just One Bite® One-Timer™ Rodenticide

Single-Use Bait Dispenser for Mice

DOMESTIC

GUARANTEE:

Bromadiolone.....0.005%

REG. NO. **27338** PCP ACT

WARNING



POISON

**KEEP OUT OF REACH OF CHILDREN
READ THE LABEL BEFORE USING**

NET CONTENTS: 43 g

Farnam Companies, Inc.
301 West Osborn Road
Phoenix, AZ 85013

Canadian Resident Agent:
Farnam (Canada) Marketing, Inc.
#700 10104-103 Avenue
Edmonton AB T5J 0H8

(602) 285-1660 M-F
7:00 a.m.-3:45 p.m., MST

DIRECTIONS FOR USE:

Just One Bite® One Timer™ Rodenticide is a self-contained disposable bait station dispenser for the control of House mice in and around buildings. One-Timer Rodenticide dispensers must be maintained in tamper-resistant bait stations or in locations not accessible to children, pets, domestic animals or wildlife. DO NOT place bait in areas where food or feed is stored or where there is a possibility of contaminating food or surfaces that may come in contact with food. Do not use in sewers. This is not a tamper resistant bait station. Place bait in locations where mice travel and will most likely find and consume the bait. Placement along walls, in corners, by gnawed openings, in and around burrows, and in concealed locations where mice feed is desirable. Maintain a steady supply of fresh bait.

Application Directions:

Place 1 bait station at intervals of 2-4 m within the treatment area. After a unit is placed for use, make the bait available to mice by holding the base of the unit with one hand and gently lifting and turning the top portion in a counterclockwise direction. Stop when resistance to further turning is encountered (after about one quarter of a revolution). Do not attempt to open the unit further. Do not open units except when they are placed ready for use.

If a bait placement is not disturbed after several days, move the placement approximately 1 meter. Two bait stations per placement may be necessary at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease. If reinfestation occurs, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations as necessary. To prevent reinfestation, eliminate food, water and harbourage as much as possible.

Securing the Bait Station: If you desire to anchor the bait station dispenser to a smooth clean surface, remove the cover sheet from the tape anchor located on the underside of the unit. If you wish to anchor the dispenser to a surface which will not allow the tape anchor to adhere, you may remove the tape anchor and place the dispenser over a nail or other such post anchored in the surface and protruding above the surface approximately 2.5 cm. in order to inhibit lateral movement of the unit by rodents. For more secure anchorage, a nail may be driven through the center post of the bait dispenser and into a wood or similar surface.

Note: Close units before removing them for disposal. Replace units which are emptied by mice or if bait becomes contaminated or spoiled.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN, PETS, DOMESTIC ANIMALS AND WILDLIFE. THIS IS NOT A TAMPER RESISTANT BAIT STATION. Keep away from feed or foodstuffs. Use gloves when handling product. Do not get in eyes, on skin, or on clothing. Wash arms, hands, and face with soap and water after handling and before eating or smoking. Harmful or fatal if swallowed or absorbed through the skin. This material may reduce the clotting ability of the blood and cause bleeding.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

FIRST AID INSTRUCTIONS:

If in eyes, flush with water for 15 minutes. If irritation persists, seek medical attention. **If on skin**, wash affected area thoroughly with soap and water. **If swallowed**, give patient one or two glasses of water and induce vomiting by touching back of throat with a finger or blunt object; repeat until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious person. Seek medical attention or contact Poison Control Centre **IMMEDIATELY**.

Take container, label or product name and PCP Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Vitamin K₁ in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE:

Store bait in original, tightly sealed container. Store in a cool, dry place. Do not contaminate water, food or feed by storage or disposal.

DISPOSAL: Do not reuse empty container. Wrap empty container and discard with household garbage.

RODENT DISPOSAL:

Dispose of dead rodents in garbage or by burying.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))