

transfer to TPP

96.02.07

SIPCO INDUSTRIAL PRODUCTS LTD.

S.I.P. (TM)

SIP-TOL (TM)

DETERGENT DISINFECTANT  
FUNGICIDE AND VIRUCIDE  
CONCENTRATE

GUARANTEE:

n-alkyl (5% C12 60% C14 30% C16 5% C18)	
dimethyl benzyl ammonium chloride	4.5%
n-alkyl (68% C12 32% C14)	
dimethyl ethyl benzyl ammonium chloride	4.5%

COMMERCIAL

READ THE LABEL BEFORE USING

DIN 02168847

KEEP OUT OF REACH OF CHILDREN

CAUTION            CORROSIVE

REGISTRATION NUMBER 24374  
PEST CONTROL PRODUCTS ACT

Net Contents    4 Litres

SIPCO INDUSTRIAL PRODUCTS LTD.  
7236 WILSON AVE.  
DELTA, B.C.  
V4G 1E5

MADE IN CANADA

FOR HOSPITAL AND INSTITUTIONAL USE:  
DIRECTIONS: SIPTOL, is a phosphate-free formulation designed

to provide effective cleaning, deodorizing, sanitization, and disinfection specifically for hospitals, nursing homes, schools, food processing plants, food service establishments and other institutions where housekeeping is of prime importance in controlling the hazard of cross-contamination. SIPTOL deodorizes those areas which generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, toilet bowls, and other areas which are prone to odours caused by microorganisms.

**DISINFECTION:** For disinfection of hard non porous surfaces (walls, floors, etc.) add 8 mL cleaner to 1 litre of water. Allow solution to wet all surfaces thoroughly and remain wet for 10 minutes. To disinfect toilet bowls, flush toilet, add 38 mL of cleaner directly to bowl water. Swab completely making sure to get under rim. Let stand for 10 minutes, flush. For heavily soiled surfaces a pre-cleaning step is required and recommended. Prepare a fresh solution for each use.

**HOSPITAL DISINFECTION:** To disinfect walls, floors, table tops and counters in operating or examining rooms in Hospitals, Nursing Homes add 8 mL cleaner in 1 litre of water. To disinfect toilet bowls, flush toilet, add 38 mL of cleaner directly to bowl water. Swab completely making sure to get rim. Let stand for 10 minutes, flush. For heavily soiled surfaces a pre-cleaning step is required and recommended. At this dilution this product exhibits effective disinfectant activity against: *Pseudomonas aeruginosa* PRD-10, *Escherichia coli*, *Salmonella choleraesuis*, *Staphylococcus aureus*.

**SANITIZATION:** To sanitize previously cleaned and rinsed non-porous food contact surfaces, prepare a 200 ppm active QUAT solution of 2.2 mL cleaner in 1 litre of water. To sanitize immobile items such as chopping blocks and counter tops, flood the area with a 200 ppm active Quat or apply with a cloth or sponge making sure to wet all surfaces completely for at least 60 seconds. Let air dry. Prepare a fresh solution for each use. To sanitize mobile items such as drinking glasses and eating utensils, immerse in a 200 ppm active Quat for at least 60 seconds making sure to immerse completely. Remove and let air dry. Prepare a fresh solution daily or more frequently as soil is apparent. When used for sanitization of previously cleaned food equipment or food contact items, limit active quaternaries to 200 ppm.

**MILDEWSTAT:** To control mold and mildew on pre-cleaned, hard, non-porous surfaces, add 80 mL of cleaner to 10 litres of

water. Wet surfaces completely. Let air dry. Repeat application at weekly intervals or when mildew growth reappears. Prepare a fresh solution for each use. At 80 mL per 10 litres of water, this product is also fungicidal against the pathogenic fungi, *Trichophyton mentagrophytes*.

**VIRUCIDAL ACTIVITY:** When used on environmental hard surfaces at 80 mL cleaner in 10 litres of solution this product exhibits effective virucidal activity against Influenza A-2 Asian, Herpes Simplex, Adeno virus type 2 and Vaccinia virus.

Efficacy tests have demonstrated that this product is an effective bactericide, fungicide and virucide in the presence of organic soil (5% blood serum). This cleaner is an effective sanitizer when diluted with water up to 250 ppm hardness (CaCO<sub>3</sub>).

**PRECAUTION:** Keep out of reach of children. Causes eye and skin damage. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Wear glasses or face shield and rubber gloves when handling. Avoid contamination of food. When used on surfaces coming in direct contact with food, rinse thoroughly with potable water.

**FIRST AID INSTRUCTIONS:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, contact a physician. Remove and wash all contaminated clothing before re-use. If swallowed, drink milk, egg whites, gelatin solution or if none of these are available, drink large quantities of water. Call a physician immediately.

**NOTICE TO USER:** This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

**TOXICOLOGICAL INFORMATION:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

**DISPOSAL:**

1. Rinse the emptied container thoroughly and add the rinsings to treatment site.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.

3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills contact the Provincial Regulatory Agency or the Manufacturer.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))