96.04.17

CFPI

ALLIZOL

INDUSTRIAL

SOLUBLE POWDER

COMMERCIAL/INDUSTRIAL

FOR WEED CONTROL IN INDUSTRIAL AND OTHER NON-CROPPED AREAS

CAUTION * POISON

READ THE LABEL BEFORE USING

REGISTRATION NO. 15999 PEST CONTROL PRODUCTS ACT

CFPI, 92233 Gennevilliers, France c/o 150-251 Midpark Blvd. S.E. Calgary, Alberta T2X 1S3

NET CONTENTS 50 KG

DIRECTIONS FOR USE

FOR WEEDS

Apply ALLIZOL any time weeds are green and actively growing. Weeds treated after they have developed strong stems may remain standing for the rest of the season, even though they are dead. These standing weeds are unsightly and a fire hazard. Therefore, for most effective control of tall weeds, cut and remove the weeds if possible, then spray the regrowth. For best results, treat when soil moisture is adequate for rapid weed growth.

Use enough water to thoroughly wet weed foliage - 550 to 1100 L per ha. In highway spraying, a strip 2.5 m wide and 4 km long is

about 1 ha. Be sure ALLIZOL is completely dissolved before spraying. Do not use for selective weed control in turf or edible crops. Some of the Weeds Controlled: Alfalfa Blue grasses Canada Thistle Carpetweed Catch fly Chickweed, common Chrysanthemum weed Cocklebur Crab grass Couch grass Dandelion Downy brome grass Foxtail (wild millet) Goose grass Groundsel Hemp nettle Jimson weed Knotweed Kochia Lambsquarters Mustard Nightshade Piqweed Plantain Poison Ivy Purslane Ragweed Shepherd's purse Smartweeds Sow thistle Stinkweed Velvet grass White cockle Wild Buckwheat Wild oats Witchgrass Yellow rocket Amounts per hectare: ALLIZOL: 2.25KG WATER: 1000L

Control or suppress annual grasses and broadleaf weeds where residual control not required. Spray when most weeds are 8 to 10 cm tall.

Amounts per hectare: ALLIZOL: 4.25KG WATER: 1000L Control annual grasses, annual and perennial broadleaf weeds, suppress perennial grasses. Spray when most weeds are 15 to 25 cm tall.

Cattails: Treat after catkins are fully formed until frost.

Amounts per hectare:

 ground application: aerial application: 	ALLIZOL:	8.5-10.5KG	WATER:	3000L
	ALLIZOL:	10.5KG	WATER:	150L
Amounts per hectare:	ALLIZOL:	2.25KG	WATER:	1000L

Poison ivy, poison oak. Treat after leaves are fully developed in the spring until leaves start to turn color in the fall.

FOR WOODY PLANTS

Use Allizol in a right-of-way maintenance program as a special tool for cleaning up spot infestations of the species listed. Most of these are killed to ground line by foliage sprays of 2,4-D but resprout badly the next year. ALLIZOL is a good translocator and kills most of the roots as well as the tops.

Allizol is specific on the species listed. It should not be expected to kill other species. Use the standard foliage stem method for high volume sprays along fencerows, highways and utility rights-of-way. Spray all foliage and stems of plants from the time foliage is fully developed until plants begin to go dormant. For effective kills, all leaves, stems and suckers must be thoroughly wet to ground line.

AMOUNTS TO USE

Species	Kilograms Allizol per 1000L WATER
Salmonberry	3.75
Western dewberry (blackberry)	3.75
Bigleaf maple	3.75
Honeysuckle	3.75
Locust	2.75 to 3.75
Ash	3.75 to 5.75
Sumac	2.75 to 3.75

Add 250 mL of spreader sticker per 1000 L water.

PRECAUTIONS

Harmful if swallowed. Avoid contact with eyes, skin, and clothing. Wear a respirator during mixing and loading of this product.

Wear lightweight, waterproof clothing (jumpsuit or coverall and boots or shoes, gloves, and a wide-brimmed plastic hard hat) when applying this product to control dense, tall vegetation. Workers applying this product in other situations must wear lightweight gloves and boots (or shoes).

Avoid spray-drift as this product may injure beans, peas, tomatoes, some ornamentals, etc. (Coarse sprays are less likely to drift). Do not use same sprayer for other purposes unless thoroughly cleaned. To clean sprayer, rinse all parts of sprayer with water several times.

Do not store near fertilizers, seeds, insecticides or fungicides or use in a greenhouse. Do not contaminate water used for irrigation or domestic purposes.

FIRST AID:

If swallowed: Induce vomiting immediately. Get medical attention or contact a poison control centre.

If on skin: Wash thoroughly with soap and water. Remove contaminated clothing and wash before re-use.

If in eyes: Flush eyes with plenty of water and get medical attention or contact a poison control centre.

NOTE TO PHYSICIAN: This product may affect thyroid function. Overexposure can be determined by monitoring thyroid function. If worker is under treatment for hypo- or hyperthyroidism, he should be removed from spray crew until the euthyroid state is achieved.

DISPOSAL:

- 1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture.
- 2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
- 3. Make the empty container unsuitable for further use.
- 4. Dispose of the container in accordance with provincial

requirements.

5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the manufacturer.

NOTICE TO USER - This control product is to be used only in accordance with the directions o this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER - Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the Buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))