



NEGUVON®

POUR-ON
CATTLE INSECTICIDE

FOR CONTROL OF CATTLE GRUBS AND LICE

AGRICULTURAL

GUARANTEE: trichlorfon.....8%

This product contains the potentially toxic formulant dipropylene glycol monomethyl ether at 89.8%.

REGISTRATION NO. 9419 PEST CONTROL PRODUCTS ACT

DANGER POISON

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

PROTECT FROM EXTREME HEAT

NET CONTENTS: 4 LITRES

Bayer Inc.
77 Belfield Road
Toronto, Ontario, Canada M9W 1G6

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

DIRECTIONS FOR USE:

Neguvon Pour-On Cattle Insecticide is a ready-to-use preparation that requires no further dilution. A single application placed on the backline of an animal gives control of cattle grubs and reduction of lice populations.

AMOUNT TO USE:

Apply Neguvon Pour-On solution at the rate of 32.5 mL per 100 kilograms of body weight. Measure the correct amount to use in relation to the animal's weight.

THE CONTENTS OF THIS CONTAINER WILL TREAT:

Weight of Animals (kilograms)	100	200	300	400	500
Number of Animals	123	62	41	31	25

HOW TO APPLY:

Pour the correct amount of solution uniformly along the animal's back. The Neguvon solution may be dipped from any clean bucket or pail using the graduated applicator supplied. Do not subsequently use dip container in any connection with feed, food or drinking water.

WHEN TO APPLY TO BEEF AND NON-LACTATING DAIRY CATTLE FOR GRUB CONTROL:

The most beneficial effects of treatment with Neguvon Pour-On will be achieved by treating as soon as possible after activity of heel fly has ceased and preferably at least 6 weeks prior to expected appearance of grubs in the back.

REDUCTION OF LICE POPULATIONS:

A backline treatment for grub control also provides significant reduction of lice populations under most conditions. Heavily or chronically infested animals should be treated in the fall with an overall spray prepared from other suitable louse control insecticides.

APPLICATION LIMITATIONS:

Do not apply during the months of December, January, or February when this type of treatment may result in toxic side reactions such as salivation, bloat, hindquarter stiffness, or neuro muscular impairment. Should these symptoms occur, consult a veterinarian.

If treatment cannot be made before the end of November, treatment during the winter months of December, January, and February should be made only under the supervision of a veterinarian.

Do not treat lactating dairy cattle, animals which are less than 3 months old, sick or convalescent, or livestock in a stressed condition.

Do not apply to non-lactating dairy cattle within 7 days of freshening. If freshening should occur within 7 days of treatment, do not use milk as human food for the balance of the 7-day interval.

Do not treat animals for 10 days before or after shipping, weaning, or dehorning; or after exposure to contagious or infectious diseases.

Neguvon (trichlorfon) is a cholinesterase inhibitor. Do not use this product on animals at the same time as, or within a few days before or after, treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals.

Do not treat cattle within 21 days of slaughter.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN

Avoid contact with solution. Wear gloves while handling. May be poisonous if swallowed. Harmful if absorbed through the skin. Do not get in eyes or on skin. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use.

Do not contaminate feed, food, or water.

Do not re-use empty container.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DISPOSAL:

1. Follow provincial instructions for any required cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

FIRST AID:

In case of poisoning call a physician or Poison Control Centre immediately. **If on skin**, remove contaminated clothing and wash skin thoroughly with soap and water. **If in eyes**, flush with water for 5-10 minutes and obtain medical attention. **If swallowed**, do not induce vomiting but rush patient to nearest hospital or doctor's office.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Trichlorfon depresses cholinesterase. Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not give morphine.

<p>NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.</p>
--

305230

®Neguvon, Bayer and Bayer Cross are registered trademarks of Bayer



NEGUVON®

POUR-ON
CATTLE INSECTICIDE

FOR CONTROL OF CATTLE GRUBS AND LICE

AGRICULTURAL

GUARANTEE: trichlorfon.....8%

This product contains the potentially toxic formulant dipropylene glycol monomethyl ether at 89.8%.

REGISTRATION NO. 9419 PEST CONTROL PRODUCTS ACT

DANGER POISON

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

PROTECT FROM EXTREME HEAT

NET CONTENTS: 8 LITRES

Bayer Inc.
77 Belfield Road
Toronto, Ontario, Canada M9W 1G6

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

DIRECTIONS FOR USE:

Neguvon Pour-On Cattle Insecticide is a ready-to-use preparation that requires no further dilution. A single application placed on the backline of an animal gives control of cattle grubs and reduction of lice populations.

AMOUNT TO USE:

Apply Neguvon Pour-On solution at the rate of 32.5 mL per 100 kilograms of body weight. Measure the correct amount to use in relation to the animal's weight.

THE CONTENTS OF THIS CONTAINER WILL TREAT:

Weight of Animals (kilograms)	100	200	300	400	500
Number of Animals	246	124	82	62	50

HOW TO APPLY:

Pour the correct amount of solution uniformly along the animal's back. The Neguvon solution may be dipped from any clean bucket or pail using the graduated applicator supplied. Do not subsequently use dip container in any connection with feed, food or drinking water.

WHEN TO APPLY TO BEEF AND NON-LACTATING DAIRY CATTLE FOR GRUB CONTROL:

The most beneficial effects of treatment with Neguvon Pour-On will be achieved by treating as soon as possible after activity of heel fly has ceased and preferably at least 6 weeks prior to expected appearance of grubs in the back.

REDUCTION OF LICE POPULATIONS:

A backline treatment for grub control also provides significant reduction of lice populations under most conditions. Heavily or chronically infested animals should be treated in the fall with an overall spray prepared from other suitable louse control insecticides.

APPLICATION LIMITATIONS:

Do not apply during the months of December, January, or February when this type of treatment may result in toxic side reactions such as salivation, bloat, hindquarter stiffness, or neuro muscular impairment. Should these symptoms occur, consult a veterinarian.

If treatment cannot be made before the end of November, treatment during the winter months of December, January, and February should be made only under the supervision of a veterinarian.

Do not treat lactating dairy cattle, animals which are less than 3 months old, sick or convalescent, or livestock in a stressed condition.

Do not apply to non-lactating dairy cattle within 7 days of freshening. If freshening should occur within 7 days of treatment, do not use milk as human food for the balance of the 7-day interval.

Do not treat animals for 10 days before or after shipping, weaning, or dehorning; or after exposure to contagious or infectious diseases.

Neguvon (trichlorfon) is a cholinesterase inhibitor. Do not use this product on animals at the same time as, or within a few days before or after, treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals.

Do not treat cattle within 21 days of slaughter.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN

Avoid contact with solution. Wear gloves while handling. May be poisonous if swallowed. Harmful if absorbed through the skin. Do not get in eyes or on skin. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use.

Do not contaminate feed, food, or water.

Do not re-use empty container.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DISPOSAL:

1. Follow provincial instructions for any required cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

FIRST AID:

In case of poisoning call a physician or Poison Control Centre immediately. **If on skin**, remove contaminated clothing and wash skin thoroughly with soap and water. **If in eyes**, flush with water for 5-10 minutes and obtain medical attention. **If swallowed**, do not induce vomiting but rush patient to nearest hospital or doctor's office.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Trichlorfon depresses cholinesterase. Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not give morphine.

<p>NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.</p>
--

305230

®Neguvon, Bayer and Bayer Cross are registered trademarks of Bayer.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.