

2000/2005

AGRICULTURAL

20 kg

AMBUSH® 1% GRANULAR

Insecticide

READ LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

GUARANTEE: permethrin 1%

REGISTRATION NO. 17673 PEST CONTROL PRODUCTS ACT

ZENECA Agro, a business of ZENECA Corp.
#250, 3115 - 12th Street NE
Calgary, Alberta T2E 7J2

A93/A-1

PRECAUTIONS

This product is toxic to fish. Do not contaminate ponds, lakes, streams or rivers. **DO NOT APPLY BY AIRCRAFT.** Do not apply within 15 meters of fishery productive bodies of water or waterfowl habitats.

FIRST AID

If in contact with eyes, wash with plenty of clean water for at least 15 minutes and see physician. **If on skin**, wash with soap and water. **If swallowed**, induce vomiting and see physician.

EMERGENCY TELEPHONE NUMBER

All hours, 1-800-327-8633 **ONLY** for health and environmental information.

DISPOSAL

Thoroughly empty the contents of the container into the application device. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Environmental Protection, Environment Canada.

NOTICE TO USER

This control product to be used only in accordance with the directions on this label. It is an offence under the Pest Control Product Act to use a control product under unsafe conditions.

DIRECTIONS

Ground Application Only

SWEET AND FIELD CORN: European Corn Borer - Apply using granular applicators on high clearance equipment so granules fall into whorl.

Application Rates

Rate	90 g/100 m row					
Row Spacing	76 cm	81 cm	86 cm	91 cm	97 cm	102 cm
kg/ha	11.7	11.1	10.6	10.0	9.4	8.8

For first generation borers, apply no later than when the first feeding is seen on the foliage or apply according to the adult flight monitoring data provided by the local agricultural authority. For sweet corn, repeat in 5 to 10 days to ensure adequate protection. In some areas, protection from second generation borers may be necessary. Apply before tassels begin to appear in the whorls or according to the monitoring data provided by the local agricultural authority. For sweet corn, a repetition of above treatment may be required in 5 to 10 days for continued protection. Once larvae enter the stalk, control is not possible. Late season sweet corn ear protection may not be adequate in two generation areas.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out in the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

AMBUSH is a registered trademark of Imperial Chemical Industries PLC, England,

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))