

09-JAN-1996

DOMESTIC

1 LITRE

C-I-L

MOTHER EARTH(R)

INSECTICIDE - FUNGICIDE

FOR FRUITS, ROSES AND OTHER ORNAMENTAL PLANTS

LIQUID LIME SULPHUR

CONCENTRATED

FOR BOTH DORMANT AND SUMMER SEASON USE

READ THE LABEL BEFORE USING

GUARANTEE: SULPHIDE SULPHUR 23%

REG. NO. 7386.05 P.C.P. ACT

Made in CANADA

NU-GRO CORPORATION  
Woodstock, Ontario N4S 8P6  
Laval, Quebec H7S 1G3  
Calgary, Alberta T2H 2S1

DIRECTIONS FOR USE:

SHAKE WELL BEFORE USING: Lime Sulphur should mainly be used as a DORMANT SPRAY to clean up overwintering insects and disease infections. During the summer further applications with Lime Sulphur or C-I-L's Insecticides or Fungicides are necessary to keep plants free from pests. Lime Sulphur may only be mixed with C-I-L Dormant Oil, not with any other products. Do not apply during or shortly before temperature reaches 27°C or higher. Shake spray solution while spraying. Lime Sulphur stains painted wood, brick and stone. Prevent from freezing.

FRUIT TREES: As Dormant Spray, apply before buds break, to control Blister and Rust Mites, Scale insects, Pear Scale, Red Mite eggs, Aphid eggs, Twig Borer, Plum Black Knot and Peach Leaf Curl, mix 100 mL in 1 Litre of water (1/2 of this one Litre container in 5 Litres of water).

APPLE AND PEAR: During summer use 15 mL (1 tablespoonful) per Litre of water just before and again after blossoming to control powdery mildew, apple and pear scab; repeat 10 and 20 days after end of blossoming. Do not apply on plum in leaf and never on apricot. The apple variety "Delicious" may be injured by summer spraying with Lime Sulphur.

RASPBERRY, BLACKBERRY AND LOGANBERRY: To control rust, anthracnose, cane and spur blight, apply 100 mL in 1 Litre of water when buds break.

CURRENT, GOOSEBERRY: To control mildew, rust and scale insects apply 100 mL in 1 Litre of water just before buds break. Apply 60 mL (4 tablespoonfuls) in 3 Litres of water just before blossoms open and repeat weekly after bloom until fruit colours. May be used up to 1 day before harvest.

ROSES: To control Rose scale, canker and black spot make dormant application before buds break, using 100 mL in 1 Litre of water. For black spot and mildew during summer use 5 mL (1 teaspoonful) in 3 Litres of water every one to two weeks.

TREES AND SHRUBS: As Dormant Spray to control scale insects on bark and needles, also Maple gall mite, use 100 mL in 1 Litre of water; for Spruce gall aphid use 30 mL (2 tablespoonfuls) in 1 Litre of water.

COMBINATION DORMANT SPRAY: Fruit trees, roses and deciduous ornamentals. For overwintering insects, eggs, oyster shell scale, San José scale, rose scale, black scale as well as overwintering fungus disease: Mix 120 mL (8 tablespoonfuls) C-I-L's Lime Sulphur and 60 mL (4 tablespoonfuls) C-I-L's Dormant Oil in 3 Litres of water.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes and skin. Wash after use. Avoid staining clothing and skin. Dispose of empty container with regular garbage.

FIRST AID: If contact with eyes, rinse with plenty of water for 15 minutes and get medical attention. If contact with skin, wash with water and apply vinegar. If swallowed induce vomiting and obtain medical attention.

C-I-L is a trademark of ICI Canada Inc.  
NU-GRO Corporation is a Registered User.  
Mother Earth is a registered trademark of NU-GRO Corporation.

82401 995

GUARANTEE: Sulphide Sulphur

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))