

93.03.29

TERMINATOR

WARFARIN RODENTICIDE BAIT

COMMERCIAL

READ THE LABEL BEFORE USING

REGISTRATION NO. 21875

PCP ACT

NET CONTENTS:

340 g

GUARANTEE:

WARFARIN 0.025%

TERMINATOR BAIT PROCESSORS

Box 560, McLaughlin, AB T0B 2Y0

DIRECTIONS FOR USE:

This is a prepared bait registered in Canada for the control of the house mouse, roof rat and Norway rat for use in dwellings, farm buildings, food service establishments, granary bins (empty), processing plants (feed, food, nonfood), storage areas (feed, food, nonfood).

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood which, upon repeated feeding, causes fatal hemorrhaging in rats and mice.

Place bait in rodent tunnels, runways, along walls, in corners, near rat holes and areas where rats feed, drink or travel. Set bait out at the rate of 50 to 340 grams (1 bag) per station for rats. Set out piles of 15-50 grams of bait every 2-3 meters for

mice.

NOTE: If bait molds, decomposes, or becomes damp or sour, replace with fresh material. Maintain an uninterrupted supply of bait in the stations for no less than 15 days or until feeding ceases. Bait can be continued indefinitely if rat and mice infestation is a problem.

In meat packing, food processing plants and kitchens, use only when plant is not in operation. Place bait stations on the floor only; do not place in overhead locations. Remove or cover all food, packaging materials and utensils before placing baits. Remove all baits and dead rodents before reusing plant for food processing. If any equipment is accidentally contaminated, all equipment should be thoroughly washed.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN AND PETS. Place bait in locations which are not accessible to children, pets and domestic animals, or in tamper proof bait boxes. If taken in large or repeated doses by humans or animals, it may reduce the clotting ability of the blood and cause internal bleeding. Do not contaminate food or feed. Do not contaminate any body of water. Bury or burn bodies of the dead rodents.

FIRST AID:

If swallowed, give patient 1-2 glasses of water. To cause vomiting, insert a finger down the throat. Repeat until vomit fluid is clear. Call a physician.

TOXICOLOGICAL INFORMATION:

Vitamin K1 in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

STORAGE: Store only in original container in a dry place

inaccessible to children and pets.

DISPOSAL:

1. Dispose of stale bait and dead rodents in accordance with local requirements.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted product and clean up of spills, contact the regional office of Conservation and Protection, Environment Canada.

NOTICE TO USERS:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))