

93.08.24

DITRAC(R)

REUSABLE REUTILISABLE
PREBAITED PRE-APPATE

MOUSE BAIT STATION
STATION D'APPATS POUR SOURIS

KILLS MICE TUE LES SOURIS

REG.NO. 22136 P.C.P. ACT
NO.D'ENR. 22136 L.P.A.

DOMESTIC DOMESTIQUE

GUARANTEE/GARANTIE:
Diphacinone.....0.005%

READ THE LABEL BEFORE USING
(See back panel for additional precautionary statements)
KEEP OUT OF REACH OF CHILDREN

LIRE L'ETIQUETTE AVANT USAGE
(Voir a l'arriere pour precautions supplementaires)
NE PAS METTRE A LA PORTEE DES ENFANTS

NET CONTENTS: 1 Prebaited Mouse Bait Station with DITRAC BLOX
(28 g) POIDS NET: 1 Station pre-appatee pour souris avec DITRAC
BLOX (28 g)

BELL LABORATORIES, INC.
MADISON, WI 53704 U.S.A.

DIRECTIONS FOR USE

USE LIMITATIONS: Use DITRAC Mouse Bait Station to control the
house mouse(Mus musculus) in and around dwellings.

SELECTION OF TREATMENT AREAS: Determine areas where mice will
most likely find and consume the bait. Generally, these are

along walls, by gnawed openings, beside burrows, or in corners and concealed places.

APPLICATION DIRECTIONS: HOUSE MICE: Place bait station at intervals of 2.5 to 3 meters along walls or in corners where mice live or travel. The bait station can be secured to the floor with a screw or double-sided tape. When bait is consumed, replace with one DITRAC BLOX.

TO REFILL:(See illustrations) 1. Fully insert 2-pronged key in holes on top of DITRAC Mouse Bait Station. 2. Bend key away from station. 3. Gently lift up. 4. Place one DITRAC BLOX in the triangular area next to the hinge. 5. Close lid until it snaps securely. Repeat steps for easy rebaiting. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment.

ENVIRONMENTAL HAZARD: Keep out of lakes, streams, or ponds.

DISPOSAL: Wrap and dispose of in household garbage.

PRECAUTIONS: Hazardous to humans and domestic animals.

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of blood and cause bleeding. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: If swallowed, call a physician or Poison Control Center at once.

TOXICOLOGICAL INFORMATION: If ingested, Vitamin K, in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

Net Contents: 28 g (1 oz.)

BELL LABORATORIES, INC.
MADISON, WI 53704 U.S.A.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))