

05-JUL-2004 Notification received of a label change. See below for this change.

2000/2005

Notification Change

~~CLEAN CROP®~~

MECOPROP

HERBICIDE

COMMERCIAL

CAUTION

POISON

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

GUARANTEE:

MECOPROP.....150 g/L
(d'isomers equivalent present as potassium salt)

REGISTRATION NO. 10915
PEST CONTROL PRODUCTS ACT

Notification Change

~~UNITED AGRI PRODUCTS
RR #2
Dorchester, Ontario
N0L 1G5~~

UNITED AGRI PRODUCTS CANADA INC.
789 DONNYBROOK DRIVE □
DORCHESTER ON N0L 1G5 □

NET: 10 L

AC 00086-2-00

DIRECTIONS FOR USE

Liquid Weed Killer for Turf, Golf Greens and Fairways as well as cereals. DO NOT APPLY BY AIR.

WEATHER - Apply in warm weather under good growing conditions. Avoid spraying in very hot weather or in drought conditions.

MIXING - To ensure even mixing half fill the sprayer tank with clean water, add the required amount of MECOPROP and stir thoroughly. Add the remainder of the water and mix before spraying.

SPRAY VOLUME - Equip the machine with nozzle tips to apply 300 to 400 litres of water per hectare using a pressure not exceeding 275 kPa. Before use calibrate the machine to ensure that the correct

application rate is obtained.

On small areas MECOPROP may be applied at the recommended rates in 10 L/100m². It is recommended that the grass is not cut for 3 or 4 days before or after spraying. Do not water for 24 hours after treatment. Treatment may be repeated if necessary. MECOPROP may act more slowly than some weedkillers and up to three weeks may elapse before effect on the weeds becomes apparent.

WEED CONTROL IN TURF, GOLF GREENS & FAIRWAYS

MECOPROP may be used on Red fescue, Blue grasses, Bent grasses and Rye grasses when applied according to the directions below.

APPLICATION RATES

Seedling grasses: 60 mL/100m²; 5.5 L/ha (should be at least 2.5 cm high at spraying).

Established Turf: 90 mL/100m²; 8.5 L/ha

FOR FAIRWAYS AND LAWNS

For mixed weed populations, add 3L of MCPA Amine 500 to 6.6L of MECOPROP.

WEEDS CONTROLLED

SUSCEPTIBLE:

Seedlings: Use 60 mL/100m²; 5.5 L/ha

Established Plants: Use 90 mL/100 m²; 8.5 L/ha

Spray when the plants are growing vigorously.

Common chickweed (*Stellaria media*)

Mouse-ear chickweed (*Cerastium vulgatum*)

Plantain (*Plantago* sp.)

Clover (*trifolium* spp.)

MODERATELY SUSCEPTIBLE:

Use 90 mL/100 m²; 8.5 L/ha

Buttercup (*Ranunculus* spp). Spray before flowering.

Creeping Charlie (*Glechoma hederacea*). In early June or September.

Black Medick (*Medicago lupulina*). When young and growing fast.

Where Dandelion is present MECOPROP + 2,4-D is recommended.

WHEAT, BARLEY AND OATS: MECOPROP should be applied under good growing conditions from the time the crop is at the 3-leaf expanded stage until the early flag leaf (shotblade) stage: grain is about 10-15 cm high. DO NOT SPRAY ON GRAIN UNDERSEEDED WITH LEGUMES.

RATES OF APPLICATION ON CEREALS FOR CONTROL OF:

Black Medick
Canada Thistle (top growth only)
Chickweed
Cleavers
Clover (volunteer)

Corn Spurry
Lamb's Quarters
Plantain
Stitchwort
Wild Mustard

FOR SEEDLING WEEDS USE5.5L/ha

FOR MATURE WEEDS USE7.L/ha in 200-300 L of water/ha

For best results apply when the weeds are in the 2-4 leaf stage.

CAUTION: Do not graze the treated crops or cut for hay; sufficient data are not available to support such use.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Store with container tightly closed, and away from seeds, fertilizers, plant foodstuffs and greenhouses. Protect from freezing. Avoid spray drift on to susceptible crops. Wash sprayer thoroughly before and after use. Avoid contamination of ponds, streams, rivers and water sources. After use wash hands and skin contaminated by spray or concentrate.

FIRST AID

If swallowed induce vomiting. Get medical attention or contact a poison control centre. If in eyes flush with water for 15 minutes. Get medical attention if irritation persists. If on skin wash with soap and water. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

DISPOSAL

1. Triple or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

CLEAN CROP®

MECOPROP

HERBICIDE COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

GUARANTEE:

MECOPROP, d'isomers equivalent present as potassium salt.....150 g/L

REGISTRATION NO. 10915
PEST CONTROL PRODUCTS ACT

CAUTION

POISON

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Store with container tightly closed, and away from seeds, fertilizers, plant foodstuffs and greenhouses. Protect from freezing. Avoid spray drift on to susceptible crops. Wash sprayer thoroughly before and after use. Avoid contamination of ponds, streams, rivers and water sources. After use wash hands and skin contaminated by spray or concentrate.

FIRST AID

If swallowed induce vomiting. Get medical attention or contact a poison control centre. If in eyes flush with water for 15 minutes. Get medical attention if irritation persists. If on skin wash with soap and water. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

DISPOSAL

1. Triple or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.

4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

NET:10 L

**United Agri Products,
RR #2,
Dorchester, Ontario N0L 1G5**

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))