

93.03.29

2 kg
NET WEIGHT

WILBUR-ELLIS
IDEAS TO GROW WITH(R)

PARATHION 15W

INSECTICIDE WETTABLE POWDER

RESTRICTED

DANGER POISON

GUARANTEE: PARATHION 15%

REGISTRATION NO. 14777 PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

WILBUR-ELLIS COMPANY OF CANADA LIMITED
555 Southgate Drive,
Unit 101,
Guelph, Ontario N1H 6J3

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Poisonous if swallowed, inhaled or absorbed through the skin. Rapidly absorbed through the skin. Repeated exposure may, without symptoms progressively increase susceptibility to parathion poisoning. If you suspect excessive exposure, see a doctor. Avoid use on excessively hot days or in confined areas. Do not breathe powder while opening or handling the container. Wear a respirator such as American Optical Respirator 5058 or Willson Agritox Respirator while handling and spraying or entering newly sprayed areas. Do not breathe spray mist. Wear protective clothing. Change contaminated clothing and wash thoroughly before re-use. Never handle the material with bare hands; use natural rubber gloves. Do not use synthetic rubber, leather or cotton gloves. Wash hands, face and arms thoroughly after handling parathion and before eating or smoking. Take a complete bath or shower at the end of the day. Always work to the windward side when handling or

applying treatments. Keep unprotected persons out of operating areas or vicinity or where there may be a danger of drift. Do not re-enter treated areas within 48 hours after treatment. Wear protective clothing including gloves when entering treated areas or handling treated foliage within 14 days of treatment. Do not store product near food, feed, fertilizers, seed, and like commodities. Do not use in dwellings or home gardens. Do not use on crops during pollinating periods.

SYMPTOMS OF POISONING: Headache, blurred vision, weakness, nausea, abdominal cramps, chest discomfort, diarrhea, salivation, sweating, vomiting.

FIRST AID: If swallowed, induce vomiting by touching the back of the throat with a finger or blunt object; repeat until vomit fluid is clear. Rush patient to nearest hospital taking this container with you. In case of skin contact, remove contaminated clothing and wash skin with soap and water. If in eyes, flush with plenty of water for 15 minutes and get medical attention at once.

TOXICOLOGICAL INFORMATION: Never give morphine or other respiratory depressants. Give atropine by injection or by tablet 0.65 - 2.2 milligrams every hour until pupils dilate. Patient should be fully atropinized to ensure adequate results. Clear chest by postural drainage. Give artificial respiration or oxygen if patient becomes comatose. Observe patient continuously for 24 - 48 hours. Warn patient against re-exposure, including contaminated clothing. Do not permit exposure to parathion or other phosphate insecticides until cholinesterase is regenerated as determined by blood tests.

STORAGE: Store in a cool dry place away from food or feed.

COMPATIBILITY: PARATHION 15W appears to be compatible and safe with all the commonly used insecticides and fungicides. It should NOT be used with Bordeaux Mixture, Captan, Morestan, Morocide, Perthane, Phygon or summer oils.

DISPOSAL:

1. Empty bag.
2. Follow provincial instructions for any required additional cleaning of the bag prior to its disposal.
3. Make the empty bag unsuitable for further use.
4. Dispose of the bag in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills contact the regional office of Environment Canada.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

RESTRICTED USES

NOTICE TO USER: This control product is to be used only in accordance with the Directions on this label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NATURE OF RESTRICTION: This product is to be stored apart from lodgings for humans, animal quarters, and normal work areas, to avoid inadvertent exposure.

APPLE, APRICOT, CHERRY, PEACH, PEAR, PLUM, PRUNE: Black Cherry Aphid, Cherry Fruit Fly, Codling Moth, Cottony Peach Scale, Eye-spotted Budmoth, Mealybug, Oriental Fruit Moth, Plant Bug, Plum Curculio, Red-banded Leafroller, San Jose, Lecanium and Oystershell Scale, Short-hole Borer, Woolly Aphid.

METHOD/RATE: Use 6 - 8.4 kilograms per hectare (kg/ha). Begin treatment when pests are first noticed. Repeat at 7 to 10 day intervals as needed to control. Avoid using higher rates than recommended. Avoid application during very hot, still weather. Avoid contact with foliage for 14 days after spraying. May cause injury to apple varieties such as McIntosh, Snow, Golden Delicious, R.I. Greening, Duchess, Jonathon, Melba and Cortland under extremely poor drying conditions. Do not apply later than 14 days before harvest.

GRAPE, STRAWBERRY: Grape Mealybug, Strawberry Leafroller.

METHOD/RATE: Use 2 kg/1000 litres (L) water. For time of application follow provincial calendar. Avoid contact with foliage for 14 days after spraying. Do not apply later than 14 days before harvest.

BEET (red), CELERY, LETTUCE (leaf, head), ONION, POTATO, SPINACH, TOMATO: Aphids, Armyworm, Cabbage Looper, Flea Beetle, Grasshopper, Leafhopper, Leafminer, Leafroller, Onion Maggot (Fly), Thrip, Green Stink Bug, Spider Mite, Tomato Caterpillar, Webworm, Whitefly.

METHOD/RATE: Use 1.5 - 2.25 kg/ha. Treat infestations as necessary to control, using water-base spray. Repeat if necessary. For lettuce, spray at 6-day intervals from transplanting or emergence until 4 weeks before harvest. Do not apply later than the following number of days to harvest: 5 days - head lettuce, potatoes; 7 days - spinach; 10 days - tomatoes;

15 days - red beets, celery, onions; and 21 days - leaf lettuce.

CARROT: Carrot Rust Fly.

METHOD/RATE: Apply 2.25 kg/ha. Treat in mid-August, then at weekly intervals. Do not apply within 15 days of harvest.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))