

17-AUG-1998

((LABEL))

CB®

PURGE®

CB-80

Insect Killer

COMMERCIAL

KILLS: Flies, Mosquitoes, Wasps, Gnats, Cockroaches, Ants, Crickets, Spiders, Silverfish, Small Flying Moths, Indianmeal Moths, Angoumois Grain Moths, Carpet Beetles, Confused Flour Beetles, Sawtoothed Grain Beetles, Granary Weevils, Rice Weevils, Red Flour Beetles, Grain Mites, Bedbugs and Fleas.

FOR USE IN Food Handling Establishments, Commercial or Institutional Kitchens, Dining Areas and Pantries of Restaurants, Supermarkets, Food Plants, Food Warehouses, Bottling Plants, Hospitals, Hotels, Motels, Homes, Stores, Dairies, Stables, Nursing Homes, Day-care Centers, Egg and Milk Handling Areas of Dairies and Chicken Houses.

Concentrated power, fast knockdown and kill

GUARANTEE:

Pyrethrins.....0.50%
Piperonyl Butoxide, technical....4.00%

READ THE LABEL AND ATTACHED LEAFLET BEFORE USING

REGISTRATION NO. 22770
PEST CONTROL PRODUCTS ACT

Product of U.S.A.

CAUTION EXPLOSIVE

NET CONTENTS 851 g

Manufactured by
WATERBURY COMPANIES, INC.
P.O. Box 1812 Waterbury, CT 06722 USA

DISPOSAL: When container is empty, press valve to release all remaining pressure. Dispose of empty container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the provincial regulatory agency or the manufacturer.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Do not use in presence of open flame or spark. Contents under pressure. Container may explode if heated. Do not expose to temperature above 50EC. Do not puncture or incinerate.

Harmful if swallowed. Avoid breathing of vapor. Avoid contact with skin and eyes. In case of contact, immediately flush eyes with plenty of water. Wash skin with warm water and soap. Get medical attention if irritation persists. Do not contaminate food or foodstuffs. Do not remain in treated areas. Vacate premises for at least 15 minutes after spraying. Ventilate the area after treatment is completed and before re-entering. Do not use in rooms where infants, ill or aged are confined. Remove pets and birds. Cover fish aquariums before spraying.

Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before use. When using the product in these areas, apply when the facility is not in operation.

Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

FIRST AID: In case of contact with eyes or skin, flush with water. If irritation persists, seek medical attention. If accidentally swallowed, seek medical attention at once.

NOTICE TO USER: This Control Product is to be used only in accordance with the directions on the label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

((LEAFLET))

CB®

PURGE®

CB-80

Insect Killer

COMMERCIAL

KILLS: Flies, Mosquitoes, Wasps, Gnats, Cockroaches, Ants, Crickets, Spiders, Silverfish, Small Flying Moths, Indianmeal Moths, Angoumois Grain Moths, Carpet Beetles, Confused Flour Beetles, Sawtoothed Grain Beetles, Granary Weevils, Rice Weevils, Red Flour Beetles, Grain Mites, Bedbugs and Fleas.

FOR USE IN Food Handling Establishments, Commercial or Institutional Kitchens, Dining Areas and Pantries of Restaurants, Supermarkets, Food Plants, Food Warehouses, Bottling Plants, Hospitals, Hotels, Motels, Homes, Stores, Dairies, Stables, Nursing Homes, Day-care Centers, Egg and Milk Handling Areas of Dairies and Chicken Houses.

Concentrated power, fast knockdown and kill

GUARANTEE:

Pyrethrins.....0.50%
Piperonyl Butoxide, technical....4.00%

READ THE LABEL LEAFLET BEFORE USING

REGISTRATION NO. 22770
PEST CONTROL PRODUCTS ACT

Product of U.S.A.

CAUTION EXPLOSIVE

NET CONTENTS 851 g

Manufactured by
WATERBURY COMPANIES, INC.
P.O. Box 1812 Waterbury, CT 06722 USA

DISPOSAL: When container is empty, press valve to release all remaining pressure. Dispose of empty container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the provincial regulatory agency or the manufacturer.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Do not use in presence of open flame or spark. Contents under pressure. Container may explode if heated. Do not expose to temperature above 50°C. Do not puncture or incinerate.

Harmful if swallowed. Avoid breathing of vapor. Avoid contact with skin and eyes. In case of contact, immediately flush eyes with plenty of water. Wash skin with warm water and soap. Get medical attention if irritation persists. Do not contaminate food or foodstuffs. Do not remain in treated areas. Vacate premises for at least 15 minutes after spraying. Ventilate the area after treatment is completed and before re-entering. Do not use in rooms where infants, ill or aged are confined. Remove pets and birds. Cover fish aquariums before spraying.

Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before use. When using the product in these areas, apply when the facility is not in operation.

Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

FIRST AID: In case of contact with eyes or skin, flush with water. If irritation persists, seek medical attention. If accidentally swallowed, seek medical attention at once.

NOTICE TO USER: This Control Product is to be used only in accordance with the directions on the label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

DIRECTIONS FOR USE

FLYING INSECTS: FLIES, MOSQUITOES, SMALL FLYING MOTHS, WASPS, GNATS: Calculate m³ of space to be treated. Close all doors and windows. Spray in sweeping motion to all parts of the room. Apply for 2-3 seconds per 30 m³ of space. Sweep up and destroy fallen insects. Repeat treatment as necessary, but not more than once per 24 hour period (day).

CRAWLING INSECTS: COCKROACHES, ANTS, SPIDERS, SILVERFISH:
Calculate m² of floor area to be treated. Open all cabinets, storage areas, closets. Spray directly on insects or in cracks and crevices where insects hide.

FOR COCKROACHES: For best results, spray directly on cockroaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind stoves and refrigerators, food storage areas, and wherever these pests are suspected of hiding. Also, spray around door and window frames. Repeat as necessary, but not more than once per 24 hour period (day).

CRACK & CREVICE APPLICATION: Food areas and/or food handling establishments - Spray in cracks and crevices, using the crack and crevice nozzle provided. A crack and crevice application may be done in food areas and/or food handling establishments while these facilities are not in operation.

FOR SPIDERS AND SILVERFISH: For control of these pests, frequent and repeated spraying of baseboards, windows, doorframes, bookcases and storage areas should be made. For adequate control, spray directly on pests.

FOR CRICKETS: Spray thoroughly and frequently behind stoves, refrigerators, under cabinets, in food storage areas, and especially in closets, basements, and other damp areas. For best results, spray directly on insects.

FOR ANTS: Spray ant trails, nests and points of entry. For initial clean-up, apply for 10-15 seconds per 9.5 m² of floor area. Then periodically spray as a space aerosol 3-5 seconds per 30 m³ of space.

FOR INDIANMEAL MOTHS AND ANGOUMOIS GRAIN MOTHS: Close doors and windows. Direct spray into all parts of room, especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 30 m³ of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.

FOR CARPET BEETLES: Spray directly on beetles, baseboards, floors, and shelves. On carpets, use as a spot treatment only. Test for color fastness first. Repeat treatment as necessary, but not more than once per 24 hour period (day).

FOR CONFUSED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, GRANARY WEEVILS, RICE WEEVILS, RED FLOUR BEETLES, GRAIN MITES: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to

congregate. Spray 30 seconds for each 9.5 m² of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary, but not more than once per 24 hour period (day).

FOR BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary, but not more than once per 24 hour period (day).

FOR FLEAS: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary, but not more than once per 24 hour period (day). Use only a recommended pet spray on pets.

Note, when using this product in an enclosed area, such as milk room, barn or stable, remove animals before spraying. Spray only when temperatures are below 32°C. Do not contaminate feed or water troughs.

DAIRY FARM USE: Milking Parlor and Milk Room - Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds for 30 m³ of space. Do not remain in treated areas. Ventilate after treatment. Use only when milkhouse is not in operation. After spraying, wash all surfaces that may contact milk and rinse thoroughly with potable water before resuming operation. Cover milking utensils and milk to prevent contamination from spray or dead and falling insects.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 2-3 seconds per 30 m³ of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat application as necessary, but not more than once per 24 hour period (day).

BEEF CATTLE USE: Thoroughly spray entire animal from approx. 0.6 m distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Do not spray directly on poultry. Spray at the rate of 2-3 seconds per 30 m³ of space. Do not remain in treated areas. After 15 minutes, ventilate the area. Treat daily or as necessary.

SWINE USE: For animal application, direct spray over backs of hogs and spray for 5 seconds per hog. Do not spray directly towards the animal's face or eyes. Repeat application daily or as necessary. Do not use more than once a day.

HOG OPERATIONS: In hog houses, close all doors and windows. Spray at the rate of 2-3 seconds per 30 m³ of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary, but not more than once per 24 hour period (day).

HORSE BARNS OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 2-3 seconds per 30 m³ of space. Do not remain in the treated area. After 15 minutes, ventilate the area. Repeat the application as necessary, but not more than once per 24 hour period (day).

HORSE OR PONEY USE: For animal application, lightly mist over backs of horses and ponies from a distance of 0.6 m. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary. Do not cover animals after treatment.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))