

93.05.21

10 L

NET CONTENTS

MULTIVAP IV

BARN AND LIVESTOCK INSECTICIDE

COMMERCIAL

READ LABEL BEFORE USING

GUARANTEE (W/W)

0.1 % Pyrethrins

1.0 % Technical Piperonyl Butoxide

To protect livestock from horn flies, stable flies, house flies, mosquitoes, gnats and horseflies.

ATTENTION / CAUTION

POISON

KEEP OUT OF REACH OF CHILDREN

20424.01 REGISTRATION NUMBER

PEST CONTROL PRODUCTS ACT

MULTI-VET LTEE/LTD.

JOLIETTE, QUEBEC

J6E 3Z1

MULTIVAP IV

MULTIVAP IV INSECTICIDE: Premium fogging insecticide for use as a spray for the control of gnats, house flies, stable flies and mosquitoes in farm buildings. Also for use as a livestock spray for horn flies, mosquitoes, house flies, stable flies and horseflies.

FOGGING IN FARM BUILDINGS SUCH AS DAIRY BARNS, MILK ROOMS, HORSE BARNS, PIGGERIES, POULTRY HOUSES, SHELTER SHEDS, AND OTHER TYPES OF LIVESTOCK BUILDINGS.

Reduce air movement in the area to be treated by closing doors, windows and ventilators. Turn off all possible sources of ignition in area to be treated. Remove all animals from building before using as a space spray. Apply as a space spray using an electric fogger or similar mist applicator, at the rate of 200 mL of insecticide per 100 m³ of space. Spray upwards to fill area with a fine mist. Keep area closed 15 minutes before ventilating. Direct spray upwards to all parts of room especially windows and other light sources and to areas where insects are present. The insects must be in contact with Multivap IV to be killed. Ventilate treated areas thoroughly for at least 30 minutes before animal reentry. Sweep up dead insects. Repeat as necessary with additional spraying of breeding areas. Spray only when temperature is under 32°C.

LIVESTOCK SPRAY FOR THE CONTROL OF FLIES, AND MOSQUITOES ON BEEF CATTLE, DAIRY CATTLE, HOGS, AND MULES.

Apply 30-60 mL per animal, using a sprayer which produces a fine mist. Cover entire animal lightly, without making hair wet. Do not spray directly on udder. Do not spray animal's eyes or mouth. Repeat once or twice daily as required. Treat dairy cattle at least 20 minutes before milking. Wash udder before milking. Spray from a distance of at least one metre.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Not for use on humans. Harmful if swallowed. Avoid contact with skin, eyes and clothing and avoid inhalation of fog or spray. Product contains petroleum distillate; wear mask or respirator while fogging. Keep container closed and store away from heat sources. Prior to fogging, all possible sources of ignition should be extinguished in the area to be treated. Wash after use. Do not remain in treated areas. Ensure proper ventilation after sealed areas have been treated. Do not contaminate food, feed or water troughs. Any treated surface that may contact food or feed products should be washed thoroughly with potable water after spraying and before reuse. Cover milk and milking utensils to prevent contamination from spray and dead insects. Do not use in areas where beekeeping equipment or supplies are stored. Product toxic to fish. Do not contaminate any body of water.

DISPOSAL: Do not reuse empty container. Make empty container unsuitable for further use and dispose in accordance with provincial requirements. Follow provincial instructions for any required cleaning of container prior to disposal. For information on the disposal of unused or unwanted product and the clean-up of

spills, contact the Regional Office of Conservation and Protection, Environment Canada.

FIRST AID: If on skin, wash with soap and water, if in eyes, flush with plenty of water. If irritation persists, get prompt medical attention. This product contains petroleum distillate. If swallowed, do not induce vomiting, if patient is unconscious, give him air. Get prompt medical attention.

LIMITATION OF WARRANTY: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))