

(Warfarin Rat Killer Meal Throw Bag)

CO-OP ®

Rat Killer Meal

250 g

Domestic

REG. NO 11609 P.C.P. ACT

GUARANTEE: Warfarin 0.025%

Contains malathion at 0.0095% and calcium propionate at 0.3% as preservatives.

READ LABEL AND OUTER BOX BEFORE USING

KEEP OUT OF REACH OF CHILDREN

**INTERPROVINCIAL COOPERATIVE LTD.
P.O. BOX 1050, SASKATOON, SASK. S7K 3M9**

1-204-233-3461

RC 486-0307

**PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS,
AND DOMESTIC ANIMALS.**

FIRST AID: **If in eyes**, flush with water for 15 minutes. **If on skin**, wash with soap and water. **If swallowed**, give patient one or two glasses of water and induce vomiting by touching back of the throat with a finger or blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or Poison Control Centre. If pet or livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention.

DISPOSAL: Wrap container and discard in household garbage. Do not re-use empty container.

RODENT DISPOSAL: Dispose of dead rodents in garbage or by burying.

(Carton label)

CO-OP®

Net Contents 500 g

**WARFARIN
RAT KILLER
RODENTICIDE MEAL**

DOMESTIC

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

GUARANTEE: WARFARIN 0.025%
Contains malathion at 0.0095% and calcium propionate at 0.3%
as preservatives.

REGISTRATION NO. 11609
PEST CONTROL PRODUCTS ACT

INTERPROVINCIAL COOPERATIVE LIMITED, P.O. BOX 1050, SASKATOON,
SASKATCHEWAN S7K 3M9

1-204-233-3461

RC 459-0307

USE LIMITATIONS

Bait **MUST** either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. **DO NOT** place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

DIRECTIONS: (indoors and against the outside walls of buildings):

RATS: Place 250 to 500 g of bait at intervals of 4.5 to 9 meters per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

MICE: Place 15 to 30 g of bait at intervals of 3 to 4 meters per placement. Larger placements may be needed at points of high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK.

May be harmful or fatal if swallowed or absorbed through the skin. Do not open pre-measured place packs. Rubber gloves must be worn when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.** Keep out of lakes, streams or ponds.

FIRST AID: **If in eyes**, flush with water for 15 minutes. **If on skin**, wash with soap and water. **If swallowed**, give patient one or two glasses of water and induce vomiting by touching back of the throat with a finger or blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. If pet or livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: This product contains an anti-coagulant chemical which, if taken accidentally by humans, domestic animals or pets, may reduce the clotting ability of the blood and haemorrhage may result. Vitamin K₁ in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case, measured by establishing prolonged prothrombin times (P.T.), will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE:

Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

DISPOSAL: Wrap container and discard in household garbage. Do not re-use empty container.

RODENT DISPOSAL: Dispose of dead rodents in garbage or by burying.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

® Registered trademark of TMC Distributing Ltd. Used under license.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.