

99/2004

SANITROL-MB

DISINFECTANT-CLEANER SANITIZER-FUNGICIDE  
DEODORIZER

WITH ORGANIC SOIL TOLERANCE FOR HOSPITALS, INSTITUTIONAL,  
INDUSTRIAL, SCHOOL, DAIRY AND OTHER FARM USE

KEEP OUT OF REACH OF CHILDREN

FOR INDUSTRIAL INSTITUTIONAL & FOOD PLANT USE

WARNING

CAUTION

CORROSIVE

READ THE LABEL BEFORE USING

REGISTRATION NO. 23573

PEST CONTROL PRODUCTS ACT

DIN NO. 02073021

AVOID CONTAMINATION OF FOOD

GUARANTEE:

n-Alkyl (40% C12, 50% C14, 10% C16)  
dimethylbenzyl ammonium chloride.....3.000%  
Octyl decyl dimethyl ammonium chloride.....2.250%  
Dioctyl dimethyl ammonium chloride.....0.900%  
Didecyl dimethyl ammonium chloride.....1.350%

NET CONTENTS

litres      20 litres      210 litres

MULTI-BLEND LTD.  
2720 Slough Street

Mississauga, Ontario  
L4T 1G3

SANITIZING OF FOOD PROCESSING EQUIPMENT AND OTHER HARD SURFACES IN FOOD CONTACT LOCATIONS. For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard nonporous surfaces. No potable rinse is required.

Wash and rinse all articles thoroughly, then apply a solution of 2 mL SANITROL-MB in 1 litre of water (150 ppm active). Surfaces should remain wet for at least 1 minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanical application, use solution may not be reused for sanitizing applications. Apply to sink tops, counter tops, refrigerated storage and display equipment and other stationary hard surfaces by cloth, brush, or mechanical spray device. No potable water rinse is required. Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 2 mL/L solution of SANITROL-MB. No potable rinse required.

PRECAUTIONARY STATEMENTS: KEEP OUT OF REACH OF CHILDREN.

Danger: Corrosive to eyes. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST-AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg white gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

TOXICOLOGICAL INFORMATION: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence

under the Pest Control Products Act to use a control product under unsafe conditions.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to treatment site.
2. Follow provincial instructions or any required additional cleaning of the container prior to its disposal or reconditioning.
3. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted product and the cleanup of spills contact the regional office of Environment Canada.
5. For small spills. Recover free liquid. Soak up residue with solids absorbent. DO NOT FLUSH INTO SURFACE STREAMS. INFORM ENVIRONMENT CANADA AS IN #4.

DIRECTIONS FOR USE: Apply SANITROL-MB with a cloth or mop. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the used solution becomes very dirty.

DISINFECTION IN HOSPITALS, NURSING HOMES AND OTHER HEALTH CARE INSTITUTIONS: For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard nonporous surfaces. Add 6 mL SANITROL-MB to 1 litre water. Apply to previously cleaned hard surfaces. At this use level, SANITROL-MB is effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella choleraesuis* in the presence of 5% blood serum, when evaluated by the AOAC Use-Dilution Test.

DISINFECTION OF INSTITUTIONS, INDUSTRY AND SCHOOLS: Add 6.0 mL of SANITROL-MB to 1 litre of water. At 1.75 mL/litre of water use level, the fungicidal effectiveness of SANITROL-MB in the presence of 5% blood serum against Trichophyten mentagrophytes has been shown, utilizing the AOAC Fungicidal Test.

DISINFECTION OF BARBER TOOLS: Precleaned barber tools (such as combs, brushes, razors and scissors), can be disinfected by immersing in a 2 mL/L solution of SANITROL-MB.

DISINFECTION OF POULTRY EQUIPMENT, ANIMAL QUARTERS AND KENNELS: Poultry brooders, watering fountains, feeding equipment and other

animal quarters (such as stalls and kennel areas) can be disinfected after thoroughly cleaning by applying a solution of 6 mL SANITROL-MB to 1 litre of water. Small utensils should be immersed in this solution. Prior to disinfection, all poultry, other animals and their foods must be removed from the premises. This includes emptying all troughs, racks and other feeding and water appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals. After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact food or drinking water must be rinsed with potable water before reuse.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))